

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/063169 A1

(51) International Patent Classification⁷: C07D 235/16,
471/04, 487/04, 235/12, 491/04, 401/04, 403/04, 233/64,
233/90, 233/68, A61K 31/4184, 31/4164, 31/4178, A61P
35/00

Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome,
Chuo-ku, Osaka-shi, Osaka 541-8514 (JP).

(21) International Application Number:

PCT/JP2004/000157

(74) Agent: TAKASHIMA, Hajime; Fujimura Yamato Seimei
Bldg., 2-14, Fushimimachi 4-chome, Chuo-ku, Osaka-shi,
Osaka 541-0044 (JP).

(22) International Filing Date: 13 January 2004 (13.01.2004)

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

2003-900116 13 January 2003 (13.01.2003) AU
2003-905406 6 October 2003 (06.10.2003) AU

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): FUJII-
SAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7,
Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-
8514 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): URANO, Ya-
suharu [JP/JP]; c/o Fujisawa Pharmaceutical Co., Ltd.,
4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka
541-8514 (JP). SATOH, Shigeki [JP/JP]; c/o Fujisawa
Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome,
Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). ISHIBASHI,
Naoki [JP/JP]; c/o Fujisawa Pharmaceutical Co., Ltd.,
4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka
541-8514 (JP). KAMIJO, Kazunori [JP/JP]; c/o Fujisawa


Published:

— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 2004/063169 A1

(54) Title: HYDROXAMID ACID DERIVATIVES AS HISTONE DEACETYLASE (HDAC) INHIBITORS



(57) Abstract: A compound having the following formula (I): wherein R_i¹ is N-containing heterocyclic ring optionally substituted with one or more suitable substituent(s), R_i² is hydroxyamino, R_i³ is hydrogen or a suitable substituent, L_i¹ is -(CH₂ⁿ)_n (wherein n is an integer of 0 to 6) optionally substituted with one or more suitable substituent(s), wherein one or more methylene(s) may be replaced with suitable heteroatom(s), and L_i² is lower alkenylene, or a salt thereof. The compound is useful as a histone deacetylase inhibitor.

DESCRIPTION

HYDROXAMID ACID DERIVATIVES AS HISTONE DEACETYLASE (HDAC) INHIBITORS


TECHNICAL FIELD

The present invention relates to a compound useful as a
5 medicament, and to a pharmaceutical composition comprising the same.

BACKGROUND ART

Histone deacetylase (hereinafter also referred as HDAC) is
known to play an essential role in the transcriptional machinery
for regulating gene expression, induce histone hyperacetylation and
10 to affect the gene expression. Therefore, it is useful as a
therapeutic or prophylactic agent for diseases caused by abnormal
gene expression such as inflammatory disorders, diabetes, diabetic
complications, homozygous thalassemia, fibrosis, cirrhosis, acute
promyelocytic leukaemia (APL), organ transplant rejections,
15 autoimmune diseases, protozoal infections, tumors, etc.

WO 01/38322 discloses an inhibitor of histone deacetylase
represented by the following formula:



20 wherein

Cy is cycloalkyl, aryl, heteroaryl or heterocyclyl, each of which
is optionally substituted;


L¹ is -(CH₂)_m-W- wherein m is an integer of 0 to 4, and W is selected
from the group consisting of -C(O)NH-, -S(O)₂NH-, etc.;

25 Ar is optionally substituted arylene, which is optionally fused to
an aryl, heteroaryl ring, etc.;

Y¹ is a chemical bond or a straight- or branched-chain saturated
alkylene, wherein said alkylene is optionally substituted; and

Z is selected from the group consisting of anilinyl, pyridyl,
30 thiadiazolyl and -O-M wherein M is H or a pharmaceutically
acceptable cation.

WO 02/22577 discloses the following hydroxamate compound as
a deacetylase inhibitor:



wherein

R₁ is H, halo or a straight chain C₁-C₆ alkyl;

R₂ is selected from H, C₁-C₁₀ alkyl, C₄-C₉ cycloalkyl, C₄-C₉
 5 heterocycloalkyl, C₄-C₉ heterocycloalkylalkyl, cycloalkylalkyl,
 aryl, heteroaryl, etc.;

R₃ and R₄ are the same or different and independently H, C₁-C₆ alkyl,
 acyl or acylamino, or

R₃ and R₄ together with the carbon to which they are bound to
 10 represent C=O, C=S, etc.; or

R₂ together with the nitrogen to which it is bound and R₃ together
 with the carbon to which it is bound to form a C₄-C₉
 heterocycloalkyl, a heteroaryl, a polyheteroaryl, a non-aromatic
 polyheterocycle, or a mixed aryl and non-aryl polyheterocycle ring;

15 R₅ is selected from H, C₁-C₆ alkyl, etc.;

n, n₁, n₂ and n₃ are the same or different and independently
 selected from 0-6, when n₁ is 1-6, each carbon atom can be
 optionally and independently substituted with R₃ and/or R₄;

X and Y are the same or different and independently selected from H,
 20 halo, C₁-C₄ alkyl, etc.;

or a pharmaceutically acceptable salt thereof.

SUMMARY OF THE INVENTION

The present invention relates to a novel compound useful as
 a medicament, and to a pharmaceutical composition comprising the
 25 same.

More particularly, the present invention relates to a
 compound having a potent inhibitory effect on the activity of
 histone deacetylase.

The inventors of the present invention have also found that
 30 histone deacetylase inhibitors, such as a compound of the formula
 (I) (hereinafter compound [I]), have a potent immunosuppressive
 effect and potent antitumor effect. Therefore, a histone

deacetylase inhibitors such as compound [I] is useful as an active ingredient for an immunosuppressant and an antitumor agent, and useful as an active ingredient for a therapeutic or prophylactic agent for diseases such as inflammatory disorders, diabetes, 5 diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), organ transplant rejections, autoimmune diseases, protozoal infections, tumors, etc.

Accordingly, one object of the present invention is to provide a compound having biological activities for treating or 10 preventing the diseases as stated above.


A further object of the present invention is to provide a pharmaceutical composition containing the compound [I] as an active ingredient.

A yet further object of the present invention is to provide 15 use of the histone deacetylase inhibitors, such as compound [I], for treating or preventing the diseases as stated above.

A yet further object of the present invention is to provide a commercial package comprising the pharmaceutical composition containing the compound [I] and a written matter associated 20 therewith, the written matter stating that the pharmaceutical composition may or should be used for treating or preventing the diseases as stated above.

Thus, the present invention provides

[1] A compound having the following formula (I):



25 wherein

R^1 is N-containing heterocyclic ring optionally substituted with one or more suitable substituent(s),

R^2 is hydroxyamino,


R^3 is hydrogen or a suitable substituent,

30 L^1 is $-(CH_2)_n-$ (wherein n is an integer of 0 to 6) optionally substituted with one or more suitable substituent(s),

wherein one or more methylene(s) may be replaced with suitable heteroatom(s), and
 L^2 is lower alkenylene,
 or a salt thereof.

5

[2] The compound of the above-mentioned [1], wherein
 R^1 is N-containing heterocyclic ring represented by the following
 formula:



10

wherein

R^4 is hydrogen or a group selected from the group
 consisting of

15


- (1) lower alkyl optionally substituted with di(lower)alkylamino or hydroxy,
- (2) lower alkoxy,
- (3) aryl optionally substituted with the substituent
 selected from the group consisting of halogen, lower
 alkanoyl, lower alkylsulfonyl, lower alkoxy and
 di(lower)alkylamino,
- (4) lower alkanoyl,
- (5) lower alkoxy-carbonyl,
- (6) arylcarbonyl,

25

- (7) aryl(lower)alkoxy,
(8) amino optionally mono- or di-substituted with
substituent(s) selected from the group consisting of
lower alkyl, lower alkanoyl and cycloalkyl,
5 (9) halo(lower)alkyl,
(10) aryloxy,
(11) aryl(lower)alkyl optionally substituted with
hydroxy,
(12) carboxyl,
10 (13) nitro,
(14) cyano,
(15) halogen,
(16) heteroaryl,
(17) non-aromatic heterocycle optionally substituted
15 with lower alkyl,
(18) hydroxy,
(19) (lower)alkylsulfonylcarbamoyl and
(20) non-aromatic heterocycle carbonyl,
R⁵ is hydrogen or a group selected from the group consisting
20 of lower alkyl and aryl(lower)alkyl, and
R⁶, R⁷ and R⁸ are each hydrogen or lower alkyl,
R⁹ is hydrogen or a group selected from the group consisting
of
(1) lower alkyl optionally substituted with
25 di(lower)alkylamino,
(2) aryl optionally substituted with lower alkoxy,
(3) (lower)alkoxy-carbonyl,
(4) cyano,
(5) carbamoyl optionally mono- or di-substituted with
30 (lower)alkyl,
(6) halogen,
(7) (lower)alkyl-carbonyl,
(8) arylcarbonyl and
(9) cyclo(lower)alkyl,
35 R¹⁰ is hydrogen or a group selected from the group consisting
of

- (1) (lower)alkylcarbamoyl,
 (2) di(lower)alkylcarbamoyl,
 (3) aryl optionally substituted with halogen,
 (4) (lower)alkoxy-carbonyl,
 5 (5) carboxy,
 (6) non-aromatic heterocycle carbonyl,
 (7) halogen,
 (8) (lower)alkyl optionally substituted with hydroxy,
 (lower)alkoxy, non-aromatic heterocycle, aryl,
 10 di(lower)alkylamino or halogen and
 (9) adamantyl,
 R^{11} is hydrogen or aryl(lower)alkyl in which the aryl portion
 is substituted with lower alkoxy,
 R^{12} is hydrogen or a group selected from the group consisting
 15 of lower alkyl and aryl optionally substituted with
 halogen,
 R^{13} is hydrogen or a group selected from the group consisting
 of lower alkyl and aryl, and
 R^{14} is hydrogen or lower alkyl,
 20 R^2 is hydroxyamino,
 R^3 is hydrogen or lower alkoxy,
 L^1 is $-(CH_2)_n-$ (wherein n is 1 to 5) optionally substituted with
 one or more substituent(s) selected from lower alkyl(s) and
 aryl(lower)alkyl, and wherein one methylene may be replaced
 25 with an oxygen atom, and
 L^2 is vinylene,
 or a salt thereof.

- [3] The compound of the above-mentioned [2], wherein
 30 R^1 is N-containing condensed heterocyclic ring represented by the
 following formula:



wherein R^4 and R^5 are each as defined in the above-mentioned

[2].

[4] The compound of the above-mentioned [3], wherein

R⁴ and R⁵ are each hydrogen,

5 R² is hydroxyamino,

R³ is hydrogen,

L¹ is -CH₂-, and


L² is vinylene,

or a salt thereof.

10

[5] The compound of the above-mentioned [2], wherein

R¹ is N-containing heterocyclic ring represented by the following formula:



15

wherein R⁹, R¹⁰ and R¹¹ are each as defined in the above-mentioned [2].

[6] The compound of the above-mentioned [5], wherein

20 R⁹ is hydrogen or aryl optionally substituted with lower alkoxy,

R¹⁰ is hydrogen or aryl optionally substituted with halogen, and

R¹¹ is hydrogen,

R² is hydroxyamino,




R³ is hydrogen,

25 L¹ is -CH₂-, and


L² is vinylene,

or a salt thereof.

[7] A compound of the following formula.




OR



or a salt thereof.

5

[8] A compound having the following formula (I'):



wherein

R¹ is N-containing condensed heterocyclic ring optionally substituted with one or more suitable substituent(s),

10 R² is hydroxyamino,

R³ is hydrogen or a suitable substituent,

L¹ is -(CH₂)_n- (wherein n is an integer of 0 to 6) optionally substituted with one or more suitable substituent(s), wherein one or more methylene(s) may be replaced with suitable heteroatom(s),


5 and

L² is lower alkenylene,

or a salt thereof.

[9] The compound of the above-mentioned [8], wherein

10 R¹ is N-containing condensed heterocyclic ring represented by the following formula:



15 wherein

R⁴ is hydrogen or a group selected from the group consisting of

- (1) lower alkyl,
- (2) lower alkoxy,
- 20 (3) aryl optionally substituted with the substituent selected from the group consisting of halogen, lower alkanoyl, lower alkylsulfonyl, lower alkoxy and di(lower)alkylamino,
- (4) lower alkanoyl,
- 25 (5) lower alkoxy-carbonyl,
- (6) arylcarbonyl,
- (7) aryl(lower)alkoxy,
- (8) amino optionally mono- or di-substituted with

substituent(s) selected from the group consisting of
lower alkyl, lower alkanoyl and cycloalkyl,

(9) halo(lower)alkyl,

(10) aryloxy,

5 (11) aryl(lower)alkyl optionally substituted with
hydroxy,

(12) carboxyl,

(13) nitro,

(14) cyano,

10 (15) halogen,

(16) heteroaryl and

(17) non-aromatic heterocycle optionally substituted
with lower alkyl,

15 R^5 is hydrogen or a group selected from the group consisting
of lower alkyl and aryl(lower)alkyl, and

R^6 , R^7 and R^8 are each hydrogen or lower alkyl,

R^2 is hydroxyamino,

R^3 is hydrogen or lower alkoxy,


20 L^1 is $-(CH_2)_n-$ (wherein n is 1 or 2) optionally substituted with
one or more substituent(s) selected from lower alkyl(s) and
aryl(lower)alkyl, and wherein one methylene may be replaced
with an oxygen atom, and

L^2 is vinylene,

or a salt thereof.

25

[10] A compound having the following formula (I''):



wherein


R^1 is N-containing condensed heterocyclic ring optionally
substituted with one or more suitable substituent(s),

30 R^2 is hydroxyamino,

L^1 is $-(CH_2)_n-$ (wherein n is an integer of 0 to 6) optionally
substituted with one or more suitable substituent(s), and

L² is lower alkenylene,
or a salt thereof.

- [11] The compound of the above-mentioned [10], wherein
5 R¹ is N-containing condensed heterocyclic ring represented by the
following formula:



wherein

- 10 R⁴ is hydrogen or a group selected from the group consisting
of lower alkyl and aryl, and
R⁵ is hydrogen or a group selected from the group consisting
of lower alkyl and aryl(lower)alkyl,
R² is hydroxyamino,
15 L¹ is -(CH₂)_n- (wherein n is 1 or 2) optionally substituted with
aryl(lower)alkyl, and
L² is vinylene,
or a salt thereof.

- 20 Of the above-mentioned compounds, the compounds represented by
the general formulas (I') (compound [I']) and (I'') (compound
[I'']) are also encompassed in the scope of the compound
represented by the general formula (I). Hereinafter "compound [I]"
also encompasses "compound [I']" and "compound [I'"]".

25






The above-mentioned compounds and salts thereof can be
prepared by the processes as illustrated in the following reaction
schemes or by the methods disclosed in the Preparations and
Examples.

30





In the above and subsequent descriptions of the present
specification, suitable examples and illustration of the various
definitions which the present invention intends to include within
the scope thereof are explained in detail as follows.

In the following Processes, the compound [I-1] and [I-2] are encompassed in the scope of the compound [I], and the compound [II-A], [II-B], [II-C], [II-C'], [II-D], [II-E], [II-F], [II-G], [II-H], [II-I], [II-J], [II-K], [II-L], [II-M] and [II-N] are encompassed
5 in the scope of the compound [II].


Process A



Process B




Process C




Step 4
Same method as
Step 5 of Process A

Step 4'
Same method as
Step 5 of Process A






[II-C]




[II-C']

5


Process D




Process E







Step 1





Step 2







Process F







Process G




Process H




(H-0)




(H-1)



(H-2)







(H-3)




[II-H]

Process I



Process J




(H-2')

N-halogenated succinimide

CH₃CN

Step 1




(J-1)

1) TFA/DCM

2) WSCD, HOBT
and R^aO-NH₂


DMF

Step 2



[II-J]

Process K




(H-2'')

N-halogenated succinimide

CH₃CN

Step 1




(K-1)

1) TFA/DCM

2) WSCD, HOBT
and R^aO-NH₂




DMF

Step 2







[II-K]


Process L



Process M



Process N



wherein

R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴ and L¹ are as
 5 defined above,

Hal and Hal' are each halogen,

L^{1'} is L¹ in which one of the carbon atoms is substituted with R⁵
 (wherein R⁵ is as defined above),

R¹⁵ and R¹⁶ are each lower alkyl (e.g., alkyl having 1 to 6 carbon

atom(s) such as methyl, ethyl, propyl, butyl, tert-butyl, pentyl, hexyl, etc.), or R^{15} , R^{16} and the nitrogen atom to which they are attached may be together to form non-aromatic heterocycle in which one or more carbon(s) of said heterocycle is(are) optionally
5 replaced with one or more heteroatom(s) selected from oxygen, nitrogen and sulfur (e.g., piperidino, morpholino, etc.), and R^a is a hydroxy protecting group.

In the above-mentioned Processes A, B, C, D, E, F, G, H, I, J, K, L, M and N, each of the starting compounds can be prepared,
10 for example, according to the procedures as illustrated in the Preparations in the present specification or in a manner similar thereto.

For example, compounds (A-1), (A-2), (A-3) and (A-4) can be obtained by the procedures as illustrated in Preparations 1, 2, 3
15 and 4, respectively; compounds (B-1), (B-2) and (B-3) can be obtained by the procedures as illustrated in Preparations 6, 7 and 8, respectively; compound (C-1) and (C-1') can be obtained by the procedures as illustrated in Preparation 10; compound (C-2) and (C-3) can be obtained by the procedure as illustrated in Preparations
20 11 and 12, respectively; compound (C-2') and (C-3') can be obtained by the procedure as illustrated in Preparations 23 and 24, respectively; compound (D-1) and (D-2) can be obtained by the procedure as illustrated in Preparations 20 and 21, respectively; compound (E-1) can be obtained by the procedure as illustrated in
25 Preparation 35; compounds (F-1), (F-2), (F-3), (F-4) and (F-5) can be obtained by the procedures as illustrated in Preparations 127, 128, 129, 130 and 131, respectively; compounds (G-1) and (G-2) can be obtained by the procedures as illustrated in Preparations 195 and 196, respectively; compounds (H-1), (H-2) and (H-3) can be
30 obtained by the procedures as illustrated in Preparations 204, 206 and 201, respectively; compounds (I-1) and (I-2) can be obtained by the procedures as illustrated in Preparations 212 and 219, respectively; compound (J-1) can be obtained by the procedure as illustrated in Preparation 226; Compound (K-1) can be obtained by
35 the procedure as illustrated in Preparation 293; Compound (L-1) can be obtained by the procedure as illustrated in Preparation 294;


Compounds (M-1) and (M-2) can be obtained by the procedures illustrated in Preparations 101 and 102, respectively; and Compounds (N-1) and (N-2) can be obtained by the procedures illustrated in Preparations 233 and 249, respectively. The

5 compounds [II-A], [II-B], [II-C], [II-C'], [II-D], [II-E], [II-F], [II-G], [II-H], [II-I], [II-J], [II-K], [II-L], [II-M] and [II-N] can be obtained, for example, by the procedures as illustrated in Preparations 5, 9, 13, 25, 22, 39, 132, 197, 208, 222, 191, 282, 284, 103 and 189, respectively.



10

The compound [I] of the present invention is obtained from compound [II], for example, according to the following processes or methods disclosed in the Examples.

Process 1



15

Process 2

wherein R¹, R³, L¹, L² and R^a are as defined above.

Process 1

5 The compound [I-1] is obtained as an acid addition salt by deprotecting the hydroxy group of the compound [II] in the presence of an acid.

10 The acid includes hydrogen chloride solution (e.g. hydrogen chloride in solvent such as methanol, dioxane, ethyl acetate, diethyl ether, etc.), acetic acid, p-toluenesulfonic acid, boric acid, etc.

 Optionally, one or more suitable solvent(s) for the deprotection is(are) used. Such solvent includes methanol, ethanol, ethyl acetate, dioxane, diethyl ether, acetic acid, etc.

15 The temperature of the reaction is not critical, and the reaction is usually carried out under from cooling to heating.

 This Process is exemplified by Examples 1, 58, etc.

Process 2

20 The compound [I-2] is obtained as a free form by reacting the above-mentioned compound [I-1], which is an acid salt addition salt, with a base.

 Suitable base includes sodium hydroxide, potassium hydroxide, lithium hydroxide, aqueous sodium hydrogen carbonate solution, aqueous potassium hydrogen carbonate solution, aqueous sodium

hydroxide solution, etc.

Optionally, one or more suitable solvent(s) for the deprotection is(are) used for this reaction. Such solvent includes such as methanol, ethanol, ethyl acetate, dioxane, diethyl ether, acetic acid, etc.

The temperature of the reaction is not critical, and the reaction is usually carried out under from cooling to heating.

This Process is exemplified by Examples 116, etc.

Furthermore, the compound [I-2] can be converted to a suitable salt, which is also encompassed in the scope of the present invention, by a conventional method or the methods explained in the present specification (e.g., Examples 113, 118, 119, 120, 123, etc.).

The compound [I] may be a salt, which is also encompassed in the scope of the present invention. For example, when a basic group such as an amino group is present in a molecule, the salt is exemplified by an acid addition salt (e.g. salt with an inorganic acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, etc., salt with an organic acid such as methanesulfonic acid, benzenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid (e.g., [(1S,4R)-7,7-dimethyl-2-oxobicyclo[2.2.1]hept-1-yl]methanesulfonic acid or an enantiomer thereof, etc.), fumaric acid, maleic acid, mandelic acid, citric acid, salicylic acid, malonic acid, glutaric acid, succinic acid, etc.), etc., and when an acidic group such as carboxyl group is present, the salt is exemplified by a basic salt (e.g. salt with a metal such as lithium, sodium, potassium, calcium, magnesium, aluminium, etc., salt with an amino acid such as lysine, etc.), etc.

In addition, solvates of the compound [I] such as hydrate, ethanolate, etc., are also encompassed in the scope of the present invention.

When the compound [I] has stereoisomers, such isomers are also encompassed in the scope of the present invention.

In the above and subsequent descriptions of the present specification, suitable examples and illustration of the various definitions which the present invention intends to include within

the scope thereof are explained in detail as follows.

Each of the terms "halogen", "halo" and "Hal" includes fluorine, chlorine, bromine and iodine.

The term "heteroatom" includes nitrogen atom, oxygen atom and
5 sulfur atom.

The term "lower" used in the description is intended to mean 1 to 6 carbon atom(s) unless otherwise indicated.

Suitable "one or more" includes the number of 1 to 6, preferably 1 to 3.

10 Suitable "lower alkyl" includes straight or branched alkyl having 1 to 6 carbon atom(s) such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, tert-pentyl, neopentyl, hexyl, isohexyl, etc.

Suitable "cyclo(lower)alkyl" includes cycloalkyl having 3 to
15 6 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.

Suitable "lower alkoxy" includes straight or branched alkoxy having 1 to 6 carbon atom(s) such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy,
20 tert-pentyloxy, neopentyloxy, hexyloxy, isohexyloxy, etc.

Suitable "lower alkanoyl" includes formyl and alkanoyl in which the alkyl portion is straight or branched alkyl having 1 to 6 carbon atom(s) such as acetyl, ethylcarbonyl, propylcarbonyl, isopropylcarbonyl, butylcarbonyl, isobutylcarbonyl, sec-
25 butylcarbonyl, tert-butylcarbonyl, pentylcarbonyl, tert-pentylcarbonyl, neopentylcarbonyl, hexylcarbonyl, isohexylcarbonyl, etc.

Suitable "lower alkoxy-carbonyl" includes alkoxy-carbonyl in which the alkyl portion is straight or branched alkyl having 1 to 6
30 carbon atom(s) such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl, tert-pentyloxycarbonyl, neopentyloxycarbonyl, hexyloxycarbonyl, isohexyloxycarbonyl, etc.

35 Suitable "halo(lower)alkyl" includes lower alkyl substituted with 1 to 3 halogen atom(s) such as monochloromethyl,

dichloromethyl, trichloromethyl, monofluoromethyl, difluoromethyl, trifluoromethyl, monobromomethyl, dibromomethyl, tribromomethyl, monochloroethyl, dichloroethyl, trichloroethyl, monofluoroethyl, difluoroethyl, trifluoroethyl, etc.

5 Suitable "lower alkenylene" includes straight or branched alkylene having 1 to 6 carbon atom(s) such as vinylene, 1-methylvinylene, 2-methylvinylene, 1-propenylene, 2-propenylene, 2-methyl-1-propenylene, 2-methyl-2-propenylene, 1-butenylene, 2-butenylene, 3-butenylene, 1-pentenylene, 2-pentenylene, 3-pentenylene, 4-pentenylene, 1-hexenylene, 2-hexenylene, 3-hexenylene, 4-hexenylene, 5-hexenylene, etc. Suitable lower alkenylene for L² is, for example, vinylene, 1-methylvinylene, 2-methylvinylene, etc.

 Suitable "aryl" includes C₆-C₁₆ aryl such as phenyl, naphthyl, anthryl, pyrenyl, phenanthryl, azulenyl, etc.

 Suitable "aryloxy" includes C₆-C₁₆ aryloxy such as phenoxy, naphthyloxy, anthryloxy, pyrenyloxy, phenanthryloxy, azulenyloxy, etc.

 Suitable "aryl(lower)alkyl" includes phenyl(C₁-C₆)alkyl such as benzyl, phenethyl, phenylpropyl, phenylbutyl, phenylhexyl, etc., naphthyl(C₁-C₆)alkyl such as naphthylmethyl, naphthylethyl, naphthylpropyl, naphthylbutyl, naphthylpentyl, naphthylhexyl, etc.

 Suitable "arylcarbonyl" includes arylcarbonyl in which the aryl portion is C₆-C₁₆ aryl such as phenylcarbonyl (benzoyl), naphthylcarbonyl, anthrylcarbonyl, pyrenylcarbonyl, phenanthrylcarbonyl, azulenylcarbonyl, etc.

 Suitable "aryl(lower)alkoxy" includes phenyl(C₁-C₆)alkoxy such as benzyloxy, phenethyloxy, phenylpropyloxy, phenylbutyloxy, phenylhexyloxy, etc., naphthyl(C₁-C₆)alkoxy such as naphthylmethyloxy, naphthylethyloxy, naphthylpropyloxy, naphthylbutyloxy, naphthylpentyloxy, naphthylhexyloxy, etc.

 Suitable "amino" includes unsubstituted amino, and amino mono- or di-substituted with substituent(s) selected from lower alkyl, lower alkanoyl and cycloalkyl such as N-(C₁-C₆ alkyl)amino (e.g., N-methylamino, N-ethylamino, N-propylamino, N-(n-butyl)amino, N-isobutylamino, N-(t-butyl)amino, etc.), N-(C₁-C₆ alkanoyl)amino

(e.g., N-acetylamino, N-ethylcarbonylamino, N-propylcarbonylamino, N-(n-butylcarbonyl)amino, N-isobutylcarbonylamino, N-(t-butylcarbonyl)amino, etc.), N-(C₃-C₆ cycloalkyl)amino (e.g., N-cyclopropylamino, N-cyclobutylamino, N-cyclopentylamino, N-cyclohexylamino, etc.), N,N-di(C₁-C₆ alkyl)amino (e.g., N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, etc.), etc.

Suitable "carbamoyl optionally mono- or di- substituted with lower alkyl(s)" includes carbamoyl; N-(lower)alkylcarbamoyl in which the alkyl portion is alkyl having 1 to 6 carbon atom(s) such as N-methylcarbamoyl, N-ethylcarbamoyl, N-propylcarbamoyl, N-butylcarbamoyl, N-isobutylcarbamoyl, N-tert-butylcarbamoyl, N-pentylcarbamoyl, N-neopentylcarbamoyl, N-isopentylcarbamoyl, N-hexylcarbamoyl, etc.; N,N-di(lower)alkylcarbamoyl in which the alkyl portions are each alkyl having 1 to 6 carbon atom(s) such as N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl, N,N-dibutylcarbamoyl, N,N-diisobutylcarbamoyl, N,N-di-tert-butylcarbamoyl, N,N-dipentylcarbamoyl, N,N-dineopentylcarbamoyl, N,N-diisopentylcarbamoyl, N,N-dihexylcarbamoyl, N-ethyl-N-methylcarbamoyl, N-methyl-N-propylcarbamoyl, N-butyl-N-methylcarbamoyl, N-methyl-N-isobutylcarbamoyl, etc.

The "heteroaryl" includes groups having 5 to 14 ring atoms and π electrons shared in a cyclic array and containing 1 to 4 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur besides carbon atoms. Suitable "heteroaryl" includes thienyl, benzothienyl, furyl, benzofuryl, dibenzofuryl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, indolyl, quinolyl, isoquinolyl, quinoxalinyll, tetrazolyl, oxazolyl, thiazolyl, isoxazolyl, etc.

The "heteroaryl" and "(lower)alkyl" of the "heteroaryl(lower)alkyl" are similar to those exemplified for the "heteroaryl" and "(lower)alkyl", respectively. Suitable "heteroaryl(lower)alkyl" includes pyridylmethyl, pyridylethyl, quinolylmethyl, etc.


Each of the two "(lower)alkyl" of the "(lower)alkyl-carbonyl(lower)alkyl" is similar to that exemplified for the "(lower)alkyl". Suitable "(lower)alkyl-carbonyl(lower)alkyl"


includes acetylmethyl, ethylcarbonylmethyl, etc.

The "non-aromatic heterocycle" includes a group having 5 to 14 ring atoms and containing 1 to 3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur besides carbon atoms. Suitable "non-aromatic heterocycle" includes pyrrolidinyl, pyrazolidinyl, imidazolidinyl, isoxazolidinyl, isothiazolidinyl, piperidyl (e.g., piperidino, etc.), piperazinyl, morpholinyl (e.g., morpholino, etc.), thiomorpholinyl (e.g., thiomorpholino, etc.), etc.

Suitable "N-containing heterocyclic ring" for the "N-containing heterocyclic ring optionally substituted with one or more substituent(s)" includes N-containing condensed heterocyclic ring such as indolyl, isoindolyl, indolidinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, quinoxalinyl, imidazopyridyl (e.g., imidazo[4,5-c]pyridyl, etc.), tetrahydroimidazopyridyl (e.g., 4,5,6,7-tetrahydro[4,5-c]pyridyl, etc.), 7-azabicyclo[2.2.1]heptyl, 3-azabicyclo[3.2.2]nonanyl, pyridoimidazolyl (e.g. pyrido[3,2-d]imidazolyl, pyrido[4,3-d]imidazolyl, etc.), azabenzimidazolyl, etc., and N-containing heteroaryl ring such as imidazolyl, thiazolyl, pyrazolyl, oxazolyl, etc.

Specifically, the preferred N-containing heterocyclic ring optionally substituted with one or more substituent(s) represented by R¹ includes, for example, the groups represented by the following formulas





wherein R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} and R^{14} are as defined above.

- 5 In the above formulas, suitable R^4 is hydrogen or a group such as (1) lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl, etc.); (2) lower alkoxy (e.g. methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, etc.); (3) aryl optionally substituted with substituent(s) selected from the group consisting
- 10 of halogen, lower alkanoyl, lower alkylsulfonyl, lower alkoxy and di(lower)alkylamino (e.g., phenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-acetylphenyl, 4-(N,N-dimethylamino)phenyl, 4-fluorophenyl, 4-methylsulfonylphenyl, etc.); (4) lower alkanoyl (e.g., acetyl, etc.); (5) lower alkoxy-carbonyl (e.g., methoxycarbonyl, etc.); (6)
- 15 arylcarbonyl (e.g., benzylcarbonyl, etc.); (7) aryl(lower)alkoxy (e.g., benzyloxy, etc.); (8) amino optionally mono- or di-substituted with substituent(s) selected from the group consisting of lower alkyl, lower alkanoyl and cycloalkyl (e.g., amino, N,N-dimethylamino, N,N-diethylamino, N-propylcarbonylamino, N-
- 20 cyclopentylamino, etc.); (9) halo(lower)alkyl (e.g., trifluoromethyl, etc.); (10) aryloxy (e.g., phenoxy, etc.); (11) aryl(lower)alkyl optionally substituted with hydroxy (e.g., hydroxyphenylmethyl, etc.); (12) carboxyl; (13) nitro; (14) cyano; (15) halogen (e.g., fluorine, chlorine, bromine, etc.); (16)
- 25 heteroaryl (e.g., thienyl, tetrazolyl, pyridyl, etc.); (17) non-aromatic heterocycle optionally substituted with lower alkyl (e.g., 4-methylpiperadinyll, morpholino, piperidino, etc.); (18) hydroxy, (19) (lower)alkylsulfonylcarbonyl (e.g., methylsulfonylcarbonyl, etc.), (20) non-aromatic heterocycle carbonyl (e.g., pyrrolidin-1-
- 30 ylcarbonyl, etc.). Furthermore, heteroaryl(lower)alkyl (e.g. pyridyl(lower)alkyl such as pyridylmethyl, etc.), lower alkyl-carbonyl(lower)alkyl (e.g., acetylmethyl, etc.), etc. can be also used for R^4 .

In the above formulas, suitable R^5 is hydrogen or a group selected from the group consisting of lower alkyl (e.g., methyl, ethyl, propyl, butyl, etc.), aryl(lower)alkyl (e.g., benzyl, phenethyl, etc.), heteroaryl(lower)alkyl (e.g., pyridyl(lower)alkyl
5 such as pyridylmethyl, etc.) and lower alkyl-carbonyl(lower)alkyl (e.g., acetylmethyl, etc.). Preferably, R^5 is hydrogen or a group selected from the group consisting of lower alkyl and aryl(lower)alkyl.

In the above formulas, suitable R^6 , R^7 and R^8 are each
10 hydrogen or lower alkyl (e.g., methyl, ethyl, propyl, butyl, etc.).

In the above formulas, suitable R^9 is hydrogen or a group such as (1) lower alkyl optionally substituted with di(lower)alkylamino (e.g., dimethylaminoethyl, etc.); (2) aryl optionally substituted with lower alkoxy (e.g., phenyl, o-
15 methoxyphenyl, m-methoxyphenyl, p-methoxyphenyl, etc.); (3) (lower)alkoxy-carbonyl (e.g., methoxycarbonyl, etc.); (4) cyano; (5) carbamoyl optionally mono- or di-substituted with (lower)alkyl (e.g., carbamoyl, N,N-dimethylcarbamoyl, N-isopropylcarbamoyl, etc.); (6) halogen (e.g., chloro, bromo, etc.); (7) (lower)alkyl-
20 carbonyl (e.g., acetyl, etc.); (8) arylcarbonyl (e.g., benzoyl, etc.); (9) cyclo(lower)alkyl (e.g., cyclohexyl, etc.), etc.

In the above formulas, suitable substituent represented by R^{10} is hydrogen or a group such as (1) (lower)alkylcarbamoyl (e.g., N-methylcarbamoyl, etc.); (2) di(lower)alkylcarbamoyl (e.g., N,N-
25 dimethylcarbamoyl, etc.); (3) aryl optionally substituted with halogen (e.g., phenyl, p-chlorophenyl, p-fluorophenyl, etc.); (4) (lower)alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, etc.); (5) carboxy; (6) non-aromatic heterocycle carbonyl (e.g., piperidinylcarbonyl, etc.); (7) halogen (e.g., chloro, etc.); (8)
30 (lower)alkyl optionally substituted with hydroxy or (lower)alkoxy, non-aromatic heterocycle, aryl, di(lower)alkylamino or halogen (e.g., hydroxymethyl, methoxymethyl, piperidinomethyl, morpholinomethyl, N,N-dimethylaminomethyl, trifluoromethyl, etc.); and (9) adamantyl, etc.

35 In the above formulas, suitable substituent represented by R^{11} is hydrogen or aryl(lower)alkyl in which the aryl portion is

substituted with lower alkoxy (e.g., p-methoxyphenylmethyl, etc.), etc.


In the above formulas, suitable substituent represented by R^{12} is hydrogen or a group selected from lower alkyl (e.g., methyl, etc.), aryl optionally substituted with halogen (e.g., phenyl, p-chlorophenyl, etc.), etc.

In the above formulas, suitable substituent represented by R^{13} is hydrogen or a group selected from lower alkyl (e.g., methyl, etc.), aryl (e.g., phenyl, etc.), etc.

In the above formulas, suitable substituent represented by R^{14} is hydrogen or lower alkyl (e.g., methyl, etc.), etc.

Suitable "n" of the " $-(CH_2)_n-$ " for L^1 is an integer of 0 to 6, preferably 1 or 2. The " $-(CH_2)_n-$ " is optionally substituted with one or more suitable substituent(s) such as lower alkyl (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, etc.), lower alkoxy (e.g., methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, etc.), aryl(lower)alkyl (e.g., benzyl, etc.), etc. Furthermore, one or more methylenes (e.g., one methylene, etc.) may be replaced with suitable heteroatoms (e.g., oxygen atom, etc.).

Suitable " $-(CH_2)_n-$ " for L^1 of the present invention includes, for example, $-CH_2-$, $-(CH_2)_2-$, $-CH(CH_3)-$, $-CH_2-O-$, $-(CH_2)_3-$, $-(CH_2)_3-O-$, $-(CH_2)_4-O-$,



Suitable "hydroxy protecting group" is as follows:

- 25 lower alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, hexyl, etc.), preferably methyl;
 lower alkoxy(lower)alkyl (e.g., methoxymethyl, etc.);
 lower alkoxy(lower)alkoxy(lower)alkyl (e.g., 2-methoxyethoxymethyl, etc.);
- 30 ar(lower)alkyl in which the aryl portion is optionally substituted with one or more suitable substituent(s) (e.g., benzyl (Bn), p-methoxybenzyl, m,p-dimethoxybenzyl, etc.), preferably benzyl;
 ar(lower)alkoxy(lower)alkyl in which the aryl portion is optionally

- substituted with one or more suitable substituent(s) (e.g., benzyloxymethyl, p-methoxybenzyloxymethyl, etc.); (lower)alkylthio(lower)alkyl (e.g., methylthiomethyl, ethylthiomethyl, propylthiomethyl, isopropylthiomethyl, butylthiomethyl, isobutylthiomethyl, hexylthiomethyl, etc.), etc., preferably methylthiomethyl;
- trisubstituted silyl such as tri(lower)alkylsilyl (e.g., trimethylsilyl, triethylsilyl, tributylsilyl, tert-butyltrimethylsilyl, tri-tert-butylsilyl, etc.), lower alkyldiarylsilyl (e.g. methyldiphenylsilyl, ethyldiphenylsilyl, propyldiphenylsilyl, tert-butyl-diphenylsilyl (TBDPS), etc.), etc., preferably tert-butyl-dimethylsilyl (TBDMS) and tert-butyl-diphenylsilyl;
- heterocyclic group (e.g., tetrahydropyranyl, etc.);
- acyl as described below [e.g., aliphatic acyl such as lower alkanoyl (e.g., acetyl, propanoyl, pivaloyl, etc.); aromatic acyl (e.g., benzoyl (Bz), toluoyl, naphthoyl, fluorenylcarbonyl, etc.); lower alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, t-butoxycarbonyl, pentyloxycarbonyl, hexyloxycarbonyl, etc.), etc.;
- ar(lower)alkoxycarbonyl in which the aryl portion is optionally substituted with one or more suitable substituent(s) (e.g., benzyloxycarbonyl, bromobenzyloxycarbonyl, etc.);
- lower alkylsulfonyl (e.g., methylsulfonyl, ethylsulfonyl, etc.); lower alkoxy-sulfonyl (e.g., methoxysulfonyl, ethoxysulfonyl, etc.); ar(lower)alkanoyl (e.g., phenylacetyl, phenylpropanoyl, phenylbutanoyl, phenylisobutanoyl, phenylpentanoyl, phenylhexanoyl, naphthylacetyl, naphthylpropanoyl, naphthylbutanoyl, naphthylisobutanoyl, naphthylpentanoyl, naphthylhexanoyl, etc.);
- ar(lower)alkenoyl such as ar(C₃-C₆)alkenoyl (e.g., phenylpropenoyl, phenylbutenoyl, phenylmethacryloyl, phenylpentenoyl, phenylhexenoyl, naphthylpropenoyl, naphthylbutenoyl, naphthylmethacryloyl, naphthylpentenoyl, naphthylhexenoyl, etc.), etc.];
- lower alkenyl (e.g., vinyl, allyl, etc.);, etc.

The preferable hydroxy protecting group for the present

invention is, for example, tetrahydropyranyl, trimethylsilyl, t-butyl dimethylsilyl, etc.

The following abbreviations are also used in the present specification: Boc (t-butyloxycarbonyl); HOBT or HOBT (1-hydroxybenzotriazole); WSCD (1-ethyl-3-(3'-dimethylaminopropyl)-carbodiimide); DMF (N,N-dimethylformamide); aq. (aqueous solution); Me (methyl); MeOH (methanol); Et (ethyl); EtOH (ethanol); tBu (t-butyl); TsCl (p-toluenesulfonyl chloride); Ac (acetyl); AcOH (acetic acid); AcOEt (ethyl acetate); AcONH₄ (ammonium acetate); Ph (phenyl); DIEA (diisopropylethylamine); THP (tetrahydropyranyl); THF (tetrahydrofuran) and TFA or TFAOH (trifluoroacetic acid).

Test Method

In order to show the usefulness of the compound [I] of the invention, the pharmacological test result of the representative compound of the present invention is shown in the following.

Test 1: Determination of histone deacetylase inhibitor activity

The partial purification of human histone deacetylase, the preparation of [³H] acetyl histones, and the assay for histone deacetylase activity were performed basically according to the method as proposed by Yoshida et al. as follows.

Partial purification of human histone deacetylase

The human histone deacetylase was partially purified from human T cell leukemia Jurkat cells. Jurkat cells (5×10^8 cells) were suspended in 40 mL of the HDA buffer consisting of 15 mM potassium phosphate, pH 7.5, 5% glycerol and 0.2 mM EDTA. After homogenization, nuclei were collected by centrifugation ($35,000 \times g$, 10 min) and homogenized in 20 mL of the same buffer supplemented with 1 M (NH₄)₂SO₄. The viscous homogenate was sonicated and clarified by centrifugation ($35,000 \times g$, 10 min), and the deacetylase was precipitated by raising the concentration of (NH₄)₂SO₄ to 3.5 M. The precipitated protein was dissolved in 10 mL of the HDA buffer and dialyzed against 4 liters of the same buffer. The dialyzate was then loaded onto a DEAE-cellulose (Whatman DE52) column (25 × 85 mm) equilibrated with the same buffer and eluted with 300 mL of a linear gradient (0-0.6 M) of NaCl. A single peak of histone deacetylase activity appeared between 0.3 and 0.4 M NaCl.

Preparation of [³H] acetyl histone

To obtain [³H] acetyl-labeled histone as the substrate for the histone deacetylase assay, 1×10^8 cells of Jurkat in 20 mL of RPMI-1640 medium (supplemented with 10% FBS, penicillin (50
5 units/mL) and streptomycin (50 μ g/mL)) were incubated with 300 MBq [³H] sodium acetate in the presence of 5 mM sodium butyrate for 30 minutes in 5% CO₂-95% air atmosphere at 37°C in a 75 cm² flask, harvested into a centrifuge tube (50 mL), collected by centrifugation at 1000 rpm for 10 minutes, and washed once with
10 phosphate-buffered saline. The washed cells were suspended in 15 mL of ice-cold lysis buffer (10 mM Tris-HCl, 50 mM sodium bisulfite, 1% Triton X-100, 10 mM MgCl₂, 8.6% sucrose, pH 6.5). After Dounce homogenization (30 stroke), the nuclei were collected by centrifugation at 1000 rpm for 10 minutes, washed 3 times with 15
15 mL of the lysis buffer, and once with 15 mL of ice-cooled washing buffer (10 mM Tris-HCl, 13 mM EDTA, pH 7.4) successively. The pellet was suspended in 6 mL of ice-cooled water using a mixer, and 68 μ l of H₂SO₄ was added to the suspension to give a concentration of 0.4 N. After incubation at 4°C for 1 hour, the suspension was
20 centrifuged for 5 minutes at 15, 000 rpm, and the supernatant was taken and mixed with 60 mL of acetone. After overnight incubation at -20°C, the coagulated material was collected by microcentrifugation, air-dried, and stored at -80°C.

Assay for histone deacetylase activity

25 For the standard assay, 10 μ l of [³H] acetyl-labeled histones were added to 90 μ l of the enzyme fraction, and the mixture was incubated at 25°C for 30 minutes. The reaction was stopped by addition of 10 μ l of HCl. The released [³H] acetic acid was extracted with 1 mL of ethyl acetate, and 0.9 mL of the solvent
30 layer was taken into 10 mL of toluene scintillation solution for determination of radioactivity.

Test 2: Determination of T-cell growth inhibitor activity

The T lymphocyte blastogenesis test was performed in microtiter plates with each well containing 1.5×10^5 splenic cells
35 of Lewis rats in 0.1 mL RPMI-1640 medium supplemented with 10% fetal bovine serum (FBS), 50 mM 2-mercaptoethanol, penicillin (100

units/mL) and streptomycin (100 µg/mL), to which Concanavalin A (1 µg/mL) was added. The cells were incubated at 37°C in a humidified atmosphere of 5% CO₂ for 72 hours. After the culture period, suppressive activities of the test compounds in T lymphocyte blastogenesis were quantified by AlamarBlue (trademark) Assay. The test samples were dissolved in DMSO and further diluted with RPMI-1640 medium and added to the culture. The activities of the test compounds were expressed as IC₅₀.

The results of those tests are shown in the Table 1.

10

Table 1: HDAC inhibitory activity and T-cell growth inhibitory activity of the compound of the present invention

Examples	Test 1: HDAC inhibitory activity IC ₅₀ (nM)	Test 2: T-cell growth inhibitory activity IC ₅₀ (nM)
Compound E1	28	69
Compound E3	140	160
Compound E5	96	310
Compound E6	150	150

The pharmaceutical composition of the present invention comprising histone deacetylase inhibitor such as the compound [I] is useful as a therapeutic or prophylactic agent for diseases caused by abnormal gene expression, such as inflammatory disorders, diabetes, diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), protozoal infection, etc. Furthermore, it is useful as an antitumor agent or immunosuppressant, which prevents an organ transplant rejection and autoimmune diseases as exemplified below:

rejection reactions by transplantation of organs or tissues such as the heart, kidney, liver, bone marrow, skin, cornea, lung, pancreas, small intestine, limb, muscle, nerve, intervertebral disc, trachea,

25

myoblast, cartilage, etc.;

graft-versus-host reactions following bone marrow transplantation;
autoimmune diseases such as rheumatoid arthritis, systemic lupus
erythematosus, Hashimoto's thyroiditis, multiple sclerosis,

5 myasthenia gravis, type I diabetes, etc.; and

infections caused by pathogenic microorganisms (e.g., *Aspergillus*
fumigatus, *Fusarium oxysporum*, *Trichophyton asteroides*, etc.).

Furthermore, pharmaceutical preparations of the histone
deacetylase inhibitor, such as the compound [I], are useful for the
10 therapy or prophylaxis of the following diseases.

Inflammatory or hyperproliferative skin diseases or
cutaneous manifestations of immunologically-mediated diseases (e.g.,
psoriasis, atopic dermatitis, contact dermatitis, eczematoid
dermatitis, seborrheic dermatitis, lichen planus, pemphigus,
15 bullous pemphigoid, epidermolysis bullosa, urticaria, angioedema,
vasculitides, erythema, dermal eosinophilia, lupus erythematosus,
acne, alopecia areata, etc.);

autoimmune diseases of the eye (e.g., keratoconjunctivitis, vernal
conjunctivitis, uveitis associated with Behcet's disease, keratitis,
20 herpetic keratitis, conical keratitis, corneal epithelial dystrophy,
keratoleukoma, ocular pemphigus, Mooren's ulcer, scleritis,
Grave's ophthalmopathy, Vogt-Koyanagi-Harada syndrome,
keratoconjunctivitis sicca (dry eye), phlyctenule, iridocyclitis,
sarcoidosis, endocrine ophthalmopathy, etc.);

25 reversible obstructive airways diseases [asthma (e.g., bronchial
asthma, allergic asthma, intrinsic asthma, extrinsic asthma, dust
asthma, etc.), particularly chronic or inveterate asthma (e.g.,
late asthma, airway hyper-responsiveness, etc.), bronchitis, etc.];
mucosal or vascular inflammations (e.g., gastric ulcer, ischemic or
30 thrombotic vascular injury, ischemic bowel diseases, enteritis,
necrotizing enterocolitis, intestinal damages associated with
thermal burns, leukotriene B₄-mediated diseases, etc.);

intestinal inflammations/allergies (e.g., coeliac diseases,
proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's
35 disease, ulcerative colitis, etc.);

food-related allergic diseases with symptomatic manifestation

- remote from the gastrointestinal tract (e.g., migraine, rhinitis, eczema, etc.);
- renal diseases (e.g., interstitial nephritis, Goodpasture's syndrome, hemolytic uremic syndrome, diabetic nephropathy, etc.);
- 5 nervous diseases (e.g., multiple myositis, Guillain-Barre syndrome, Meniere's disease, multiple neuritis, solitary neuritis, cerebral infarction, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS), radiculopathy, etc.);
- cerebral ischemic diseases (e.g., head injury, hemorrhage in brain
10 (e.g., subarachnoid hemorrhage, intracerebral hemorrhage, etc.), cerebral thrombosis, cerebral embolism, cardiac arrest, stroke, transient ischemic attack (TIA), hypertensive encephalopathy, etc.);
- endocrine diseases (e.g., hyperthyroidism, Basedow's disease,
15 etc.);
- hematic diseases (e.g., pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia, etc.);
- 20 bone diseases (e.g., osteoporosis, etc.);
- respiratory diseases (e.g., sarcoidosis, pulmonary fibrosis, idiopathic interstitial pneumonia, etc.);
- skin diseases (e.g., dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photosensitivity, cutaneous T-cell lymphoma,
25 etc.);
- circulatory diseases (e.g., arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa, myocarditis, etc.);
- collagen diseases (e.g., scleroderma, Wegener's granuloma, Sjögren's syndrome, etc.);
- 30 adiposis;
- eosinophilic fasciitis;
- periodontal diseases (e.g., damage to gingiva, periodontium, alveolar bone or substantia ossea dentis, etc.);
- nephrotic syndrome (e.g., glomerulonephritis, etc.);
- 35 male pattern alopecia, alopecia senile;
- muscular dystrophy;

- pyoderma and Sezary syndrome;
chromosome abnormality-associated diseases (e.g., Down's syndrome, etc.);
Addison's disease;
- 5 active oxygen-mediated diseases (e.g., organ injury [e.g., ischemic circulation disorders of organs (e.g., heart, liver, kidney, digestive tract, etc.) associated with preservation, transplantation, ischemic diseases (e.g., thrombosis, cardiac infarction, etc.), etc.];
- 10 intestinal diseases (e.g., endotoxin shock, pseudomembranous colitis, drug- or radiation-induced colitis, etc.);
renal diseases (e.g., ischemic acute renal insufficiency, chronic renal failure, etc.);
pulmonary diseases (e.g., toxicosis caused by pulmonary oxygen or
- 15 drugs (e.g., paracort, bleomycin, etc.), lung cancer, pulmonary emphysema, etc.);
ocular diseases (e.g., cataracta, iron-storage disease (siderosis bulbi), retinitis, pigmentosa, senile plaques, vitreous scarring, corneal alkali burn, etc.);
- 20 dermatitis (e.g., erythema multiforme, linear immunoglobulin A bullous dermatitis, cement dermatitis, etc.); and
other diseases (e.g., gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution (e.g., air pollution, etc.), aging, carcinogen, metastasis of carcinoma,
- 25 hypobaropathy, etc.));
diseases caused by histamine release or leukotriene C4 release;
restenosis of coronary artery following angioplasty and prevention of postsurgical adhesions;
autoimmune diseases and inflammatory conditions (e.g., primary
- 30 mucosal edema, autoimmune atrophic gastritis, premature menopause, male sterility, juvenile diabetes mellitus, pemphigus vulgaris, pemphigoid, sympathetic ophthalmitis, lens-induced uveitis, idiopathic leukopenia, active chronic hepatitis, idiopathic cirrhosis, discoid lupus erythematosus, autoimmune orchitis,
- 35 arthritis (e.g., arthritis deformans, etc.), polychondritis, etc.);
Human Immunodeficiency Virus (HIV) infection, AIDS;

allergic conjunctivitis;
hypertrophic cicatrix, keloid due to trauma, burn or surgery, etc.

Therefore, the pharmaceutical composition of the present invention is useful for the therapy and prophylaxis of liver
5 diseases [e.g., immunogenic diseases (e.g., chronic autoimmune liver diseases such as autoimmune hepatic diseases, primary biliary cirrhosis, sclerosing cholangitis, etc.), partial liver resection, acute liver necrosis (e.g., necrosis caused by toxins, viral hepatitis, shock, anoxia, etc.), hepatitis B, non-A non-B hepatitis,
10 hepatocirrhosis, hepatic failure (e.g., fulminant hepatitis, late-onset hepatitis, "acute-on-chronic" liver failure (acute liver failure on chronic liver diseases, etc.), etc.), etc.].

The pharmaceutical composition of the present invention can be used in the form of pharmaceutical preparation, for example, in
15 a solid, semisolid or liquid form, which contains the histone deacetylase inhibitor, such as the compound [I], as an active ingredient in admixture with an organic or inorganic carrier or excipient suitable for external, enteral or parenteral administrations. The active ingredient may be compounded, for
20 example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, injections, ointments, liniments, eye drops, lotion, gel, cream, and any other form suitable for use.

The carriers those can be used for the present invention
25 include water, glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea and other carriers suitable for use in manufacturing preparations in a solid, semisolid, or liquid form. Furthermore, auxiliary, stabilizing, thickening,
30 solubilizing and coloring agents and perfumes may be used.

For applying the composition to human, it is preferable to apply it by intravenous, intramuscular, topical or oral administration, or by a vascular stent impregnated with the compound [I]. While the dosage of therapeutically effective amount
35 of the histone deacetylase inhibitor, such as the compound [I], varies from and also depends upon the age and condition of each

individual patient to be treated, when an individual patient is to be treated, in the case of intravenous administration, a daily dose of 0.01-10 mg of the histone deacetylase inhibitor, such as the compound [I], per kg weight of human being, in the case of
5 intramuscular administration, a daily dose of 0.1-10 mg of the histone deacetylase inhibitor, such as the compound of the formula [I], per kg weight of human being, and in the case of oral administration, a daily dose of 0.5-50 mg of the histone deacetylase inhibitor, such as the compound [I], per kg weight of
10 human being, is generally given for treatment.

During the preparation of the above-mentioned pharmaceutical administration forms, the compound [I] or a salt thereof can also be combined together with other immunosuppressive substances, for example repamycin, mycophenolic acid, cyclosporin A, tacrolimus or
15 brequinar sodium.

Hereinafter the reactions in each Preparations and Examples for preparing the compound [I] of the present invention are explained in more detail. The invention should not be restricted by the following Preparations and Examples in any way.
20

Preparation 1

To a solution of 4-iodophenylacetic acid (1346 mg) in N,N-dimethylformamide (15 mL) were added tert-butyl 2-aminophenylcarbamate (1.07 g), 1-hydroxybenzotriazole (HOBT) (764
25 mg), 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (1.08 g), and the mixture was stirred at ambient temperature for 3 hours. The mixture was poured into water and extracted with ethyl acetate. The organic phase was sequentially washed with saturated aqueous ammonium chloride solution, saturated aqueous sodium
30 hydrogen carbonate solution and brine. The organic phase was dried over magnesium sulfate and evaporated in vacuo. The residue was purified by silica gel chromatography eluting with a mixture of hexane and ethyl acetate (4:1 to 2:1) to give Compound (1) as a pale yellow amorphous (2.03 g).

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.50 (3x3H, s), 3.66 (2H, s), 6.62 (1H, brs), 7.07-7.20 (4H, m), 7.33 (1H, m), 7.47 (1H, m), 7.69 (2x1H, d,

J=8.3 Hz), 8.00 (1H, brs);

MASS (ES+): m/e 453.

Preparation 2

To a stirred solution of Compound (1) (25.6 g) in ethanol
5 (300 mL) was added concentrated hydrochloric acid (30 mL), and the
mixture was refluxed for 1 hour. The solvent was evaporated *in*
vacuo azeotropically with toluene. The residual solid was
collected with the mixture of ethanol and ethyl acetate (1:10) to
give Compound (2) as an orange solid (20.0 g).

10 ¹H-NMR (300 MHz, CDCl₃, δ): 4.52 (2H, s), 7.30 (2x1H, d, J=8.3 Hz),
7.49-7.57 (2H, m), 7.73-7.82 (4H, m);

MASS (ES+): m/e 335.

Preparation 3

To a stirred solution of Compound (2) (114 mg) in dioxane (3
15 mL) and 1N-sodium hydroxide (0.8 mL) was added p-toluenesulfonyl
chloride (70 mg) at 0°C. The mixture was allowed to warm to
ambient temperature and stirred for 30 minutes. Additional p-
toluenesulfonyl chloride (70 mg) was added, then 1N-sodium
hydroxide (0.5 mL) was added so that the final pH was 9. The
20 mixture was stirred at ambient temperature for 2 hours. The
solvent was evaporated *in vacuo* and the resulting solution was
extracted with ethyl acetate. The organic phase was washed with
brine, dried over magnesium sulfate and evaporated *in vacuo*. The
residue was purified by preparative thin layer chromatography
25 (hexane: ethyl acetate=2:1) to give Compound (3) as a pale yellow
amorphous (130 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.35 (3H, s), 4.56 (2H, s), 7.05 (2x1H,
d, J=8.5 Hz), 7.32-7.44 (4H, m), 7.63-7.70 (3H, m), 7.78 (2x1H, d,
J=8.5 Hz), 7.94 (1H, d, J=6.5 Hz);

30 MASS (ES+): m/e 489.

Preparation 4

To a stirred solution of Compound (3) (1,137 mg) in N,N-
dimethylformamide (15 mL) were added acrylic acid (0.8 mL),
palladium(II) acetate (26 mg), tris(2-methylphenyl)phosphine (142
35 mg) and N,N-diisopropylethylamine (1.25 mL). The mixture was
stirred at 120°C for 90 minutes. The resulting mixture was allowed

to cool to ambient temperature, poured into water and extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica column chromatography eluting with a mixture of
5 chloroform and methanol (20:1) to give Compound (4) as a pale yellow amorphous (455 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.32 (3H, s), 4.63 (2H, s), 6.51 (1H, d, $J=16$ Hz), 7.26-7.44 (6H, m), 7.54-7.69 (4H, m), 7.79 (2x1H, d, $J=8.4$ Hz), 7.94 (1H, m);

10 MASS (ES+): m/e 433.

Preparation 5

To a stirred solution of Compound (4) (70 mg) in *N,N*-dimethylformamide (3 mL) were added 1-hydroxybenzotriazole (HOBT) (26 mg), 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide
15 hydrochloride (37 mg) and *O*-(tetrahydro-2H-pyran-2-yl)hydroxylamine (23 mg), and the resulting mixture was stirred at ambient temperature for 14 hours. To the reaction mixture were added additional 1-hydroxybenzotriazole (13 mg), 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (19 mg) and *O*-
20 (tetrahydro-2H-pyran-2-yl)hydroxylamine (12 mg), and the mixture was stirred for 6 hours. The reaction mixture were diluted with ethyl acetate and washed succesively with water, saturated ammonium chloride solution, saturated sodium hydrogen carbonate solution and brine. The organic phase was dried over magnesium sulfate and
25 concentrated *in vacuo*. The residue was purified by preparative thin layer chromatography (chloroform:methanol=10:1) to give Compound (5) as a white amorphous (503 mg). The Compound (5) was used in Example 1.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.44-1.76 (6H, m), 3.52 (1H, m), 3.95
30 (1H, m), 4.19 (1H, m), 4.90 (1H, m), 6.47 (1H, d, $J=15.8$ Hz), 7.09-7.19 (2H, m), 7.34-7.58 (7H, m), 11.23 (1H, s), 12.30 (1H, s);

MASS (ES+): m/e 378.

Preparation 6

To a stirred solution of (4-bromophenyl)acetic acid (80.0 g,
35 372 mmol) in *N,N*-dimethylformamide (640 mL) were added acrylic acid *t*-butyl ester (95.4 g), palladium(II) acetate (1.67 g),

triphenylphosphine (3.91 g) and N,N-diisopropylethylamine (162 mL). The mixture was stirred at 100°C for 7 hours. The resulting mixture was allowed to cool to ambient temperature, poured into 1N-hydrochloric acid and extracted with ethyl acetate twice. The
5 combined organic phase was extracted with saturated sodium hydrogen carbonate solution three times. The combined aqueous phase was acidified with concentrated hydrogen chloride to pH 2 and extracted with ethyl acetate. The organic phase was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give Compound
10 (6) as a pale yellow solid (78.1 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 3.67 (2H, s), 6.35 (1H, d, J=16 Hz), 7.29 (2H, d, J=8 Hz), 7.47 (2H, d, J=8 Hz), 7.56 (1H, d, J=16 Hz).

Preparation 7

15 To a solution of Compound (6) (77.7 g), tert-butyl 2-aminophenylcarbamate (61.7 g) and 1-hydroxybenzotriazole (HOBT) (44.0 g) in N,N-dimethylformamide (777 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (62.5 g) at 4°C. The mixture was warmed to ambient temperature and stirred for
20 2 hours. The reaction mixture was added saturated aqueous sodium hydrogencarbonate (777 mL) and water (3.1 L), and extracted with ethyl acetate (1.5 L). The organic layer was washed with 5% aqueous potassium hydrogen sulfate (500 mL), saturated aqueous sodium hydrogencarbonate (500 mL) and brine (500 mL), dried over
25 magnesium sulfate, filtered and evaporated in vacuo to give Compound (7) (135 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.49 (9H, s), 1.54 (9H, s), 3.74 (2H, s), 6.36 (1H, d, J=16 Hz), 6.66 (1H, brs), 7.10-7.20 (2H, m), 7.33-7.40 (3H, m), 7.44-7.54 (3H, m), 7.57 (1H, d, J=16 Hz), 7.98 (1H, brs).

30 Preparation 8

A solution of Compound (7) (47.6 g) in 1N-hydrogen chloride in acetic acid (60 mL) was heated at 120°C for 1 hour. The resulting mixture was allowed to cool to ambient temperature and diluted with ethyl acetate. The resulted precipitate was filtered
35 and the residue was washed with ethyl acetate to give Compound (8) (28.9 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.56 (2H, s), 6.56 (1H, d, J=16 Hz), 7.48-7.55 (4H, m), 7.59 (1H, d, J=16 Hz), 7.72-7.80 (4H, m).

Preparation 9

To a solution of Compound (8) (50.0 g), O-tetrahydro-2H-pyran-2-ylhydroxylamine (29.8 g) and 1-hydroxybenzotriazole (34.3 g) in N,N-dimethylformamide (795 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (39.5 g) at 9°C. The mixture was warmed to ambient temperature and stirred for 2 hours. The reaction mixture was added saturated aqueous sodium hydrogencarbonate (795 mL) and water (3.2 L). The resulting precipitate was collected by filtration, and washed with saturated aqueous sodium hydrogencarbonate (250 × 2 mL) and water (250 × 2 mL) to give Compound (9) (57.2 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48-1.75 (6H, m), 3.48-3.57 (1H, m), 3.89-4.00 (1H, m), 4.20 (2H, s), 4.90 (1H, brs), 6.47 (1H, d, J=16 Hz), 7.07-7.16 (2H, m), 7.34-7.57 (7H, m), 11.2 (1H, brs), 12.3 (1H, brs).

Preparation 10

To a stirred solution of 2-(4-iodobenzyl)-1H-benzimidazole (451 mg) in dimethylformamide (5 mL) was added portionwise sodium hydride (81 mg, 60% oil dispersion) at 0°C. After 30 minutes, benzyl bromide (0.19 mL) was added dropwise to the mixture, and the mixture was stirred for 30 minutes. The resulting mixture was poured into saturated ammonium chloride solution and extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, and concentrated *in vacuo*. The residue was purified by preparative thin layer chromatography (chloroform:methanol=10:1) to give Compound (10) as a pale yellow oil (225 mg). In this preparation, a by-product (1-benzyl-2-[1-(4-iodophenyl)-2-phenylethyl]-1H-benzimidazole) (306 mg) was also obtained and was used in Preparation 23 described below.

¹H-NMR (300 MHz, CDCl₃, δ): 4.18 (2H, s), 5.19 (2H, s), 6.86-6.97 (4H, m), 7.19-7.32 (6H, m), 7.56 (2x1H, J=8.5 Hz), 7.81 (1H, d, J=7.5 Hz);

MASS (ES⁺): m/e 425.

Preparation 11

Compound (11) was obtained from Compound (10) according to a manner similar to Preparation 4 as a pale yellow oil (142 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.33 (3H, t, $J=7$ Hz), 4.26 (2H, s), 4.26 (2H, q, $J=7$ Hz), 5.21 (2H, s), 6.38 (1H, d, $J=16$ Hz), 6.88–6.96 (2H, m), 7.18–7.32 (8H, m), 7.41 (2x1H, d, $J=8$ Hz), 7.62 (1H, d, $J=16$ Hz), 7.81 (1H, d, $J=8$ Hz);

MASS (ES+): m/e 397.

Preparation 12

To a stirred solution of Compound (11) (140 mg) in methanol (6 mL) was added 1N sodium hydroxide solution (0.71 mL). The mixture was stirred at ambient temperature for 7 hours. The solvent was evaporated *in vacuo*, and the residue was dissolved in water and washed with diethyl ether. The aqueous phase was acidified to pH 3 with hydrochloric acid, and extracted three times with ethyl acetate. The combined organic phase was washed with brine, dried over sodium sulfate and evaporated *in vacuo* to give Compound (12) as a pale yellow powder (111 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 4.29 (2H, s), 5.23 (2H, s), 6.36 (1H, d, $J=15.7$ Hz), 6.88–6.96 (2H, m), 7.16–7.34 (8H, m), 7.41 (2x1H, d, $J=8$ Hz), 7.62 (1H, d, $J=15.7$ Hz), 7.81 (1H, d, $J=7.5$ Hz);

MASS (ES+): m/e 368.

Preparation 13

Compound (13) was obtained from Compound (12) according to a manner similar to Preparation 9 as a white amorphous (111 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.52–1.95 (6H, m), 3.61 (1H, m), 3.94 (1H, m), 4.25 (2H, s), 5.02 (1H, m), 5.20 (2H, s), 6.88–6.96 (2H, m), 7.12–7.41 (11H, m), 7.66 (1H, d, $J=15.5$ Hz), 7.82 (1H, d, $J=8$ Hz);

MASS (ES+): m/e 468.

Preparation 14

To a stirred solution of 3-phenylpropanoic acid (7.51 g) in acetic acid (70 mL) were added periodic acid (2.39 g), iodine (5.08 g), concentrated sulfuric acid (1.5 mL) and water (10 mL), and the mixture was stirred at 70°C for 7 hours. The solvent was evaporated *in vacuo*, and the residue was diluted with water and extracted with ethyl acetate. The organic phase was washed with

10% sodium thiosulfate solution twice, then washed with brine, dried over magnesium sulfate and evaporated *in vacuo*. The precipitate was crystallized from ethyl acetate and hexane to give Compound (14) (5.80 g).

- 5 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 2.66 (2H, t, $J=7$ Hz), 2.90 (2H, t, $J=7$ Hz), 6.97 (2x1H, d, $J=8.5$ Hz), 7.61 (2x1H, d, $J=8.5$ Hz);
MASS (ES⁻): m/e 275.

Preparation 15

Compound (15) was obtained from Compound (14) according to a
10 manner similar to Preparation 1 (9.50 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.51 (3x3H, s), 2.64 (2H, t, $J=7.5$ Hz), 3.00 (2H, t, $J=7.5$ Hz), 6.69 (1H, s), 7.00 (2x1H, d, $J=8.5$ Hz), 7.12-7.20 (2H, m), 7.33 (1H, m), 7.45 (1H, m), 7.62 (2x1H, d, $J=8.5$ Hz), 7.97 (1H, brs);

- 15 MASS (ES⁺): m/e 467.

Preparation 16

Compound (16) was obtained from Compound (15) according to a
manner similar to Preparation 2 (1.55 g).

20 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.20 (2H, t, $J=7.5$ Hz), 3.41 (2H, t, $J=7.5$ Hz), 7.10 (2x1H, d, $J=8.5$ Hz), 7.48-7.56 (2H, m), 7.66 (2x1H, d, $J=8.5$ Hz), 7.74-7.82 (2H, m);

MASS (ES⁺): m/e 349.

Preparation 17

Compound (17) was obtained from Compound (16) according to a
25 manner similar to Preparation 3 (7.10 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 2.38 (3H, s), 3.18 (2H, t, $J=7$ Hz), 3.43 (2H, t, $J=7$ Hz), 7.05 (2x1H, d, $J=8.5$ Hz), 7.25 (2x1H, d, $J=8.5$ Hz), 7.30-7.40 (2H, m), 7.61 (2x1H, d, $J=8.5$ Hz), 7.67 (1H, m), 7.71 (2x1H, d, $J=8.5$ Hz), 8.03 (1H, m);

- 30 MASS (ES⁺): m/e 503.

Preparation 18

Compound (18) was obtained from Compound (17) according to a
manner similar to Preparation 4 (3.59 g).

35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 2.38 (3H, s), 3.27 (2H, t, $J=7$ Hz), 3.47 (2H, t, $J=7$ Hz), 6.44 (1H, d, $J=16$ Hz), 7.25 (2x1H, d, $J=8$ Hz), 7.31-7.40 (4H, m), 7.50 (2x1H, d, $J=8$ Hz), 7.66-7.81 (4H, m), 8.04

(1H, m);

MASS (ES+): m/e 447.

Preparation 19

Compound (19) was obtained from Compound (18) according to a
5 manner similar to Preparation 5 (2.20 g). The Compound (19) was
used in Example 3.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.46-1.76 (6H, m), 3.08-3.18 (4H, m),
3.53 (1H, m), 3.95 (1H, m), 4.90 (1H, m), 6.45 (1H, d, J=15.5 Hz),
7.08-7.16 (2H, m), 7.31 (2x1H, d, J=8 Hz), 7.40-7.54 (5H, m), 11.21
10 (1H, s), 12.28 (1H, br);

MASS (ES+): m/e 392.

Preparation 20

To a stirred solution of 1H-benzimidazole (500 mg) in N,N-
dimethylformamide (10 mL) was added sodium hydride (186 mg, 60% of
15 oil suspension) at 0°C. After 90 minutes, 4-iodobenzyl bromide was
added to the mixture and the mixture was stirred at ambient
temperature for 1 hour. The reaction mixture was quenched with
saturated ammonium chloride solution, diluted with water and
extracted with ethyl acetate. The organic phase was washed with
20 brine, dried over sodium sulfate and concentrated *in vacuo*. The
residue was triturated with hexane to give Compound (20) as a white
solid. (1.20 g).

¹H-NMR (300 MHz, CDCl₃, δ): 5.31 (2H, s), 6.92 (2x1H, d, J=8.5 Hz),
7.21-7.33 (3H, m), 7.67 (2x1H, d, J=8.5 Hz), 7.84 (1H, m), 7.95 (1H,
25 s);

MASS (ES+): m/e 335.

Preparation 21

Compound (21) was obtained from Compound (20) according to a
manner similar to Preparation 4 (614 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 5.53 (2H, s), 6.50 (1H, d, J=16 Hz),
30 7.15-7.24 (2H, m), 7.32 (2x1H, d, J=8.5 Hz), 7.51 (1H, m), 7.52 (1H,
d, J=16 Hz), 7.61-7.70 (3H, m), 8.43 (1H, s);

MASS (ES+): m/e 279.

Preparation 22

35 Compound (22) was obtained from Compound (21) according to a
manner similar to Preparation 5 (536 mg). The obtained Compound

(22) was used in Example 4.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.45-1.76 (6H, m), 3.52 (1H, m), 3.94 (1H, m), 4.89 (1H, m), 5.53 (2H, s), 6.46 (1H, d, J=16 Hz), 7.16-7.25 (2H, m), 7.33 (2x1H, d, J=8.5 Hz), 7.44 (1H, d, J=16 Hz), 7.51 (1H, m), 7.54 (2x1H, d, J=8.5 Hz), 7.67 (1H, m), 8.42 (1H, s), 11.24 (1H, s);

MASS (ES+): m/e 378.

Preparation 23

Compound (23) was obtained from the by-product obtained in Preparation 10 according to a manner similar to Preparation 4 (150 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.32 (3H, t, J=7 Hz), 3.35 (1H, dd, J=13.5, 7.5 Hz), 3.85 (1H, dd, J=13.5, 7.5 Hz), 4.25 (2H, q, J=7 Hz), 4.28 (1H, dd, J=7.5, 7.5 Hz), 5.05 (1H, d, J=16.5 Hz), 5.11 (1H, d, J=16.5 Hz), 6.35 (1H, d, J=16 Hz), 6.75 (2x1H, dd, J=7.5, 1 Hz), 6.93-7.00 (2H, m), 7.09-7.38 (13H, m), 7.59 (1H, d, J=16 Hz), 7.91 (1H, d, J=8 Hz);

MASS (ES+): m/e 487.

Preparation 24

Compound (24) was obtained from Compound (23) according to a manner similar to Preparation 12 (135 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 3.37 (1H, dd, J=13.5, 7.5 Hz), 3.86 (1H, dd, J=13.5, 7.5 Hz), 4.31 (1H, dd, J=7.5, 7.5 Hz), 5.06 (1H, d, J=15.7 Hz), 5.11 (1H, d, J=15.7 Hz), 6.39 (1H, d, J=15.7 Hz), 6.74 (2x1H, d, J=7 Hz), 6.93-7.02 (2H, m), 7.08-7.33 (11H, m), 7.36 (2x1H, d, J=8 Hz), 7.68 (1H, d, J=15.7 Hz), 7.94 (2x1H, d, J=7.5 Hz);

MASS (ES+): m/e 459.

Preparation 25

Compound (25) was obtained from Compound (24) according to a manner similar to Preparation 9 (140 mg). The obtained Compound (25) was used in Example 5.

¹H-NMR (300 MHz, CDCl₃, δ): 1.55-1.92 (6H, m), 3.35 (1H, dd, J=13.5, 7.5 Hz), 3.64 (1H, m), 3.84 (1H, dd, J=13.5, 7.5 Hz), 3.95 (1H, m), 4.28 (1H, dd, J=7.5, 7.5 Hz), 5.00 (1H, m), 5.04 (1H, d, J=17 Hz), 5.11 (1H, d, J=17 Hz), 6.75 (2x1H, d, J=7 Hz), 6.92-7.00 (2H, m),

7.08-7.37 (14H, m), 7.64 (1H, d, J=15 Hz), 7.90 (1H, d, J=8 Hz);
MASS (ES+): m/e 558.

Preparation 26

Compound (26) was obtained from (3-bromophenyl)acetic acid
5 according to a manner similar to Preparation 6 (6.20 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (3x3H, s), 3.06 (2H, s), 6.37 (1H,
d, J=15.8 Hz), 7.25-7.46 (4H, m), 7.56 (1H, d, J=15.8 Hz);

MASS (ES-): m/e 261.

Preparation 27

10 Compound (27) was obtained from Compound (26) according to a
manner similar to Preparation 7 (6.96 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.48 (3x3H, s), 1.53 (3x3H, s), 3.74 (2H,
s), 6.39 (1H, d, J=15.8 Hz), 6.70 (1H, brs), 7.09-7.20 (2H, m),
7.32-7.52 (6H, m), 7.56 (1H, d, J=15.8 Hz), 8.04 (1H, brs):

15 MASS (ES+): m/e 453.

Preparation 28

Compound (28) was obtained from Compound (27) according to a
manner similar to Preparation 8 (4.19 g).

20 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.58 (2H, s), 6.58 (1H, d, J=16 Hz),
7.42-7.58 (5H, m), 7.58 (1H, d, J=16 Hz), 7.66 (1H, m), 7.74-7.82
(2H, m), 7.87 (1H, brs);

MASS (ES+): m/e 279.

Preparation 29

Compound (29) was obtained from Compound (28) according to a
25 manner similar to Preparation 9 (3.34 g). The obtained Compound
(29) was used in Example 6.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.44-1.76 (6H, m), 3.53 (1H, m), 3.95
(1H, m), 4.20 (2H, s), 4.90 (1H, m), 6.50 (1H, d, J=16 Hz), 7.08-
7.16 (2H, m), 7.32-7.60 (7H, m), 11.25 (1H, s), 12.31 (1H, brs);

30 MASS (ES+): m/e 378.

Preparation 30

Compound (30) was obtained from (4-[(1E)-3-tert-butoxy-3-
oxo-1-propenyl]phenyl)acetic acid according to a manner similar to
Preparation 1 (324 mg).

35 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.48 (3x3H, s), 3.176 (1H, s), 3.723
(1H, s), 5.00 (1H, s), 5.01 (1H, s), 6.51 (1H, d, J=15.7 Hz), 6.82.

(1H, m), 7.19-7.60 (10H, m), 7.66 (2x1H, d, J=8 Hz), 9.45 (0.5H, s), 9.47 (0.5H, s);

MASS (ES+): m/e 429.

Preparation 31

5 Compound (31) was obtained from Compound (30) according to a manner similar to Preparation 8 (216 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.59 (2H, s), 6.57 (1H, d, J=16 Hz), 7.42 (1H, m), 7.51 (2x1H, d, J=7.5 Hz), 7.53 (2x1H, d, J=8.5 Hz), 7.60 (1H, d, J=16 Hz), 7.69-7.88 (6H, m), 7.96 (1H, s);

10 MASS (ES+): m/e 355.

Preparation 32

Compound (32) was obtained from Compound (31) according to a manner similar to Preparation 9 (231 mg). The obtained Compound (32) was used in Example 7.

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.46-1.78 (6H, m), 3.52 (1H, m), 3.95 (1H, m), 4.22 (2H, s), 4.90 (1H, m), 6.48 (1H, d, J=15 Hz), 7.30-7.81 (13H, m), 11.23 (1H, s), 12.38 (1/2H, s), 12.41 (1/2H, s);

MASS (ES+): m/e 454.

Preparation 33

20 Compound (33) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (5.48 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.50 (2H, s), 6.55 (1H, d, J=16 Hz), 7.47 (2H, d, J=8 Hz), 7.59 (1H, d, J=16 Hz), 7.56-7.75 (4H, m), 7.97 (1H, s).

25 MASS (ESI): m/z 357 (M+1).

Preparation 34

Compound (34) was obtained from Compound (33) according to a manner similar to Preparation 9 (557 mg). The Compound (34) was used in Example 8.

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48-1.74 (6H, m), 3.48-3.57 (1H, m), 3.89-4.00 (1H, m), 4.20 (2H, s), 4.90 (1H, brs), 6.47 (1H, d, J=16 Hz), 7.26 (1H, dd, J=2, 8 Hz), 7.34-7.56 (6H, m), 7.68 (1H, brs).

MASS (ESI): m/z 456 (M+1).

Preparation 35

35 To a mixture of Compound (34) (200 mg), 4-acetylphenylboronic acid (167 mg) and

dichlorobis(triphenylphosphine)palladium(II) (10.7 mg) in dioxane (10 mL) was added 2M sodium carbonate (2.5 mL), and the mixture was heated at 90°C for 5 hours. After cooling, the reaction mixture was partitioned between ethyl acetate and water. The inorganic layer was separated and acidified with 1N hydrochloric acid. The resulting precipitate was collected by filtration, and washed with water and ethyl acetate to give Compound (35) (193 mg). The Compound (35) was used in Example 9.

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.63 (3H, s), 4.57 (2H, s), 6.56 (1H, d, J=16 Hz), 7.52 (2H, d, J=8 Hz), 7.59 (1H, d, J=16 Hz), 7.74 (2H, d, J=8 Hz), 7.85-7.92 (4H, m), 8.03-8.10 (3H, m);

MASS (ESI): m/z 397 (M+1).

Preparation 36

Compound (36) was obtained from Compound (34) according to a manner similar to Preparation 35 (163 mg). The Compound (36) was used in Example 10.

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.37 (2H, s), 6.53 (1H, d, J=16 Hz), 7.12-7.16 (1H, m), 7.45-7.71 (9H, m), 7.81 (1H, s);

MASS (ESI): m/z 361 (M+1).

Preparation 37

Compound (37) was obtained from Compound (34) according to a manner similar to Preparation 35 (183 mg). The Compound (37) was used in Example 11.

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.55 (2H, s), 6.56 (1H, d, J=16 Hz), 7.40-8.01 (11H, m);

MASS (ESI): m/z 361 (M+1).

Preparation 38

Compound (38) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (2.07 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.47 (2H, s), 6.54 (1H, d, J=16 Hz), 7.46 (2H, d, J=8 Hz), 7.58 (1H, d, J=16 Hz), 7.70 (2H, d, J=8 Hz), 7.77 (1H, d, J=8 Hz), 7.83 (1H, d, J=8 Hz), 8.26 (1H, s);

MASS (ESI): m/z 304 (M+1)...

Preparation 39

Compound (39) was obtained from Compound (38) according to a manner similar to Preparation 9 (2.14 g). The Compound (39) was

used in Examples 12 and 18.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.49–1.72 (6H, m), 3.50–3.56 (1H, m),
3.91–3.99 (1H, m), 4.26 (2H, s), 4.91 (1H, brs), 6.47 (1H, d, J=16
5 Hz), 7.37 (2H, d, J=8 Hz), 7.46 (1H, d, J=16 Hz), 7.51–7.57 (3H, m),
7.65 (1H, d, J=8 Hz), 8.32 (1H, brs);

MASS (ESI): m/z 401 (M-1).

Preparation 40

Compound (40) was obtained from Compound (6) according to
manners similar to Preparations 7 and 8 (955 mg). The Compound
10 (40) was used in Example 13.

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.50 (2H, s), 6.55 (1H, d, J=16 Hz),
7.32–7.39 (1H, m), 7.47 (2H, d, J=8 Hz), 7.59 (1H, d, J=16 Hz),
7.62 (1H, dd, J=2, 8 Hz), 7.72 (2H, d, J=8 Hz), 7.74–7.79 (1H, m);
MASS (ESI): m/z 297 (M+1).

Preparation 41

Compound (41) was obtained from Compound (6) according to
manners similar to Preparations 7 and 8 (1.02 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.52 (2H, s), 6.55 (1H, d, J=16 Hz),
7.48 (2H, d, J=8 Hz), 7.50 (1H, dd, J=2, 8 Hz), 7.59 (1H, d, J=16
20 Hz), 7.72 (2H, d, J=8 Hz), 7.75 (1H, d, J=8 Hz), 7.85 (1H, J=2 Hz);
MASS (ESI): m/z 313 (M+1).

Preparation 42

Compound (42) was obtained from Compound (41) according to a
manner similar to Preparation 9 (839 mg). The Compound (42) was
25 used in Example 14.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.49–1.71 (6H, m), 3.49–3.57 (1H, m),
3.88–4.02 (1H, m), 4.20 (2H, s), 4.88–4.93 (1H, m), 6.43–6.52 (1H,
m), 7.12–7.18 (1H, m), 7.36 (2H, d, J=8 Hz), 7.46 (1H, d, J=16 Hz),
7.53 (2H, d, J=8 Hz), 7.53–7.60 (1H, m), 7.64 (1H, d, J=8 Hz);
30 MASS (ESI): m/z 412 (M+1).

Preparation 43

Compound (43) was obtained from Compound (6) according to
manners similar to Preparations 7 and 8 (411 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.78–2.84 (4H, m), 3.13–3.20 (4H, m),
35 4.54 (2H, s), 6.55 (1H, d, J=16 Hz), 7.16–7.76 (8H, m);
MASS (ESI): m/z 377 (M+1).

Preparation 44

Compound (44) was obtained from Compound (43) according to a manner similar to Preparation 9 (23 mg). The Compound (44) was used in Example 15.

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.54–1.87 (6H, m), 2.36 (3H, s), 2.58–2.64 (4H, m), 3.12–3.20 (4H, m), 3.58–3.66 (1H, m), 3.92–4.04 (1H, m), 4.15 (2H, s), 5.02–5.10 (1H, m), 6.92–7.60 (9H, m);
MASS (ESI): m/z 476 (M+1).

Preparation 45

- 10 Compound (45) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (358 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.16–3.22 (4H, m), 3.76–3.83 (4H, m), 4.53 (2H, s), 6.56 (1H, d, $J=16$ Hz), 7.34–7.76 (8H, m);
MASS (ESI): m/z 364 (M+1).

- 15 Preparation 46

Compound (46) was obtained from Compound (45) according to a manner similar to Preparation 9 (59 mg). The Compound (46) was used in Example 16.

- 20 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.51–1.87 (6H, m), 3.05–3.13 (4H, m), 3.57–3.64 (1H, m), 3.83–3.91 (4H, m), 3.95–4.04 (1H, m), 4.15 (2H, s), 5.04–5.13 (1H, m), 6.91–7.61 (9H, m);
MASS (ESI): m/z 463 (M+1).

Preparation 47

- 25 Compound (47) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (517 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.57–1.69 (2H, m), 1.83–1.94 (4H, m), 3.37–3.50 (4H, m), 4.52 (2H, s), 6.48–6.58 (1H, m), 7.40–7.74 (8H, m);
MASS (ESI): m/z 362 (M+1).

- 30 Preparation 48

Compound (48) was obtained from Compound (47) according to a manner similar to Preparation 9 (47 mg). The Compound (48) was used in Example 17.

- 35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.50–1.87 (12H, m), 3.05–3.11 (4H, m), 3.57–3.66 (1H, m), 3.94–4.05 (1H, m), 4.14 (2H, s), 5.02–5.12 (1H, m), 6.94–7.48 (9H, m);

MASS (ESI): m/z 461 (M+1).

Preparation 49

Compound (49) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (1.17 g).

5 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.90 (3H, s), 4.49 (2H, s), 6.54 (1H, d, J=16 Hz), 7.47 (2H, d, J=8 Hz), 7.59 (1H, d, J=16 Hz), 7.72 (2H, d, J=8 Hz), 7.78 (1H, d, J=8 Hz), 8.00 (1H, dd, J=2, 8 Hz), 8.24 (1H, brs);

MASS (ESI): m/z 337 (M+1).

10 Preparation 50

Compound (50) was obtained from Compound (49) according to a manner similar to Preparation 9 (1.30 g). The Compound (50) was used in Example 19.

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48-1.74 (6H, m), 3.48-3.58 (1H, m), 3.85 (3H, s), 3.87-4.00 (1H, m), 4.25 (2H, brs), 4.90 (1H, brs), 6.47 (1H, d, J=16 Hz), 7.35-7.65 (6H, m), 7.74-8.16 (2H, m);

MASS (ESI): m/z 436 (M+1).

Preparation 51

20 To a solution of Compound (50) (299 mg) in dioxane (7 mL) was added 1N sodium hydroxide (2.1 mL). After stirring at 80°C for 1 hour, the reaction mixture was added water (25 mL) and acidified with 1N hydrochloric acid (to pH 3-4). A resulting precipitate was collected by filtration and washed with water to give Compound (51) (255 mg). The Compound (51) was used in Example 20.

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.49-1.74 (6H, m), 3.47-3.59 (1H, m), 3.87-4.01 (1H, m), 4.24 (2H, s), 4.90 (1H, brs), 6.47 (1H, d, J=16 Hz), 7.38 (2H, d, J=8 Hz), 7.43-7.62 (4H, m), 7.73-8.14 (2H, m);

MASS (ESI): m/z 422 (M+1).

Preparation 52

30 Compound (52) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (384 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.68 (3H, s), 4.55 (2H, s), 6.55 (1H, d, J=16 Hz), 7.50 (2H, d, J=8 Hz), 7.59 (1H, d, J=16 Hz), 7.73 (2H, d, J=8 Hz), 7.81 (1H, d, J=8 Hz), 8.04 (1H, dd, J=2, 8 Hz), 8.27 (1H, s);

35 MASS (ESI): m/z 321 (M+1).

Preparation 53

Compound (53) was obtained from Compound (52) according to a manner similar to Preparation 9 (394 mg). The Compound (53) was used in Example 21.

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.47–1.73 (6H, m), 2.61 (3H, s), 3.48–3.57 (1H, m), 3.90–4.01 (1H, m), 4.25 (2H, brs), 4.90 (1H, brs), 6.47 (1H, d, $J=16$ Hz), 7.36–8.23 (8H, m);
MASS (ESI): m/z 418 (M-1).

Preparation 54

- 10 Compound (54) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (335 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.58 (2H, s), 6.53 (1H, d, $J=16$ Hz), 7.30–7.36 (1H, m), 7.47 (2H, d, $J=8$ Hz), 7.54–7.73 (5H, m);
MASS (ESI): m/z 357 (M+1).

- 15 Preparation 55

Compound (55) was obtained from Compound (54) according to a manner similar to Preparation 9 (364 mg). The Compound (55) was used in Example 22.

- 20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.48–1.74 (6H, m), 3.48–3.58 (1H, m), 3.89–4.02 (1H, m), 4.24 (2H, s), 4.90 (1H, brs), 6.47 (1H, d, $J=16$ Hz), 7.05–7.12 (1H, m), 7.33–7.58 (7H, m);
MASS (ESI): m/z 456 (M+1).

Preparation 56

- 25 Compound (56) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (805 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.24 (3H, s), 1.27 (3H, s), 3.02–3.15 (1H, m), 4.54 (2H, s), 6.56 (1H, d, $J=16$ Hz), 7.33–7.75 (8H, m);
MASS (ESI): m/z 321 (M+1).

Preparation 57

- 30 Compound (57) was obtained from Compound (56) according to a manner similar to Preparation 9 (70 mg). The Compound (57) was used in Example 23.

- 35 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.22 (3H, s), 1.24 (3H, s), 1.48–1.73 (6H, m), 2.89–3.03 (1H, m), 3.48–3.57 (1H, m), 3.89–4.00 (1H, m), 4.17 (2H, s), 4.90 (1H, brs), 6.47 (1H, d, $J=16$ Hz), 7.01 (1H, d, $J=8$ Hz), 7.20–7.57 (7H, m);

MASS (ESI): m/z 420 (M+1).

Preparation 58

Compound (58) was obtained from Compound (6) according to manners similar to Preparations 7 and 8 (925 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.50 (2H, s), 5.37 (2H, s), 6.54 (1H, d, $J=16$ Hz), 7.13-7.75 (13H, m);

MASS (ESI): m/z 385 (M+1).

Preparation 59

10 Compound (59) was obtained from Compound (58) according to a manner similar to Preparation 9. The Compound (59) was used in Example 24.

Preparation 60

Compound (60) was obtained according to a manner similar to Preparation 20 (1.25 g).

15 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 2.56 (3H, s), 5.27 (2H, s), 6.79 (2x1H, d, $J=8.4$ Hz), 7.15-7.29 (3H, m), 7.63 (2x1H, d, $J=8.4$ Hz), 7.73 (1H, m);

MASS (ES+): m/e 349.

Preparation 61

20 Compound (61) was obtained from Compound (60) according to a manner similar to Preparation 4 (625 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 2.57 (3H, s), 5.36 (2H, s), 6.40 (1H, d, $J=16.2$ Hz), 7.07 (2x1H, d, $J=8.5$ Hz), 7.20-7.29 (3H, m), 7.48 (2x1H, d, $J=8.5$ Hz), 7.64 (1H, d, $J=16.2$ Hz), 7.71 (1H, m);

25 MASS (ES+): m/e 293.

Preparation 62

Compound (62) was obtained from Compound (61) according to a manner similar to Preparation 9 (556 mg). The Compound (62) was used in Example 25.

30 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.46-1.75 (6H, m), 2.52 (3H, s), 3.52 (1H, m), 3.94 (1H, m), 4.89 (1H, m), 5.50 (2H, s), 6.46 (1H, d, $J=16$ Hz), 7.11-7.20 (4H, m), 7.38-7.62 (5H, m);

MASS (ES+): m/e 392.

Preparation 63

35 To a stirred solution of 1-phenylcyclopropanecarboxylic acid (3.25 g) in acetic acid (30 mL) were added periodic acid (959 mg),

iodine (2.03 g), concentrated H₂SO₄ (0.6 mL) and water (4 mL), and the mixture was stirred at 70°C for 12 hours. Water (100 mL) was added to the mixture and the precipitated solid was collected by filtration and washed with water to give Compound (63) (4.31 g).

5 ¹H-NMR (300 MHz, CDCl₃, δ): 1.23 (2H, ddd, J=7, 4, 4 Hz), 1.67 (2H, ddd, J=7, 4, 4 Hz), 7.09 (2x1H, d, J=8.4 Hz), 7.63 (2x1H, d, J=8.4 Hz);

MASS (ES⁻): m/e 287.

Preparation 64

10 Compound (64) was obtained from Compound (63) according to a manner similar to Preparation 6 (432 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.16 (2H, ddd, J=7, 4, 4 Hz), 1.45 (2H, ddd, J=7, 4, 4 Hz), 1.48 (3x3H, s), 6.49 (1H, d, J=15.7 Hz), 7.35 (2x1H, d, J=8 Hz), 7.53 (1H, d, J=15.7 Hz), 7.61 (2x1H, d, J=8 Hz);

15 MASS (ES⁻): m/e not determined.

Preparation 65

Compound (65) was obtained from Compound (64) according to a manner similar to Preparation 7 (1.67 g).

20 ¹H-NMR (300 MHz, CDCl₃, δ): 1.20 (2H, m), 1.45 (3x3H, s), 1.51 (3x3H, s), 1.74 (2H, m), 6.40 (1H, d, J=15.7 Hz), 6.63 (1H, br-s), 6.98-7.62 (10H, m);

MASS (ES⁺): m/e 479.

Preparation 66

25 Compound (66) was obtained from Compound (65) according to a manner similar to Preparation 8 (710 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.71 (2H, m), 1.95 (2H, m), 6.59 (1H, d, J=15.8 Hz), 7.40-7.55 (4H, m), 7.63 (1H, d, J=15.8 Hz), 7.64-7.72 (2H, m), 7.76 (2x1H, d, J=8.4 Hz);

MASS (ES⁺): m/e 305.

30 Preparation 67

Compound (67) was obtained from Compound (66) according to a manner similar to Preparation 9 (647 mg).

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.39 (2H, m), 1.48-1.76 (8H, m), 3.34 (1H, m), 3.54 (1H, m), 3.96 (1H, m), 4.91 (2H, s), 6.51 (1H, d, J=16 Hz), 7.07-7.15 (2H, m), 7.31-7.43 (3H, m), 7.46-7.63 (4H, m);

MASS (ES⁺): m/e 404.

Preparation 68

To a stirred solution of 4-bromo-2-nitroaniline (1.37 g) in dioxane (20 mL) were added 4-fluorophenylboronic acid (1.06 g), PdCl₂(PPh₃)₂ (133 mg), and 2M sodium carbonate solution (12.7 ml), and the mixture was stirred at 100°C for 2 hours. The solvent was evaporated and the residue was partitioned between ethyl acetate and water. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The resulting solid was recrystallized from toluene to give Compound (68) (1.13 g) as a white solid.

¹H-NMR (300 MHz, CDCl₃, δ): 6.18 (2H, br-s), 6.89 (1H, d, J=8.7 Hz), 7.12 (2x1H, dd, J=8.7, 8.7 Hz), 7.51 (2x1H, dd, J=8.7, 5 Hz), 7.59 (1H, dd, J=8.7, 2.2 Hz), 8.31 (1H, d, J=2.2 Hz);

MASS (ES-) m/e 231.

15 Preparation 69

To a stirred solution of Compound (68) (1.08 g) in EtOH (15 mL) was added tin (II) chloride (1.32 g). The mixture was stirred at 100°C for 6 hours. The solvent was evaporated to the half volume and the residue was basified with 1N-NaOH to pH 9 and extracted with ethyl acetate. The organic phase was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was triturated with ethyl acetate to give Compound (69) (800 mg) as an orange powder.

¹H-NMR (300 MHz, CDCl₃, δ): 3.45 (2x2H, br-s), 6.76 (2x1H, d, J=8.3 Hz), 6.89 (1H, d, J=2.1 Hz), 6.91 (1H, dd, J=8.3, 2.1 Hz), 7.06 (2x1H, dd, J=8.8, 8.8 Hz), 7.46 (2x1H, dd, J=8.8, 5.4 Hz);

MASS (ES+): m/e 203.

Preparation 70

Compound (70) was obtained from Compound (69) according to a manner similar to Preparation 7 (552 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (3x3H, s), 3.81 (2H, s), 6.39 (1H, sd, J=16 Hz), 6.80-7.62 (15H, m);

MASS (ES+): m/e 446.

Preparation 71

Compound (71) was obtained from Compound (70) according to a manner similar to Preparation 8 (375 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.58 (2H, s), 6.56 (1H, d, $J=16$ Hz), 7.23-7.86 (11H, m), 7.94 (1H, s);

MASS (ES+): m/e 373.

Preparation 72

5 Compound (72) was obtained from Compound (71) according to a manner similar to Preparation 9 (321 mg). The Compound (71) was used in Example 26.

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.45-1.75 (6H, m), 3.52 (1H, m), 3.95 (1H, m), 4.22 (2H, s), 4.90 (1H, m), 6.48 (1H, d, $J=15.8$ Hz), 7.22-10 7.80 (12H, m), 11.23 (1H, br-s), 12.40 (H, br-s);

MASS (ES+): m/e 472.

Preparation 73

Compound (73) was obtained from Compound (6) according to a manner similar to Preparation 7 (228 mg).

15 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.53 (3x3H, s), 3.75-3.87 (4H, br), 6.41 (1H, d, $J=15.7$ Hz), 6.81-7.54 (13H, m), 7.58 (1H, d, $J=15.7$ Hz);

MASS (ES+): m/e 429.

Preparation 74

20 Compound (74) was obtained from Compound (73) according to a manner similar to Preparation 8 (165 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.59 (2H, s), 6.59 (1H, d, $J=16$ Hz), 7.38-7.55 (5H, m), 7.59 (1H, d, $J=16$ Hz), 7.67 (1H, m), 7.73 (2x1H, d, $J=7.5$ Hz), 7.78-7.85 (2H, m), 7.87 (1H, br), 7.96 (1H, br-s);

25 MASS (ES+): m/e 355.

Preparation 75

Compound (75) was obtained from Compound (74) according to a manner similar to Preparation 9 (185 mg). The Compound (75) was used in Example 27.

30 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.46-1.76 (6H, m), 3.53 (1H, m), 3.94 (1H, m), 4.23 (2H, s), 4.90 (1H, m), 6.50 (1H, d, $J=16$ Hz), 7.28-7.52 (10H, m), 7.57 (1H, m), 7.66 (2x1H, d, $J=7.5$ Hz), 11.25 (1H, br), 12.38 (1H, br);

MASS (ES+): m/e 454.

35 Preparation 76

Compound (76) was obtained according to a manner similar to

Preparation 35 (320 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.24 (3H, s), 7.16 (1H, d, $J=8.8$ Hz), 7.68 (2H, s), 7.85-8.07 (5H, m), 8.34 (1H, d, $J=2.5$ Hz);

MASS (ES-): m/e 291.

5 Preparation 77

To a stirred solution of 4'-(methylsulfonyl)-3-nitro-1,1'-biphenyl-4-ylamine (305 g) in EtOH (15 mL) were added iron powder (583 mg), NH_4Cl (56 mg) and water (1 mL). The mixture was refluxed for 5 hours. The iron powder was filtered off and the filtrate was
10 evaporated in vacuo. The residue was partitioned between CHCl_3 and saturated NaHCO_3 solution. The organic phase was washed with brine, dried over Na_2SO_4 , and concentrated in vacuo to give Compound (77) (150 mg) as an orange powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.44 (3H, s), 4.45-5.20 (4H, m), 6.65
15 (1H, br), 6.82-7.22 (2H, m), 7.62-8.22 (4H, m);

MASS (ES+): m/e 263.

Preparation 78

Compound (78) was obtained from Compound (77) according to a manner similar to Preparation 7 (200 mg).

20 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (3x3H, s), 3.07 (3x1/2H, s), 3.08 (3x1/2H, s), 3.83 (2H, br-s), 6.39 (1H, d, $J=16$ Hz), 6.84-7.10 (2H, m), 7.29 (1H, m), 7.42 (2H, m), 7.53-7.73 (5H, m), 7.95 (2H, m);

MASS (ES+): m/e 507.

Preparation 79

25 Compound (79) was obtained from Compound (78) according to a manner similar to Preparation 8 (162 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.28 (3H, s), 4.59 (2H, s), 6.57 (1H, d, $J=15.8$ Hz), 7.52 (2x1H, d, $J=8.2$ Hz), 7.59 (1H, d, $J=15.8$ Hz), 7.74 (2x1H, d, $J=8.2$ Hz), 7.86-7.90 (2H, m), 7.95-8.10 (5H, m);

30 MASS (ES+): m/e 432.

Preparation 80

Compound (80) was obtained from Compound (79) according to a manner similar to Preparation 9 (190 mg). The Compound (80) was used for Example 28.

35 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.46-1.76 (6H, m), 3.25 (3H, s), 3.53 (1H, m), 3.94 (1H, m), 4.24 (2H, m), 4.90 (1H, m), 6.48 (1H, d,

J=16 Hz), 7.39 (2x1H, d, J=8 Hz), 7.42-8.02 (10H, m), 11.23 (0.5H, br), 12.50 (0.5H, br);

MASS (ES+): m/e 532.

Preparation 81

5 Compound (81) was obtained from Compound (6) according to a manner similar to Preparation 7 (137 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.33 (3H, t, J=7.5 Hz), 4.26 (2H, q, J=7.5 Hz), 4.38 (2H, s), 6.41 (1H, d, J=16 Hz), 7.18 (1H, dd, J=8, 4.5 Hz), 7.38 (2x1H, d, J=8 Hz), 7.51 (2x1H, d, J=8 Hz), 7.65 (1H, 10 d, J=16 Hz), 7.99-8.10 (2H, m);

MASS (ES+): m/e 308.

Preparation 82

Compound (82) was obtained from Compound (81) according to a manner similar to Preparation 12 (362 mg).

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.23 (2H, s), 6.50 (1H, d, J=16 Hz), 7.17 (1H, dd, J=8, 5 Hz), 7.38 (2x1H, d, J=8 Hz), 7.56 (1H, d, J=16 Hz), 7.65 (2x1H, d, J=8 Hz), 7.91 (1H, br), 8.25 (1H, br);

MASS (ES+): m/e 280.

Preparation 83

20 Compound (83) was obtained from Compound (82) according to a manner similar to Preparation 9 (312 mg). The Compound (83) was used in Example 29.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.46-1.76 (6H, m), 3.53 (1H, m), 3.95 (1H, m), 4.21 (2H, br-s), 4.90 (1H, m), 6.48 (1H, d, J=15.5 Hz), 25 7.17 (1H, dd, J=8, 4.5 Hz), 7.39 (2x1H, d, J=8 Hz), 7.46 (1H, d, J=15.5 Hz), 7.54 (2x1H, br-d, J=8 Hz), 7.88 (1H, m), 8.27 (1H, m), 11.24 (1H, br-s), 12.58 (0.5H, br), 13.00 (0.5H, br);

MASS (ES+): m/e 379.

Preparation 84

30 Compound (84) was obtained from Compound (6) according to a manner similar to Preparation 7 (9.77 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.26 (3H, t, J=7 Hz), 3.74 (2H, s), 4.19 (2H, q, J=7 Hz), 6.54 (2H, s), 6.62 (1H, d, J=16 Hz), 6.77 (1H, d, J=8.8 Hz), 7.41 (2x1H, d, J=8 Hz), 7.65 (1H, d, J=16 Hz), 7.70 35 (2x1H, d, J=8 Hz), 7.85 (1H, dd, J=8.8, 2.5 Hz), 8.20 (1H, d, J=2.5 Hz), 9.46 (1H, s);

MASS (ES+): m/e 370.

Preparation 85

Compound (85) was obtained from Compound (84) according to a manner similar to Preparation 8 (6.83 g).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.25 (3H, t, $J=7$ Hz), 4.18 (2H, q, $J=7$ Hz), 4.44 (2H, s), 6.64 (1H, d, $J=16$ Hz), 7.46 (2x1H, d, $J=8$ Hz), 7.64 (1H, d, $J=16$ Hz), 7.73 (2x1H, d, $J=8$ Hz), 7.80 (1H, d, $J=9$ Hz), 8.20 (1H, dd, $J=9$, 2.2 Hz), 8.50 (1H, d, $J=2.2$ Hz);

MASS (ES+): m/e 352.

10 Preparation 86

Compound (86) was obtained from Compound (85) according to a manner similar to Preparation 77 (872 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.31 (3H, br-t, $J=7$ Hz), 4.15-4.36 (4H, m), 6.26 (1H, br-d, $J=16$ Hz), 6.64 (1H, m), 6.78 (1H, m), 7.20-7.40
15 (5H, m), 7.49 (1H, br-d, $J=16$ Hz);

MASS (ES+): m/e 322.

Preparation 87

To a stirred suspension of Compound (86) (303 mg) in dioxane (10 mL) were added di-tert-butylidicarbonate (618 mg) in dioxane
20 (3mL) and then 1N-NaOH (2.8 mL), and the mixture was stirred at ambient temperature for 12 hours. The solvent was evaporated in vacuo and the residue was partitioned between ethyl acetate and water. The organic phase was washed with brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue was purified by preparative
25 thin layer chromatography (hexane: ethyl acetate = 1: 1) to give Compound (87) (379 mg) as a pale brown amorphous (379 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.33 (3H, t, $J=7$ Hz), 1.52 (4.5H, s), 1.53 (4.5H, s), 1.58 (4.5H, s), 1.60 (4.5H, s), 4.25 (2H, q, $J=7$ Hz), 4.61 (2H, s), 6.39 (1H, d, $J=16$ Hz), 6.58 (1H, br-d, $J=6$ Hz),
30 7.09 (0.5H, dd, $J=8.8$, 2.2 Hz), 7.23-7.30 (2H, m), 7.42-7.48 (2.5H, m), 7.567.68 (2H, m), 7.77 (0.5H, d, $J=8.8$ Hz), 8.24 (0.5H, br);

MASS (ES+): m/e 522.

Preparation 88

To a stirred solution of Compound (87) (360 mg) in methanol
35 (5 mL) was added 1N-NaOH solution (1.4 mL). The mixture was stirred at ambient temperature for 2.5 hours. The reaction mixture

was neutralized by 1N-HCl solution, and the solvent was evaporated in vacuo. The residue was partitioned between ethyl acetate and water. The organic phase was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by preparative
5 thin layer chromatography (chloroform :methanol = 10: 1) to give Compound (88) (57 mg) as an orange powder and a methyl ester of Compound (88) as a by-product.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48 (3x3H, s), 4.16 (2H, s), 6.49 (1H, d, J=16 Hz), 7.12 (1H, br), 7.31-7.42 (3H, m), 7.55 (1H, d, J=16
10 Hz), 7.60-7.72 (3H, m), 9.25 (1H, br), 12.12 (1H, br);

MASS (ES+): m/e 394.

Preparation 89

Compound (89) was obtained from the methyl ester of Compound (88) according to a manner similar to Preparation 12 (245 mg).

15 [The Compound (89) is similar to Compound (88).]

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48 (3x3H, s), 4.22 (2H, s), 6.50 (1H, d, J=16 Hz), 7.19 (1H, br-d, J=8.5 Hz), 7.34-7.42 (3H, m), 7.56 (1H, d, J=16 Hz), 7.65 (2x1H, d, J=8.5 Hz), 7.72 (1H, br-s), 9.30 (1H, br-s), 12.38 (1H, br);

20 MASS (ES+): m/e 394.

Preparation 90

Compound (90) was obtained from Compound (88) according to a manner similar to Preparation 9 (230 mg). The Compound (90) was used for Example 30.

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48 (3x3H, s), 1.48-1.75 (6H, m), 3.53 (1H, m), 3.94 (1H, m), 4.15 (2H, s), 4.90 (1H, m), 6.47 (1H, br-d, J=16 Hz), 7.06-7.414 (4H, m), 7.46 (1H, d, J=16 Hz), 7.53 (2x1H, br-d, J=8.5 Hz), 7.67 (1H, m), 9.15 (1/3H, br-s), 9.26 (2/3H, br-s), 11.22 (1H, br-s), 12.10 (2/3H, br-s), 12.13 (1/3H, br-s);

30 MASS (ES+): m/e 493.

Preparation 91

To a stirred solution of Compound (86) (150 mg) in DMF (2 mL) were added butyric acid (49 mg), HOBT (76 mg), and EDCI hydrochloride (107 mg), and the resulting mixture was stirred at
35 ambient temperature for 12 hours. The reaction mixture was diluted with ethyl acetate and washed successively with water, saturated

NaHCO₃ solution and brine. The organic phase was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by preparative thin layer chromatography (chloroform: methanol = 10: 1) to give Compound (91) (30 mg) as a pale yellow amorphous.

5 ¹H-NMR (300 MHz, CDCl₃, δ): 0.98 (3H, t, J=7.3 Hz), 1.33 (3H, t, J=7 Hz), 1.73 (2H, tq, J=7.3, 7.3 Hz), 2.33 (2H, t, J=7.3 Hz), 4.09 (2H, s), 4.24 (2H, q, J=7 Hz), 6.31 (1H, d, J=16 Hz), 7.05 (1H, br-d, J=8 Hz), 7.15 (2x1H, d, J=8 Hz), 7.32 (2x1H, d, J=8 Hz), 7.36 (1H, d, J=8 Hz), 7.53 (1H, s), 7.58 (1H, s);

10 MASS (ES+): m/e 392.

Preparation 92

Compound (92) was obtained from Compound (91) according to a manner similar to Preparation 11 (28 mg).

¹H-NMR (300 MHz, CD₃OD, δ): 1.00 (3H, t, J=7.3 Hz), 1.73 (2H, tq, J=7.3, 7.3 Hz), 2.42 (2H, t, J=7.3 Hz), 4.58 (2H, s), 6.52 (1H, d, J=16 Hz), 7.47-7.58 (3H, m), 7.64-7.74 (4H, m), 8.29 (1H, s);

15 MASS (ES+): m/e 364.

Preparation 93

20 Compound (93) was obtained from Compound (92) according to a manner similar to Preparation 9 (19 mg). The Compound (93) was used in Example 31.

¹H-NMR (300 MHz, CD₃OD-CDCl₃, δ): 1.00 (3H, t, J=7.4 Hz), 1.52-1.96 (8H, m), 2.35 (2H, t, J=7.5 Hz), 3.65 (1H, m), 4.03 (1H, m), 4.11 (2H, s), 5.03 (1H, m), 6.24 (1H, m), 7.06-7.29 (4H, m), 7.38 (2x1H, d, J=8.5 Hz), 7.48 (1H, m), 7.82 (1H, s);

25 MASS (ES+): m/e 463.

Preparation 94

Compound (94) was obtained from Compound (6) according to a manner similar to Preparation 7 (641 mg).

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.26 (3H, t, J=7 Hz), 3.76 (2H, s), 4.19 (2H, q, J=7 Hz), 6.62 (1H, d, J=16 Hz), 6.64 (1H, dd, J=8.5, 8.5 Hz), 7.12 (2H, s), 7.40 (2x1H, d, J=8.5 Hz), 7.45 (1H, dd, J=8.5, 1 Hz), 7.65 (1H, d, J=16 Hz), 7.69 (2x1H, d, J=8.5 Hz), 7.91 (1H, dd, J=8.5, 1 Hz), 9.57 (1H, s);

35 MASS (ES+): m/e 370.

Preparation 95

Compound (95) was obtained from Compound (94) according to a manner similar to Preparation 8 (512 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.25 (3H, t, J=7 Hz), 4.18 (2H, q, J=7 Hz), 4.45 (2H, s), 6.62 (1H, d, J=16 Hz), 7.46 (2x1H, d, J=8.3 Hz),
5 7.49 (1H, dd, J=8, 8 Hz), 7.63 (1H, d, J=16 Hz), 7.71 (2x1H, d, J=8.3 Hz), 8.11 (1H, d, J=8 Hz), 8.20 (1H, d, J=8 Hz);

MASS (ES+): m/e 352.

Preparation 96

10 Compound (96) was obtained from Compound (95) according to a manner similar to Preparation 12 (119 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.34 (2H, s), 6.49 (1H, d, J=16 Hz), 7.36 (1H, dd, J=8, 8 Hz), 7.41 (2x1H, d, J=8 Hz), 7.55 (1H, d, J=16 Hz), 7.64 (2x1H, d, J=8 Hz), 8.04 (1H, d, J=8 Hz), 8.09 (1H, d, J=8 Hz), 13.26 (1H, br-s);

15 MASS (ES+): m/e 324.

Preparation 97

Compound (97) was obtained from Compound (96) according to a manner similar to Preparation 9 (70 mg). The Compound (97) was used in Example 32.

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.45-1.77 (6H, m), 3.53 (1H, m), 3.95 (1H, m), 4.34 (2H, s), 4.90 (1H, m), 6.47 (1H, d, J=16 Hz), 7.37 (1H, dd, J=8, 8 Hz), 7.42 (2x1H, d, J=8 Hz), 7.46 (1H, d, J=16 Hz), 7.53 (2x1H, d, J=8 Hz), 8.05 (1H, d, J=8 Hz), 8.10 (1H, d, J=8 Hz);
MASS (ES+): m/e 423.

25 Preparation 98

Compound (98) was obtained from Compound (6) according to a manner similar to Preparation 7 (160 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.26 (3H, t, J=7 Hz), 3.75 (2H, s), 4.19 (2H, q, J=7 Hz), 6.47 (2H, br-s), 6.62 (1H, d, J=16 Hz), 6.70
30 (1H, d, J=6 Hz), 7.40 (2x1H, d, J=8 Hz), 7.65 (1H, d, J=16 Hz), 7.69 (2x1H, d, J=8 Hz), 7.91 (1H, d, J=6 Hz), 8.18 (1H, s), 9.61 (1H, s);

MASS (ES+): m/e 326.

Preparation 99

35 Compound (99) was obtained from Compound (6) according to a manner similar to Preparation 8 (91 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.26 (2H, s), 6.50 (1H, d, J=16 Hz), 7.38 (2x1H, d, J=8 Hz), 7.48 (1H, m), 7.56 (1H, d, J=16 Hz), 7.65 (2x1H, d, J=8 Hz), 8.25 (1H, d, J=3 Hz), 8.81 (1H, s), 12.75 (1H, br);

5 MASS (ES+): m/e 280.

Preparation 100

Compound (100) was obtained from Compound (99) according to a manner similar to Preparation 9 (33 mg). The Compound (100) was used for Example 33.

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.40-1.80 (6H, m), 3.52 (1H, m), 3.96 (1H, m), 4.26 (2H, s), 4.91 (1H, m), 6.50 (1H, br-d, J=15.5 Hz), 7.35-7.60 (6H, m), 8.22 (1H, d, J=5.5 Hz), 8.81 (1H, s);

MASS (ES+) m/e 379.

Preparation 101

15 Compound (101) was obtained from Compound (6) according to a manner similar to Preparation 7 (520 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.26 (3H, t, J=7 Hz), 3.11 (3H, s), 3.31 (3H, s), 3.62 (2H, s), 4.19 (2H, q, J=7 Hz), 6.54-6.68 (3H, m), 7.41 (2x1H, d, J=8 Hz), 7.58-7.70 (3H, m), 8.66 (1H, s);

20 MASS (ES+): m/e 387.

Preparation 102

Compound (102) was obtained from Compound (101) according to a manner similar to Preparation 8 (416 mg).

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.22 (3H, s), 3.40 (3H, s), 4.08 (2H, s), 6.49 (1H, d, J=16 Hz), 7.33 (2x1H, d, J=8.3 Hz), 7.55 (1H, d, J=16 Hz), 7.64 (2x1H, d, J=8.3 Hz), 12.39 (1H, s), 13.48 (1H, s);

MASS (ES+): m/e 341.

Preparation 103

30 Compound (103) was obtained from Compound (102) according to a manner similar to Preparation 9 (254 mg). The Compound (103) was used in Example 34.

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.46-1.76 (6H, m), 3.22 (3H, s), 3.40 (3H, s), 3.53 (1H, m), 3.95 (1H, m), 4.07 (2H, s), 4.90 (1H, m), 6.47 (1H, d, J=16 Hz), 7.33 (2x1H, d, J=7.5 Hz), 7.46 (1H, d, J=16 Hz), 7.53 (2x1H, d, J=7.5 Hz), 11.23 (1H, br-s), 13.47 (1H, br-s);

MASS (ES-): m/e 438.

Preparation 104

Compound (104) was obtained from Compound (85) according to a manner similar to Preparation 12 (1.54 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.38 (2H, s), 6.52 (1H, d, J=16.2 Hz),
5 7.42 (2x1H, d, J=8 Hz), 7.57 (1H, d, J=16.2 Hz), 7.68 (2x1H, d, J=8 Hz),
7.74 (1H, d, J=8.8 Hz), 8.14 (1H, dd, J=8.8, 2.2 Hz), 8.46 (1H, d, J=2.2 Hz);

MASS (ES+): m/e 324.

Preparation 105

10 Compound (105) was obtained from Compound (104) according to a manner similar to Preparation 9 (1.42 g). The Compound (105) was used in Example 35.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.45-1.76 (6H, m), 3.53 (1H, m), 3.95 (1H, m), 4.30 (2H, s), 4.90 (1H, m), 6.48 (1H, d, J=15.8 Hz), 7.39
15 (2x1H, d, J=8.5 Hz), 7.47 (1H, d, J=15.8 Hz), 7.55 (2x1H, d, J=8.5 Hz), 7.67 (1H, d, J=8.8 Hz), 8.08 (2x1H, d, J=8.8 Hz), 8.41 (1H, d, J=2.2 Hz), 11.25 (1H, br), 13.02 (1H, br);

MASS (ES+): m/e 423.

Preparation 106

20 Thionyl chloride (1.75 mL) was dropwise added under stirring to methanol at 0°C. After 30 minutes 4-hydroxycinnamic acid (3.29 g) was added and the mixture was refluxed for 1.5 h. The solvent was evaporated in vacuo and the residue was crystalized from diisopropyl ether and hexane to give Compound (106) (2.41 g) as a
25 white crystal.

¹H-NMR (300 MHz, CDCl₃, δ): 3.80 (3H, s), 5.44 (1H, s), 6.31 (1H, d, J=16 Hz), 6.85 (2x1H, d, J=8.5 Hz), 7.43 (2x1H, d, J=8.5 Hz), 7.64 (1H, d, J=16 Hz);

MASS (ES-): m/e 177.

30 Preparation 107

To a stirred solution of Compound (106) (609 mg) in dimethylformamide (15 mL) was added sodium hydride (164 mg, 60% oil dispersion) at 0°C. After 30 minutes, tert-butyl bromoacetate (733 mg) was added dropwise, and the mixture was stirred at ambient
35 temperature for 2 hours. The resulting mixture was poured into 10% citric acid solution and extracted with ethyl acetate. The organic

phase was washed with saturated NaHCO₃ solution and brine, dried over sodium sulfate, and concentrated in vacuo. The residue was purified by chromatography (hexane: ethyl acetate = 4: 1) to give Compound (107) (962 mg) as a solid.

5 ¹H-NMR (300 MHz, CDCl₃, δ): 1.49 (3x3H, s), 3.80 (3H, s), 4.55 (2H, s), 6.32 (1H, d, J=15.8 Hz), 6.90 (2x1H, d, J=8.8 Hz), 7.47 (2x1H, d, J=8.8 Hz), 7.65 (1H, d, J=15.8 Hz);

MASS (ES+): m/e not detected.

Preparation 108

10 A solution of Compound (107) (906 mg) in 0.5 N-hydrogen chloride in acetic acid (10 mL) was heated at 60°C for 2 hours. The solvent was evaporated in vacuo and the residue was triturated with diisopropyl ether to give Compound (108) (645 mg) as a white solid.

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.71 (3H, s), 4.74 (2H, s), 6.51 (1H, d, J=16 Hz), 6.96 (2x1H, d, J=8.5 Hz), 7.62 (1H, d, J=16 Hz), 7.67 (2x1H, d, J=8.5 Hz), 13.07 (1H, br-s);

MASS (ES-): m/e 235.

Preparation 109

20 Compound (109) was obtained from Compound (108) according to a manner similar to Preparation 7 (950 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.45 (3x3H, s), 3.71 (3H, s), 4.79 (2H, s), 6.53 (1H, d, J=16 Hz), 7.07 (2x1H, d, J=8.8 Hz), 7.08-7.20 (2H, m), 7.46 (1H, m), 7.56 (1H, m), 7.63 (1H, d, J=16 Hz), 7.72 (2x1H, d, J=8.8 Hz), 8.73 (1H, br-s), 9.55 (1H, br-s);

25 MASS (ES+): m/e 427.

Preparation 110

Compound (110) was obtained from Compound (109) according to a manner similar to Preparation 8 (800 mg).

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.71 (3H, s), 5.69 (2H, s), 6.56 (1H, d, J=16 Hz), 7.20 (2x1H, d, J=8.8 Hz), 7.49-7.57 (2H, m), 7.65 (1H, d, J=16 Hz), 7.73-7.85 (4H, m);

MASS (ES+): m/e 309.

Preparation 111

35 Compound (111) was obtained from Compound (110) according to a manner similar to Preparation 12 (580 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 5.45 (2H, s), 6.41 (1H, d, $J=16$ Hz), 7.14 (2x1H, d, $J=8.8$ Hz), 7.23-7.32 (2H, m), 7.55 (1H, d, $J=16$ Hz), 7.56-7.66 (2H, m), 7.68 (2x1H, d, $J=8.8$ Hz), 12.28 (1H, br);

MASS (ES+): m/e 295.

5 Preparation 112

Compound (112) was obtained from Compound (111) according to a manner similar to Preparation 9 (503 mg). The Compound (112) was used in Example 36.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.47-1.76 (6H, m), 3.53 (1H, m), 3.95 (1H, m), 4.89 (1H, m), 5.37 (2H, s), 6.37 (1H, d, $J=16$ Hz), 7.10-7.25 (4H, m), 7.39-7.67 (5H, m), 11.16 (1H, s), 12.68 (1H, s);

MASS (ES+): m/e 394.

Preparation 113

To a stirred solution of Compound (86) (165 mg) in methanol (3 mL) were added cyclopentanone (52 mg) and sodium cyanoborohydride (39 mg). To the mixture was added acetic acid so that final pH was set to 5. The mixture was stirred at ambient temperature for 2 hours. The resulting mixture was poured into water and extracted with ethyl acetate. The organic phase was washed with brine, dried over magnesium sulfate, and concentrated in vacuo. The residue was purified by preparative thin layer chromatography (chloroform: methanol = 10: 1) to give Compound (113) (140 mg) as a brown oil.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.34 (3H, t, $J=7.3$ Hz), 1.40-1.77 (6H, m), 1.95-2.08 (2H, m), 3.76 (1H, m), 4.24 (2H, s), 4.26 (2H, q, $J=7.3$ Hz), 6.41 (1H, d, $J=16$ Hz), 6.56 (1H, dd, $J=7, 2.2$ Hz), 6.64 (1H, s), 7.31 (2x1H, d, $J=8$ Hz), 7.36 (1H, d, $J=8.8$ Hz), 7.48 (2x1H, d, $J=8$ Hz), 7.65 (1H, d, $J=16$ Hz);

MASS (ES+): m/e 390.

30 Preparation 114

Compound (114) was obtained from Compound (113) according to a manner similar to Preparation 12 (91 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.36-1.76 (6H, m), 1.89 (2H, m), 4.08 (2H, s), 6.44-6.56 (3H, m), 7.17 (1H, m), 7.34 (2x1H, d, $J=8$ Hz), 7.55 (1H, d, $J=16$ Hz), 7.62 (2x1H, d, $J=8$ Hz), 11.72 (1H, br);

MASS (ES+): m/e 362.

Preparation 115

Compound (115) was obtained from Compound (114) according to a manner similar to Preparation 9 (40 mg). The Compound (115) was used in Example 37.

- 5 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.38–2.08 (14H, m), 3.62 (1H, m), 3.75 (1H, m), 4.00 (1H, m), 4.11 (2H, s), 5.08 (1H, m), 6.20 (1H, br), 6.56 (1H, d, $J=8$ Hz), 6.68 (1H, s), 6.94–7.55 (6H, m);
MASS (ES+): m/e 461.

Preparation 116

- 10 Compound (116) was obtained according to a manner similar to Preparation 106 (1.77 g).

- $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.72 (3H, s), 6.53 (1H, d, $J=16$ Hz), 6.84 (1H, br-dd, $J=8$, 2 Hz), 7.04 (1H, br-d, $J=2$ Hz), 7.14 (1H, br-d, $J=8$ Hz), 7.22 (1H, dd, $J=8$, 8 Hz), 7.57 (1H, d, $J=16$ Hz), 9.63
15 (1H, s);
MASS (ES-): m/e 177.

Preparation 117

Compound (117) was obtained from Compound (116) according to a manner similar to Preparation 107 (1.61 g).

- 20 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.49 (3x3H, s), 3.81 (3H, s), 4.54 (2H, s), 6.41 (1H, d, $J=16$ Hz), 6.92 (1H, dd, $J=8$, 2.5 Hz), 7.04 (1H, dd, $J=2.5$, 2.5 Hz), 7.15 (1H, br-d, $J=8$ Hz), 7.31 (1H, dd, $J=8$, 8 Hz), 7.64 (1H, d, $J=16$ Hz);
MASS (ES+): m/e not detected.

25 Preparation 118

Compound (118) was obtained from Compound (117) according to a manner similar to Preparation 8 (1.25 g).

- $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 3.82 (3H, s), 4.72 (2H, s), 6.43 (1H, d, $J=16$ Hz), 6.96 (1H, dd, $J=8$, 2.5 Hz), 7.07 (1H, br-s), 7.20 (1H, br-d, $J=8$ Hz), 7.34 (1H, dd, $J=8$, 8 Hz), 7.65 (1H, d, $J=16$ Hz);
30 MASS (ES-): m/e 235.

Preparation 119

Compound (119) was obtained from Compound (118) according to a manner similar to Preparation 7 (1.92 g).

- 35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.49 (3x3H, s), 3.82 (3H, s), 4.69 (2H, s), 6.45 (1H, d, $J=16$ Hz), 6.69 (1H, br-s), 7.03 (1H, dd, $J=8$, 2.5

Hz), 7.15-7.26 (4H, m), 7.37 (1H, dd, J=8, 8 Hz), 7.41 (1H, m),
7.63 (1H, m), 7.66 (1H, d, J=16 Hz), 9.00 (1H, br-s);

MASS (ES+): m/e 427.

Preparation 120

5 Compound (120) was obtained from Compound (119) according to
a manner similar to Preparation 8 (1.67 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.74 (3H, s), 5.68 (2H, s), 6.74 (1H,
d, J=16 Hz), 7.21 (1H, m), 7.40-7.46 (2H, m), 7.49-7.59 (3H, m),
7.68 (1H, d, J=16 Hz), 7.77-7.85 (2H, m);

10 MASS (ES+): m/e 309.

Preparation 121

Compound (121) was obtained from Compound (120) according to
a manner similar to Preparation 12 (1.24 g).

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 5.37 (2H, s), 6.58 (1H, d, J=16 Hz),
7.11-7.26 (3H, m), 7.27-7.40 (2H, m), 7.45 (1H, br-s), 7.50 (1H, m),
7.57 (1H, d, J=16 Hz), 7.63 (1H, m), 12.69 (1H, s);

MASS (ES+): m/e 295.

Preparation 122

20 Compound (122) was obtained from Compound (120) according to
a manner similar to Preparation 9 (1662 mg). The Compound (122)
was used in Example 38.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.46-1.78 (6H, m), 3.54 (1H, m), 3.96
(1H, m), 4.91 (1H, m), 5.36 (2H, s), 6.53 (1H, d, J=16 Hz), 7.12
(1H, dd, J=8, 2.5 Hz), 7.15-7.26 (3H, m), 7.32 (1H, br-s), 7.36 (1H,
25 dd, J=8, 8 Hz), 7.46-7.68 (2H, m), 7.47 (1H, d, J=16 Hz), 11.27 (1H,
br-s), 12.69 (1H, br-s);

MASS (ES+): m/e 394.

Preparation 123

30 Compound (123) was obtained from Compound (86) according to
a manner similar to Preparation 113 (415 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.12 (2x3H, t, J=7 Hz), 1.34 (3H, t, J=7
Hz), 3.34 (2x2H, q, J=7 Hz), 4.23 (2H, s), 4.26 (2H, q, J=7 Hz),
6.39 (1H, d, J=16 Hz), 6.79 (1H, dd, J=9, 2 Hz), 6.91 (1H, d, J=2
Hz), 7.30 (2x1H, d, J=8 Hz), 7.44 (1H, d, J=9 Hz), 7.45 (2x1H, d,
35 J=8 Hz), 7.63 (1H, d, J=16 Hz);

MASS (ES+): m/e 378.

Preparation 124

Compound (124) was obtained from Compound (123) according to a manner similar to Preparation 12 (227 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.05 (2x3H, t, J=7 Hz), 3.28 (2x2H, q, J=7 Hz), 4.12 (2H, s), 6.49 (1H, d, J=16 Hz), 6.61-6.72 (2H, m), 7.28 (1H, br-d, J=8 Hz), 7.35 (2x1H, d, J=8 Hz), 7.56 (1H, d, J=16 Hz), 7.63 (2x1H, d, J=8 Hz), 11.82 (1H, br-s);

MASS (ES+): m/e 350.

Preparation 125

Compound (125) was obtained from Compound (124) according to a manner similar to Preparation 9 (132 mg). The Compound (125) was used in Example 39.

¹H-NMR (300 MHz, CDCl₃, δ): 1.12 (2x3H, t, J=7 Hz), 1.50-1.92 (6H, m), 3.32 (2x3H, q, J=7 Hz), 3.62 (1H, m), 4.00 (1H, m), 4.15 (2H, br-s), 5.07 (1H, m), 6.22 (1H, br), 6.71-6.83 (2H, m), 6.96-7.28 (3H, m), 7.38-7.54 (3H, m);

MASS (ES+): m/e 449.

Preparation 126

Compound (126) was obtained according to a manner similar to Preparation 106 (4.19 g).

¹H-NMR (300 MHz, CDCl₃, δ): 3.80 (3H, s), 3.93 (3H, s), 5.85 (1H, br-s), 6.30 (1H, d, J=16 Hz), 6.92 (1H, d, J=8 Hz), 7.03 (1H, d, J=2 Hz), 7.08 (1H, dd, J=8, 2 Hz), 7.63 (1H, d, J=16 Hz);

MASS (ES+): m/e not detected.

Preparation 127

Compound (127) was obtained from Compound (126) according to a manner similar to Preparation 107 (5.16 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.47 (3x3H, s), 3.80 (3H, s), 3.92 (3H, s), 4.62 (2H, s), 6.32 (1H, d, J=16 Hz), 6.76 (1H, d, J=8.7 Hz), 7.04-7.09 (2H, m), 7.63 (1H, d, J=16 Hz);

MASS (ES+): m/e not detected.

Preparation 128

Compound (128) was obtained from Compound (127) according to a manner similar to Preparation 8 (4.28 g).

¹H-NMR (300 MHz, CDCl₃, δ): 3.81 (3H, s), 3.93 (3H, s), 4.74 (2H, s), 6.35 (1H, d, J=16 Hz), 6.89 (1H, d, J=9 Hz), 7.07-7.12 (2H, m),

7.63 (1H, d, J=16 Hz);

MASS (ES-): m/e 265.

Preparation 129

Compound (129) was obtained from Compound (128) according to
5 a manner similar to Preparation 7 (5.43 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.42 (3x3H, s), 3.72 (3H, s), 3.86 (3H,
s), 4.76 (2H, s), 6.61 (1H, d, J=16 Hz), 7.02 (1H, d, J=8.5 Hz),
7.08-7.19 (2H, m), 7.25 (1H, dd, J=8.5, 2 Hz), 7.40-7.48 (2H, m),
7.57-7.66 (2H, m), 8.71 (1H, s), 9.46 (1H, s);

10 MASS (ES+): m/e 457.

Preparation 130

Compound (130) was obtained from Compound (129) according to
a manner similar to Preparation 8 (4.35 g).

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.72 (3H, s), 3.85 (3H, s), 5.67 (2H,
s), 6.65 (1H, d, J=16 Hz), 7.20-7.32 (2H, m), 7.49 (1H, s), 7.54-
7.62 (2H, m), 7.63 (1H, d, J=16 Hz), 7.80-7.88 (2H, m);

MASS (ES+): m/e 339.

Preparation 131

Compound (131) was obtained from Compound (130) according to
20 a manner similar to Preparation 12 (1.63 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.83 (3H, s), 5.33 (2H, s), 6.47 (1H,
d, J=16 Hz), 7.14-7.26 (4H, m), 7.36 (1H, s), 7.50 (1H, m), 7.52
(1H, d, J=16 Hz), 7.63 (1H, m), 12.70 (1H, s);

MASS (ES+): m/e 325.

25 Preparation 132

Compound (132) was obtained from Compound (131) according to
a manner similar to Preparation 9 (1.93 g). The Compound (132) was
used in Example 40.

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.45-1.77 (6H, m), 3.53 (1H, m), 3.82
(3H, s), 3.95 (1H, m), 4.90 (1H, m), 5.32 (2H, s), 6.41 (1H, d,
J=16 Hz), 7.10-7.28 (5H, m), 7.43 (1H, d, J=16 Hz), 7.50 (1H, m),
7.62 (1H, m), 11.13 (1H, s), 12.69 (1H, s);

MASS (ES+): m/e 424.

Preparation 133

35 Compound (133) was obtained from Compound (86) according to
a manner similar to Preparation 113 (437 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.34 (3H, t, $J=7$ Hz), 2.92 (2x3H, s), 4.21 (2H, s), 4.26 (2H, q, $J=7$ Hz), 6.39 (1H, d, $J=16$ Hz), 6.75–6.84 (2H, m), 7.24–7.32 (2H, m), 7.42–7.50 (3H, m), 7.63 (1H, d, $J=16$ Hz);

5 MASS (ES+): m/e 350.

Preparation 134

The ethylcarbonyl group of Compound (133) was deprotected according to a manner similar to Preparation 12. The obtained compound (300 mg) was suspended in dioxane (10 mL). To the suspension were added 1N-NaOH (3 mL) and di-tert-butylidicarbonate (407 mg) in dioxane (4mL) and the mixture was stirred at ambient temperature for 12 hours. Additional di-tert-butylidicarbonate (407 mg) and 1N-NaOH (3 mL) were added and the mixture was stirred at ambient temperature for 6 hours. The solvent was evaporated in vacuo and the residue was partitioned between diisopropyl ether and water. The aqueous phase was acidified with hydrochloric acid to pH 5 and the precipitate was collected and washed with water to give Compound (134) (326 mg) as a pale brown powder. The obtained compound (134) was used in Example 41.

20 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.45–1.95 (15H, m), 2.94–3.03 (6H, m), 3.65 (1H, m), 3.96 (1H, m), 4.54–4.64 (2H, m), 5.00 (1H, m), 6.30–7.80 (9H, m);

MASS (ES+) m/e 521.

Preparation 135

25 Compound (135) was obtained according to a manner similar to Preparation 106 (2.11 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 3.79 (3H, s), 3.93 (3H, s), 6.30 (1H, d, $J=16$ Hz), 6.85 (1H, d, $J=8$ Hz), 7.03 (1H, dd, $J=8, 2$ Hz), 7.14 (1H, d, $J=2$ Hz), 7.60 (1H, d, $J=16$ Hz);

30 MASS (ES+): m/e 209.

Preparation 136

Compound (136) was obtained from Compound (135) according to a manner similar to Preparation 107 (3.12 g).

35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.48 (3x3H, s), 3.79 (3H, s), 3.91 (3H, s), 4.61 (2H, s), 6.26 (1H, d, $J=16$ Hz), 6.89 (1H, d, $J=8.3$ Hz), 6.97 (1H, d, $J=2$ Hz), 7.14 (1H, dd, $J=8.3, 2$ Hz), 7.60 (1H, d, $J=16$

Hz);

MASS (ES+): m/e 323.

Preparation 137

Compound (137) was obtained from Compound (136) according to
5 a manner similar to Preparation 8 (2.03 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.71 (3H, s), 3.81 (3H, s), 4.75 (2H, s), 6.54 (1H, d, J=16 Hz), 7.01 (1H, d, J=8 Hz), 7.27 (1H, dd, J=8, 2 Hz), 7.31 (1H, d, J=2 Hz), 7.57 (1H, d, J=16 Hz), 12.87 (1H, br);
MASS (ES+): m/e 267.

10 Preparation 138

Compound (138) was obtained from Compound (137) according to
a manner similar to Preparation 8 (2.97 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.47 (3x3H, s), 3.80 (3H, s), 3.93 (3H, s), 4.72 (2H, s), 6.33 (1H, d, J=16 Hz), 6.95 (1H, d, J=8 Hz), 6.96
15 (1H, br), 7.12-7.29 (4H, m), 7.47 (1H, br-d, J=7.5 Hz), 7.62 (1H, d, J=16 Hz), 7.64 (1H, br-d, J=7.5 Hz), 9.02 (1H, s);
MASS (ES+) m/e 457.

Preparation 139

Compound (139) was obtained from Compound (138) according to
20 a manner similar to Preparation 7 (2.29 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.71 (3H, s), 3.83 (3H, s), 5.63 (2H, s), 6.61 (1H, d, J=16 Hz), 7.11 (1H, d, J=8.4 Hz), 7.42 (1H, dd, J=8.4, 2 Hz), 7.50-7.58 (2H, m), 7.61 (1H, d, J=16 Hz), 7.65 (1H, d, J=2 Hz), 7.78-7.96 (2H, m);
25 MASS (ES+) m/e 339.

Preparation 140

Compound (140) was obtained from Compound (139) according to
a manner similar to Preparation 12 (1.84 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.81 (3H, s), 5.36 (2H, s), 6.45 (1H, d, J=16 Hz), 7.04 (1H, d, J=8.8 Hz), 7.20-7.28 (2H, m), 7.28 (1H, dd, J=8.8, 2 Hz), 7.52 (1H, d, J=16 Hz), 7.55-7.64 (2H, m), 7.57
30 (1H, d, J=2 Hz), 12.26 (1H, br-s);
MASS (ES+) m/e 325.

Preparation 141

35 Compound (141) was obtained from Compound (140) according to
a manner similar to Preparation 9 (609 mg). The Compound (141) was

used in Example 42.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.42-1.78 (6H, m), 3.53 (1H, m), 3.81 (3H, s), 3.96 (1H, m), 4.90 (1H, m), 5.32 (2H, s), 6.40 (1H, br-d, J=16 Hz), 7.05 (1H, d, J=8.4 Hz), 7.16-7.26 (3H, m), 7.36-7.47 (2H, m), 7.50-7.66 (2H, m), 11.18 (1H, br), 12.69 (1H, br);
5 MASS (ES+) m/e 424.

Preparation 142

Compound (142) was obtained from Compound (6) according to a manner similar to Preparation 7 (400 mg).

10 ¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.76 (2H, s), 5.86 (2H, s), 6.33 (1H, s), 6.38 (1H, d, J=16.1 Hz), 6.77 (1H, s), 7.37 (2H, d, J=8.1 Hz), 7.54 (2H, d, J=8.1 Hz), 7.58 (1H, d, J=16.5 Hz);
MASS (ES+): m/e 397 (M+1).

Preparation 143

15 Compound (143) was obtained from Compound (142) according to a manner similar to Preparation 8 (305 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.44 (2H, s), 6.15 (2H, s), 6.55 (1H, d, J=16.2 Hz), 7.29 (2H, s), 7.42 (2H, d, J=8.1 Hz), 7.58 (1H, d, J=15.7 Hz), 7.72 (2H, d, J=8.4 Hz);
20 MASS (ES+): m/e 323 (M+1).

Preparation 144

Compound (144) was obtained from Compound (143) according to a manner similar to Preparation 9 (196 mg). The Compound (144) was used in Example 43.

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.69 (3H, br), 3.53 (1H, m), 3.95 (1H, m), 4.11 (2H, s), 4.90 (1H, s), 5.95 (2H, s), 6.46 (1H, d, J=15.4 Hz), 6.96 (1H, s), 7.03 (1H, s), 7.34 (2H, d, J=8.1 Hz), 7.46 (1H, d, J=15.8 Hz), 7.52 (2H, d, J=8.1 Hz);
MASS (ES+): m/e 422 (M+1).

Preparation 145

Compound (145) was obtained from Compound (6) according to a manner similar to Preparation 7 (460 mg).

35 ¹H-NMR (300 MHz, CDCl₃, δ): (1:1 mixture) 1.54 (9H, s), 3.71 (3H, s), 3.82 (2H, s), 6.30 (1H, d, J=8.1 Hz), 6.39 (1H, d, J=16.1 Hz), 6.41 (1H, d, J=8.0 Hz), 6.98 (1H, t, J=8.4 Hz), 7.40 (2H, d, J=8.4 Hz), 7.54 (2H, d, J=8.4 Hz), 7.59 (1H, d, J=15.7 Hz), 1.54 (9H, s), 3.77

(2H, s), 3.83 (3H, s), 6.37 (1H, d, J=15.8 Hz), 6.65-6.74 (sH, m), 6.80 (1H, dd, J=7.7, 1.8 Hz), 7.37 (2H, d, J=8.4 Hz), 7.53 (2H, d, J=8.4 Hz), 7.57 (1H, d, J=17.2 Hz);

MASS (ES+): m/e 383 (M+1).

5 Preparation 146

Compound (146) was obtained from Compound (145) according to a manner similar to Preparation 8 (326 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.00 (3H, s), 4.44 (2H, s), 6.54 (1H, d, J=15.7 Hz), 7.06 (1H, d, J=8.7 Hz), 7.24 (1H, d, J=8.1 Hz), 7.40 (1H, t, J=8.1 Hz), 7.44 (2H, d, J=8.4 Hz), 7.58 (1H, d, J=16.0 Hz), 7.71 (2H, d, J=8.2 Hz);

MASS (ES+): m/e 309 (M+1).

Preparation 147

15 Compound (147) was obtained from Compound (146) according to a manner similar to Preparation 9 (308 mg). The Compound (147) was used in Example 44.

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (3H, br), 1.69 (3H, br), 3.52 (1H, m), 3.90 (3H, s), 3.95 (1H, m), 4.15 (2H, s), 4.90 (1H, s), 6.46 (1H, d, J=16.2 Hz), 6.68 (1H, br), 7.04 (2H, m), 7.35 (2H, d, J=8.0 Hz), 7.45 (1H, d, J=16.4 Hz), 7.52 (2H, d, J=8.0 Hz);

MASS (ES+): m/e 408 (M+1).

Preparation 148

Compound (148) was obtained according to a manner similar to Preparation 6 (929 mg).

25 ¹H-NMR (300 MHz, CDCl₃, δ): 1.50 (3H, d, J=6.9 Hz), 1.53 (9H, s), 3.75 (1H, q, J=7.1 Hz), 6.33 (1H, d, J=16.1 Hz), 7.32 (2H, d, J=8.5 Hz), 7.45 (2H, d, J=8.1 Hz), 7.55 (1H, d, J=16.1 Hz);

MASS (ES+): not detected.

Preparation 149

30 Compound (149) was obtained from Compound (148) according to a manner similar to Preparation 7 (1.09 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 1.60 (3H, d, J=7.0 Hz), 3.79 (1H, q, J=7.0 Hz), 6.36 (1H, d, J=16.0 Hz), 6.70-6.78 (2H, m), 6.98-7.04 (1H, m), 7.08-7.15 (1H, m), 7.40 (2H, d, J=8.6 Hz), 7.51 (2H, d, J=8.4 Hz), 7.56 (1H, d, J=16.0 Hz);

MASS (ES+): m/e 367 (M+1).

Preparation 150

Compound (150) was obtained from Compound (149) according to a manner similar to Preparation 8 (567 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.86 (3H, d, J=7.0 Hz), 4.83 (1H, q, J=7.0 Hz), 6.55 (1H, d, J=15.8 Hz), 7.49 (2H, d, J=8.1 Hz), 7.52-7.55 (2H, m), 7.58 (1H, d, J=16.4 Hz), 7.73 (2H, d, J=8.1 Hz), 7.76-7.79 (2H, m);

MASS (ES+): m/e 293 (M+1).

Preparation 151

10 Compound (151) was obtained from Compound (150) according to a manner similar to Preparation 9 (598 mg). The Compound (151) was used in Example 45.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.68 (3H, br), 1.70 (3H, d, J=7.3 Hz), 3.52 (1H, m), 3.93 (1H, m), 4.41 (1H, q, J=7.0 Hz), 15 4.89 (1H, s), 6.47 (1H, d, J=16.5 Hz), 7.08-7.15 (2H, m), 7.34-7.40 (2H, m), 7.37 (2H, d, J=8.3 Hz), 7.44 (1H, d, J=16.5 Hz), 7.52 (2H, d, J=8.0 Hz);

MASS (ES+): m/e 392 (M+1).

Preparation 152

20 Compound (152) was obtained according to a manner similar to Preparation 6 (1.2 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.51 (3H, d, J=6.9 Hz), 1.53 (9H, s), 3.76 (1H, q, J=7.1 Hz), 6.34 (1H, d, J=16.0 Hz), 7.33 (2H, d, J=8.5 Hz), 7.47 (2H, d, J=8.1 Hz), 7.56 (1H, d, J=16.0 Hz);

25 MASS (ES+): not detected.

Preparation 153

Compound (153) was obtained from Compound (152) according to a manner similar to Preparation 7 (1.298 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 1.61 (3H, d, J=7.0 Hz), 30 3.79 (1H, q, J=7.0 Hz), 6.36 (1H, d, J=16.0 Hz), 6.70-6.75 (2H, m), 6.97-7.05 (1H, m), 7.09-7.14 (1H, m), 7.24 (2H, d, J=8.4 Hz), 7.51 (2H, d, J=8.5 Hz), 7.56 (1H, d, J=16.2 Hz);

MASS (ES+): m/e 367 (M+1).

Preparation 154

35 Compound (154) was obtained from Compound (153) according to a manner similar to Preparation 8 (611 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.86 (3H, d, $J=7.4$ Hz), 4.83 (1H, q, $J=7.1$ Hz), 6.55 (1H, d, $J=15.7$ Hz), 7.49 (2H, d, $J=8.0$ Hz), 7.52-7.55 (2H, m), 7.58 (1H, d, $J=16.2$ Hz), 7.74 (2H, d, $J=8.4$ Hz), 7.76-7.79 (2H, m);

5 MASS (ES+): m/e 293 (M+1).

Preparation 155

Compound (155) was obtained from Compound (154) according to a manner similar to Preparation 9 (700 mg). The Compound (155) was used in Example 46.

10 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.52 (3H, br), 1.68 (3H, br), 1.70 (3H, d, $J=6.9$ Hz), 3.50 (1H, m), 3.93 (1H, m), 4.41 (1H, q, $J=7.1$ Hz), 4.89 (1H, s), 6.48 (1H, d, $J=16.1$ Hz), 7.08-7.14 (2H, m), 7.34-7.40 (2H, m), 7.37 (2H, d, $J=8.3$ Hz), 7.47 (2H, d, $J=16.1$ Hz), 7.51 (2H, d, $J=8.9$ Hz);

15 MASS (ES+): m/e 392 (M+1).

Preparation 156

Compound (156) was obtained according to a manner similar to Preparation 7 (536 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.34 (3H, t, $J=7.0$ Hz), 1.46 (9H, s), 1.48 (9H, s), 4.05 (2H, q, $J=7.0$ Hz), 6.50 (1H, d, $J=16.1$ Hz), 6.98 (2H, d, $J=8.4$ Hz), 7.01 (1H, d, $J=7.7$ Hz), 7.40 (2H, d, $J=7.7$ Hz), 7.50 (2H, d, $J=8.8$ Hz), 7.52 (1H, d, $J=8.4$ Hz), 7.55 (1H, d, $J=16.3$ Hz), 7.66 (2H, d, $J=8.1$ Hz), 7.68 (1H, s);

MASS (ES+): m/e 573 (M+1).

25 Preparation 157

Compound (157) was obtained from Compound (156) according to a manner similar to Preparation 8 (345 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.36 (3H, t, $J=7.0$ Hz), 4.08 (2H, q, $J=7.0$ Hz), 4.55 (2H, s), 6.56 (1H, d, $J=16.1$ Hz), 7.05 (2H, d, $J=8.8$ Hz), 7.49 (2H, d, $J=8.4$ Hz), 7.60 (1H, d, $J=16.1$ Hz), 7.65 (2H, d, $J=9.1$ Hz), 7.74 (2H, d, $J=8.4$ Hz), 7.77 (2H, m), 7.88 (1H, s);

MASS (ES+): m/e 399 (M+1).

Preparation 158

35 Compound (158) was obtained from Compound (157) according to a manner similar to Preparation 9 (370 mg). The Compound (158) was

used in Example 47.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.35 (3H, t, J=7.0 Hz), 1.54 (3H, br),
1.68 (3H, br), 3.53 (1H, m), 3.94 (1H, m), 4.06 (2H, q, J=7.0 Hz),
4.21 (2H, s), 4.90 (1H, s), 6.48 (1H, d, J=16.1 Hz), 7.00 (2H, dd,
5 J=8.8, 2.2 Hz), 7.38 (3H, m), 7.44 (1H, s), 7.50 (1H, m), 7.53-7.60
(5H, m), 7.72 (1H, s);

MASS (ES+): m/e 498 (M+1).

Preparation 159

Compound (159) was obtained from Compound (6) according to a
10 manner similar to Preparation 7 (652 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.47 (9H, s), 1.49 (9H, s), 3.77 (2H,
s), 6.50 (1H, d, J=16.1 Hz), 7.40 (2H, d, J=7.9 Hz), 7.49 (1H, d,
J=7.9 Hz), 7.50 (1H, m), 7.60 (1H, m), 7.67 (2H, d, J=8.1 Hz), 8.00
15 (1H, m), 8.51 (1H, m), 8.55 (1H, m), 8.83 (1H, m), 9.76 (1H, d,
J=7.7 Hz);

MASS (ES+): m/e 530 (M+1).

Preparation 160

Compound (160) was obtained from Compound (159) according to
a manner similar to Preparation 8 (458 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.58 (2H, s), 6.56 (1H, d, J=16.1 Hz),
7.51 (2H, d, J=8.1 Hz), 7.59 (1H, d, J=16.1 Hz), 7.74 (2H, d, J=8.4
Hz), 7.90 (2H, s), 8.14 (1H, s), 8.53 (1H, d, J=7.7 Hz), 8.76 (1H,
d, J=5.1 Hz), 9.15 (1H, s);

MASS (ES+): m/e 356 (M+1).

25 Preparation 161

Compound (161) was obtained from Compound (160) according to
a manner similar to Preparation 9 (316 mg). The Compound (161) was
used in Example 48.

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.69 (3H, br), 3.53 (1H,
m), 3.95 (1H, m), 4.23 (2H, s), 4.90 (1H, s), 6.48 (1H, d, J=15.7
Hz), 7.39 (2H, d, J=8.1 Hz), 7.44-7.52 (4H, m), 7.54 (2H, d, J=8.4
Hz), 7.80 (1H, br), 8.08 (1H, m), 8.53 (1H, m), 8.90 (1H, m);

MASS (ES+): m/e 455 (M+1).

Preparation 162

35 Compound (162) was obtained according to a manner similar to
Preparation 7 (748 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (9H, s), 3.80 (2H, s), 6.38 (1H, d, $J=16.1$ Hz), 6.76 (1H, d, $J=8.1$ Hz), 7.11 (sH,), 7.38 (2H, d, $J=7.7$ Hz), 7.47 (2H, d, $J=7.7$ Hz), 7.52-7.60 (5H, m), 7.71 (2H, d, $J=7.0$ Hz);

5 MASS (ES+): m/e 457 (M+1).

Preparation 163

Compound (163) was obtained from Compound (162) according to a manner similar to Preparation 8 (521 mg).

10 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.49 (2H, s), 6.54 (1H, d, $J=16.1$ Hz), 7.47 (2H, d, $J=7.7$ Hz), 7.56-7.62 (3H, m), 7.67-7.77 (3H, m), 7.81 (2H, s), 7.98 (1H, s);

MASS (ES+): m/e 383 (M+1).

Preparation 164

15 Compound (164) was obtained from Compound (163) according to a manner similar to Preparation 9 (185 mg). The Compound (164) was used in Example 49.

20 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.53 (3H, br), 1.69 (3H, br), 3.53 (1H, m), 3.95 (1H, m), 4.27 (2H, s), 4.90 (1H, s), 6.48 (1H, d, $J=16.1$ Hz), 7.39 (2H, d, $J=8.0$ Hz), 7.46 (1H, d, $J=15.7$ Hz), 7.56 (5H, m), 7.62 (2H, s), 7.67 (3H, m), 7.73 (2H, d, $J=7.0$ Hz), 7.86 (1H, s);

MASS (ES+): m/e 482 (M+1).

Preparation 165

To a solution of Compound (60) in EtOH (3 mL) was added sodium borohydride (28 mg) at 5°C and the mixture was allowed to
25 warm to ambient temperature. After stirred for 0.5 hr, sodium borohydride (14 mg) was added to the mixture, and sodium borohydride (14 mg) was then additionally added three times to the mixture before Compound (60) was disappeared. The mixture was
30 poured into water and washed with ether. The aqueous phase was acidified with 1N hydrochloric acid to pH 4 and extracted with ethyl acetate, washed with brine. The organic phase was dried over Na_2SO_4 and the solvent was removed in vacuo to give Compound (165) (120 mg) as colorless form.

35 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.44 (2H, s), 5.86 (1H, s), 6.06 (1H, br), 6.53 (1H, d, $J=16.1$ Hz), 7.19 (1H, t, $J=7.3$ Hz), 7.29 (2H, t, $J=7.3$ Hz), 7.38 (3H, d, $J=7.0$ Hz), 7.42 (2H, d, $J=8.1$ Hz), 7.56 (1H,

d, J=5.5 Hz), 7.60 (1H, d, J=1.8 Hz), 7.67 (1H, s), 7.70 (2H, d, J=8.4 Hz);

MASS (ES+): m/e 385 (M+1).

Preparation 166

5 Compound (166) was obtained from Compound (165) according to a manner similar to Preparation 9 (104 mg). The Compound (166) was used in Example 50.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.68 (3H, br), 3.50 (1H, m), 3.93 (1H, m), 4.16 (2H, s), 4.90 (1H, s), 5.77 (1H, s), 5.82
10 (1H, br), 6.45 (1H, d, J=16.5 Hz), 7.12 (1H, m), 7.18 (1H, d, J=6.6 Hz), 7.27 (2H, d, J=7.5 Hz), 7.32 (2H, t, J=7.3 Hz), 7.34 (1H, s), 7.36 (2H, d, J=7.4 Hz), 7.42 (2H, m), 7.51 (2H, d, J=8.1 Hz);

MASS (ES+): m/e 484 (M+1).

Preparation 167

15 Compound (167) was obtained from Compound (6) according to a manner similar to Preparation 7 (234 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 2.96 (3H, s), 2.98 (3H, s), 3.80 (2H, s), 6.38 (1H, d, J=16.8 Hz), 6.75 (2H, d, J=8.5 Hz), 6.95 (1H, s), 6.99
20 (1H, d, J=7.3 Hz), 7.14 (1H, d, J=7.3 Hz), 7.40 (2H, d, J=7.0 Hz), 7.43 (2H, d, J=7.0 Hz), 7.54 (2H, d, J=7.3 Hz), 7.59 (1H, d, J=15.0 Hz);

MASS (ES+): m/e 472 (M+1).

Preparation 168

25 Compound (168) was obtained from Compound (167) according to a manner similar to Preparation 8 (263 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.04 (6H, s), 4.60 (2H, s), 6.57 (1H, d, J=16.1 Hz), 7.42 (1H, d, J=8.1 Hz), 7.60 (1H, d, J=16.1 Hz),
7.61-7.72 (2H, m), 7.74 (2H, d, J=8.4 Hz), 7.80 (2H, s), 7.90 (1H, s);

30 MASS (ES+): m/e 398 (M+1).

Preparation 169

Compound (169) was obtained from Compound (168) according to a manner similar to Preparation 9 (116 mg). The Compound (169) was used in Example 51.

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br), 1.69 (3H, br), 2.93 (6H, s), 3.53 (1H, m), 3.94 (1H, m), 4.20 (2H, s), 4.90 (1H, s), 6.47

(1H, d, J=15.8 Hz), 6.81 (2H, d, J=9.1 Hz), 7.38 (4H, d, J=8.4 Hz), 7.43 (1H, m), 7.50 (2H, m), 7.54 (2H, d, J=8.4 Hz), 7.69 (1H, s);
MASS (ES+): m/e 497 (M+1).

Preparation 170

5 Compound (170) was obtained from Compound (6) according to a manner similar to Preparation 7 (334 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.81 (2H, s), 4.02 (2H, br), 6.39 (1H, d, J=16.1 Hz), 7.03 (1H, br), 7.29 (1H, br), 7.38 (2H, d, J=8.0 Hz), 7.44 (1H, br), 7.55 (2H, dd, J=8.0 Hz), 7.58 (1H, d, J=16.2 Hz);

MASS (ES+): m/e 421 (M+1).

Preparation 171

Compound (171) was obtained from Compound (170) according to a manner similar to Preparation 8 (247 mg).

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.39 (2H, s), 6.52 (1H, d, J=15.6 Hz), 7.42 (2H, d, J=7.7 Hz), 7.58 (1H, d, J=15.8 Hz), 7.61 (1H, d, J=8.8 Hz), 7.69 (2H, d, J=8.1 Hz), 7.79 (1H, d, J=9.5 Hz), 7.97 (1H, s);
MASS (ES+): m/e 347 (M+1).

Preparation 172

20 Compound (172) was obtained from Compound (171) according to a manner similar to Preparation 9 (308 mg). The Compound (172) was used in Example 52.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.76 (3H, br), 3.52 (1H, m), 3.96 (1H, m), 4.27 (2H, s), 4.90 (1H, s), 6.48 (1H, d, J=16.4 Hz), 7.38 (2H, d, J=8.5 Hz), 7.46 (1H, d, J=8.4 Hz), 7.47 (1H, d, J=16.0 Hz), 7.55 (2H, d, J=7.9 Hz), 7.69 (1H, d, J=8.1 Hz), 7.85 (1H, s);

MASS (ES+): m/e 346 (M+1).

Preparation 173

30 Compound (173) was obtained from Compound (6) according to a manner similar to Preparation 7 (617 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.48 (9H, s), 3.69 (2H, s), 5.25 (2H, br), 6.50 (1H, d, J=15.7 Hz), 6.49-6.55 (1H, m), 6.95-7.00 (1H, m), 7.37 (2H, d, J=8.1 Hz), 7.54 (1H, d, J=16.1 Hz), 7.65 (2H, d, J=8.4 Hz);

MASS (ES+): m/e 389 (M+1).

Preparation 174

Compound (174) was obtained from Compound (173) according to a manner similar to Preparation 8 (466 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.36 (2H, s), 6.52 (1H, d, J=16.1 Hz),
5 7.31-7.40 (2H, m), 7.43 (2H, d, J=8.1 Hz), 7.57 (1H, d, J=15.7 Hz),
7.69 (2H, d, J=8.1 Hz);

MASS (ES+): m/e 315 (M+1).

Preparation 175

10 Compound (175) was obtained from Compound (174) according to a manner similar to Preparation 9 (542 mg). The Compound (175) was used in Example 53.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.69 (3H, br), 3.52 (1H,
m), 3.96 (1H, m), 4.22 (2H, s), 4.90 (1H, s), 6.47 (1H, d, J=16.1
Hz), 7.13-7.22 (2H, m), 7.37 (2H, d, J=8.0 Hz), 7.47 (1H, d, J=15.7
15 Hz), 7.54 (2H, d, J=8.0 Hz);

MASS (ES+): m/e 414 (M+1).

Preparation 176

Compound (176) was obtained from Compound (6) according to a manner similar to Preparation 7 (555 mg).

20 ¹H-NMR (300 MHz, CDCl₃, δ): 0.93 (6H, d, J=6.6 Hz), 1.54 (9H, s),
1.62 (2H, dt, J=7.0, 7.0 Hz), 1.79 (1H, hept, J=6.8 Hz), 3.76 (2H,
s), 3.90 (2H, t, J=6.5 Hz), 6.30 (2H, m), 6.38 (1H, d, J=16.1 Hz),
6.91 (1H, d, J=9.5 Hz), 7.53 (2H, d, J=8.7 Hz), 7.55 (1H, d, J=16.6
Hz);

25 MASS (ES+): m/e 439 (M+1).

Preparation 177

Compound (177) was obtained from Compound (176) according to a manner similar to Preparation 8 (398 mg).

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 0.94 (6H, d, J=6.7 Hz), 1.64 (2H, dt,
J=6.7, 6.7 Hz), 1.79 (1H, hept, J=6.7 Hz), 4.07 (2H, t, J=6.7 Hz),
4.50 (2H, s), 6.55 (1H, d, J=16.3 Hz), 7.10 (1H, dd, J=8.4, 2.2 Hz),
7.21 (1H, d, J=2.2 Hz), 7.46 (2H, d, J=8.4 Hz), 7.59 (1H, d, J=16.1
Hz), 7.62 (1H, d, J=8.7 Hz), 7.73 (2H, d, J=8.6 Hz);

MASS (ES+): m/e 365 (M+1).

35 Preparation 178

Compound (178) was obtained from Compound (177) according to

a manner similar to Preparation 9 (176 mg). The Compound (178) was used in Example 54.

¹H-NMR (300 MHz, CDCl₃, δ): 0.92 (6H, d, J=6.7 Hz), 1.42-1.87 (6H, m), 1.64 (2H, dt, J=6.7 Hz), 1.80 (1H, hept, J=6.7 Hz), 3.52 (1H, m), 3.94 (2H, t, J=6.7 Hz), 3.97 (1H, m), 4.11 (2H, s), 5.13 (1H, s), 6.18 (1H, br.s), 6.84 (1H, dd, J=8.8, 2.5 Hz), 6.90 (4H, br.s), 7.04 (1H, br.s), 7.29 (1H, br.d), 7.45 (1H, d, J=8.8 Hz);

MASS (ES+): m/e 464 (M+1).

Preparation 179

10 Compound (179) was obtained from Compound (6) according to a manner similar to Preparation 7 (414 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.28 (6H, d, J=6.3 Hz), 3.76 (2H, s), 4.44 (1H, hept, J=6.3 Hz), 6.28 (2H, m), 6.37 (1H, d, J=15.8 Hz), 6.91 (1H, d, J=9.0 Hz), 7.37 (2H, d, J=8.1 Hz), 7.52 (2H, d, J=8.1 Hz), 7.57 (1H, d, J=15.7 Hz);

MASS (ES+): m/e 411 (M+1).

Preparation 180

Compound (180) was obtained from Compound (179) according to a manner similar to Preparation 8 (330 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.29 (6H, d, J=6.1 Hz), 4.69 (1H, hept, J=6.1 Hz), 6.55 (1H, d, J=16.2 Hz), 7.08 (1H, dd, J=2.6, 9.2 Hz), 7.19 (1H, d, J=2.6 Hz), 7.59 (1H, d, J=16.1 Hz), 7.62 (1H, d, J=9.2 Hz), 7.73 (2H, d, J=8.2 Hz);

MASS (ES+): m/e 337 (M+1).

25 Preparation 181

Compound (181) was obtained from Compound (180) according to a manner similar to Preparation 9 (281 mg). The obtained Compound (181) was used in Example 55.

30 ¹H-NMR (300 MHz, CDCl₃, δ): 1.31 (6H, d, J=6.0 Hz), 1.53 (3H, br), 1.69 (3H, br), 3.64 (1H, m), 4.01 (1H, m), 4.22 (2H, s), 4.47 (1H, hept, J=6.0 Hz), 5.05 (1H, m), 6.30 (1H, br), 6.85 (1H, d, J=9.0 Hz), 7.00 (1H, s), 7.07-7.20 (2H, br), 7.26-7.35 (1H, br), 7.42 (2H, d, J=8.8 Hz);

MASS (ES+): m/e 436 (M+1).

35 Preparation 182

Compound (182) was obtained from Compound (6) according to a

manner similar to Preparation 7 (546 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.78 (2H, s), 6.38 (1H, d, J=16.1 Hz),
6.40 (2H, m), 6.89 (1H, s), 7.00 (2H, m), 7.09 (1H, t, J=7.3 Hz),
7.31 (2H, dd, J=8.7,7.6 Hz), 7.38 (2H, d, J=8.5 Hz), 7.54 (2H, d,
5 J=7.4 Hz), 7.58 (1H, d, J=16.2 Hz);
MASS (ES+): m/e 445 (M+1).

Preparation 183

Compound (183) was obtained from Compound (182) according to
a manner similar to Preparation 8 (417 mg).

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.52 (2H, s), 6.56 (1H, d, J=16.1 Hz),
7.17 (1H, t, J=7.3 Hz), 7.21 (1H, dd, J=9.1,2.3 Hz), 7.30 (1H, d,
J=2.3 Hz), 7.41 (2H, t, J=8.1 Hz), 7.48 (2H, d, J=8.5 Hz), 7.59 (1H,
d, J=16.3 Hz), 7.73 (2H, d, J=8.8 Hz), 7.76 (1H, d, J=9.1 Hz);
MASS (ES+): m/e 371 (M+1).

15 Preparation 184

Compound (184) was obtained from Compound (183) according to
a manner similar to Preparation 9 (442 mg). The Compound (184) was
used in Example 56.

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br), 1.69 (3H, br), 3.52 (1H,
m), 3.95 (1H, m), 4.19 (2H, s), 4.90 (1H, s), 6.47 (1H, d, J=16.5
Hz), 6.88 (1H, m), 6.94 (2H, m), 7.06 (2H, m), 7.35 (2H, m), 7.37
(3H, m), 7.46 (1H, d, J=16.4 Hz), 7.54 (2H, m);
MASS (ES+): m/e 470 (M+1).

Preparation 185

25 Compound (185) was obtained from Compound (6) according to a
manner similar to Preparation 7 (163 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 3.74 (3H, s), 3.78 (2H, s), 6.31 (2H, m),
6.38 (1H, d, J=16.1 Hz), 6.93 (1H, d, J=10.0 Hz), 7.38 (2H, d,
J=8.0 Hz), 7.54 (2H, d, J=8.4 Hz), 7.58 (1H, d, J=16.1 Hz);
30 MASS (ES+): m/e 383 (M+1).

Preparation 186

Compound (186) was obtained from Compound (185) according to
a manner similar to Preparation 8 (127 mg).

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.84 (3H, s), 4.52 (2H, s), 6.55 (1H,
d, J=16.1 Hz), 7.12 (1H, dd, J=8.8,2.2 Hz), 7.20 (1H, d, J=2.6 Hz),
7.47 (2H, d, J=8.1 Hz), 7.59 (1H, d, J=16.1 Hz), 7.65 (1H, d, J=9.2

Hz), 7.73 (2H, d, J=8.1 Hz);

MASS (ES+): m/e 309 (M+1).

Preparation 187

Compound (187) was obtained from Compound (186) according to
5 a manner similar to Preparation 9 (127 mg). The Compound (187) was
used in Example 57.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br), 1.69 (3H, br), 3.53 (1H,
m), 3.75 (3H, s), 3.95 (1H, m), 4.15 (2H, s), 6.48 (1H, d, J=15.7
Hz), 6.75 (1H, dd, J=8.8, 2.6 Hz), 6.98 (1H, s), 7.35 (3H, d, J=8.4
10 Hz), 7.46 (1H, d, J=15.7 Hz), 7.53 (2H, d, J=8.0 Hz);

MASS (ES+): m/e 408 (M+1).

Preparation 188

To a stirred solution of Compound (247) (described in
Preparation 247 below, 1.75 g) in dichloromethane (20 mL) was added
15 trifluoroacetic acid (20 mL) and the resulting mixture was stirred
at ambient temperature for 3 hrs. The mixture was concentrated in
vacuo and the residue was dissolved in N,N-dimethylformamide (25
mL). To this solution were successively added 1-
hydroxybenzotriazole (731 mg), O-(tetrahydro-2H-pyran-2-
20 yl)hydroxylamine (588 mg) and 1-[3-(dimethylamino)propyl]-3-
ethylcarbodiimide (840 mg) in an ice bath, and the resulting
mixture was stirred at ambient temperature. After 16 hrs,
additional 1-hydroxybenzotriazole (157 mg), O-(tetrahydro-2H-pyran-
2-yl)hydroxylamine (136 mg) and 1-[3-(dimethylamino)propyl]-3-
25 ethylcarbodiimide (180 mg) were added to the mixture and the
mixture was stirred for additional 4 hrs. The mixture was
extracted with ethyl acetate (300 mL) and successively washed with
5% aqueous potassium hydrogen sulfate (100 mL), saturated aqueous
sodium bicarbonate (100 mL) and brine (100 mL). The organic layer
30 was dried over anhydrous sodium sulfate and concentrated in vacuo.
The residue was purified by flash chromatography eluting with ethyl
acetate to afford Compound (188) (1.82 g) as a colorless amorphous
solid. The Compound (188) was used in Example 60.

¹H-NMR (300 MHz, CDCl₃, δ): 1.50-1.91 (6H, m), 3.58-3.71 (1H, m),
35 3.82 (3H, s), 3.88-4.21 (1H, m), 4.87-5.09 (3H, m), 6.17-6.40 (1H,
m), 7.03-7.61 (5H, m), 7.74-7.87 (4H, m);

MASS (ES+): m/z 496 (M+1).

Preparation 189

Compound (189) was obtained from Compound (249) described later according to a manner similar to Preparation 188 (142 mg).

5 The Compound (189) was used in Example 64.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.46-1.92 (6H, m), 3.56-3.69 (1H, m), 3.89-4.18 (2H, m), 4.96-5.12 (2H, m), 6.11-6.37 (1H, m), 7.00-7.36 (7H, m), 7.64 (2H, d, $J=8.4$ Hz);

MASS (ES+): m/z 438 (M+1).

10 Preparation 190

Compound (190) was obtained from Compound (250) described later according to a manner similar to Preparation 188 (110 mg). The Compound (190) was used in Example 68.

15 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.43-1.90 (6H, m), 3.58-3.71 (1H, m), 3.91-4.12 (3H, m), 5.03 (1H, br.s), 5.95-6.21 (1H, m), 7.00-7.42 (9H, m), 7.67-7.76 (2H, m);

MASS (ES+): m/z 404 (M+1).

Preparation 191

20 Compound (191) was obtained from Compound (226) described later according to a manner similar to Preparation 188 (78 mg). The Compound (191) was used in Example 69.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.41-2.05 (6H, m), 3.54-3.69 (1H, m), 3.89-4.05 (3H, m), 4.93 (1H, br.s), 6.04-6.28 (1H, m), 6.95-7.53 (8H, m), 7.73 (2H, d, $J=7.0$ Hz);

25 MASS (ES+): m/z 438 (M+1).

Preparation 192

Compound (192) was obtained from Compound (242) described later according to a manner similar to Preparation 188 (638 mg). The Compound (192) was used in Example 86.

30 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.26 (3H, t, $J=7.0$ Hz), 1.54 (3H, br.), 1.69 (3H, br.), 3.53 (1H, m), 3.95 (1H, m), 4.07 (2H, s), 4.24 (2H, q, $J=7.0$ Hz), 4.90 (1H, s), 6.46 (1H, d, $J=16.1$ Hz), 7.36 (5H, m), 7.52 (3H, m), 7.85 (2H, d, $J=7.3$ Hz);

MASS (ES+): m/e 476 (M+1).

35 Preparation 193

To a stirred solution of Compound (326) (200 mg, described

later in Preparation 326) in N,N-dimethylformamide (2 mL) were added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide (90 mg) and 1-hydroxybenzotriazole (79 mg) at ambient temperature, and the resulting mixture was stirred at the same temperature for 2 hrs.

5 To this mixture was added 28% ammonium hydroxide solution in water (0.08 mL), and the mixture was stirred at ambient temperature for 16 hrs. The mixture was extracted with ethyl acetate (50 mL), and successively washed with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous
10 sodium sulfate and concentrated in vacuo. The residue was purified by preparative thin-layer chromatography eluting with methanol-chloroform (10:90 v/v) to afford Compound (193) (141 mg) as a colorless amorphous solid. The Compound (193) was used in Example 61.

15 ¹H-NMR (300 MHz, CDCl₃-CD₃OD (9:1 v/v), δ): 1.51-1.94 (6H, m), 3.59-3.72 (3H, m), 3.95-4.12 (1H, m), 4.99 (2H, br.s), 6.23-6.47 (1H, m), 7.22-7.30 (2H, m), 7.32-7.47 (4H, m), 7.55-7.69 (4H, m);
MASS (ES⁺): m/z 447 (M+1).

Preparation 194

20 To a stirred solution of Compound (193) (138 mg) in pyridine (2 mL) was added trifluoroacetic anhydride (84 mg) in an ice bath. The resulting mixture was stirred at the same temperature for 2 hrs and then allowed to warm to ambient temperature. After 2 hrs, additional trifluoroacetic anhydride (84 mg) was added to the
25 reaction mixture at ambient temperature, and the mixture was stirred at the same temperature for an hour. The mixture was concentrated in vacuo and the residue was extracted with chloroform (50 mL), and successively washed with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried
30 over anhydrous sodium sulfate and evaporated. The residue was purified by preparative thin-layer chromatography eluting with ethyl acetate to afford Compound (194) (55 mg) as a colorless amorphous solid. The Compound (194) was used in Example 62.

35 ¹H-NMR (300 MHz, CDCl₃, δ): 1.32-2.01 (6H, m), 3.48-4.10 (3H, m), 4.16 (2H, br.s), 4.86-5.06 (1H, m), 6.97-7.52 (8H, m), 7.81-7.92 (2H, m);

MASS (ES+): m/z 429 (M+1).

Preparation 195

To a stirred solution of Compound (6) (500 mg) in N,N-dimethylformamide (8 mL) were successively added 2,2-diethoxyethanamine (305 mg), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (439 mg) and 1-hydroxybenzotriazole (309 mg) in an ice bath, and the resulting solution was stirred at the same temperature for 15 min. The mixture was then allowed to warm to ambient temperature and stirred at the same temperature for 16 hrs. The mixture was extracted with ethyl acetate (50 mL), and successively washed with 5% aqueous potassium sulfate (25 mL), saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous sodium sulfate and evaporated to give crude Compound (195) (861 mg) as a colorless oil, which was used in the next step without further purification.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.13 (3H, t, $J=7.3$ Hz), 1.54 (9H, s), 3.36 (2H, t, $J=5.5$ Hz), 3.40-3.52 (1H, m), 3.58 (2H, s), 3.59-3.70 (1H, m), 4.42 (1H, t, $J=5.5$ Hz), 5.62 (1H, br.s), 6.36 (1H, d, $J=16.1$ Hz), 7.28 (2H, d, $J=8.1$ Hz), 7.49 (2H, d, $J=8.1$ Hz), 7.57 (1H, d, $J=16.1$ Hz);

MASS (ES+): m/z 378 (M+1).

Preparation 196

A mixture of Compound (195) (603 mg), ammonium acetate (616 mg) and acetic acid (1.9 g) in xylene (12 mL) was heated at 170°C for five days, during which time additional same amounts of ammonium acetate and acetic acid were added to the mixture every 12 hrs. The mixture was concentrated in vacuo and extracted with chloroform (100 mL). The organic phase was successively washed with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous sodium sulfate and evaporated. The residue was triturated with acetonitrile to afford Compound (196) (200 mg) as a pale tan amorphous solid.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.47 (9H, s), 3.98 (2H, s), 6.46 (1H, d, $J=16.1$ Hz), 6.78 (1H, br.s), 7.00 (1H, br.s), 7.25 (2H, d, $J=8.1$ Hz), 7.51 (1H, d, $J=16.1$ Hz), 7.61 (2H, d, $J=8.1$ Hz);

MASS (ES+): m/z 285 (M+1).

Preparation 197

Compound (197) was obtained according to a manner similar to Preparation 188 (51 mg). The obtained Compound (197) was used in Example 112.

5 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.18–2.23 (6H, m), 3.48–4.20 (5H, m), 5.02–5.12 (1H, m), 6.90–7.53 (7H, m);

MASS (ES+): m/z 328 (M+1).

Preparation 198

Compound (198) was obtained from Compound (242) described later according to a manner similar to Preparation 188 (638 mg). The obtained Compound (198) was used in Example 86.

(Compound (198) is the same compound as Compound (192).)

15 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.26 (3H, t, $J=7.0$ Hz), 1.54 (3H, br.), 1.69 (3H, br.), 3.53 (1H, m), 3.95 (1H, m), 4.07 (2H, s), 4.24 (2H, q, $J=7.0$ Hz), 4.90 (1H, s), 6.46 (1H, d, $J=16.1$ Hz), 7.36 (5H, m), 7.52 (3H, m), 7.85 (2H, d, $J=7.3$ Hz);

MASS (ES+): m/e 476 (M+1).

Preparation 199

A solution of Compound (248) (427 mg, described later in Preparation 248) in 1N-hydrogen chloride in acetic acid (10 mL) was heated at 60°C for 2 hrs. The solvent was evaporated in vacuo, and the residue was triturated with the mixture of ethyl acetate and acetic acid (10: 1 (v/v)) to give Compound (199) (353 mg) as an off-white solid.

25 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.41 (2H, s), 6.55 (1H, d, $J=16$ Hz), 7.39 (2x1H, dd, $J=8.8$, 8.8 Hz), 7.47 (2x1H, d, $J=8.2$ Hz), 7.58 (1H, d, $J=16$ Hz), 7.71 (2x1H, d, $J=8.2$ Hz), 7.91 (2x1H, dd, $J=8.8$, 5 Hz), 8.05 (1H, s);

MASS (ES+): m/z 323.

Preparation 200

Compound (200) was obtained from Compound (207) described later according to a manner similar to Preparation 199 (361 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.51 (3H, s), 2.53 (3H, s), 4.28 (2H, s);

35 MASS (ES+): m/z 285.

Preparation 201

Compound (201) was obtained from Compound (206) described later according to a manner similar to Preparation 199 (433 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.25 (3H, s), 4.31 (2H, s), 6.55 (1H, d, $J=16$ Hz), 7.45 (2x1H, d, $J=8$ Hz), 7.54-7.63 (3H, m), 7.67-7.75

5 (3H, m), 7.86 (2x1H, d, $J=8$ Hz);

MASS (ES+): m/z 347.

Preparation 202

Compound (202) was obtained from Compound (256) described later according to a manner similar to Preparation 199 (181 mg).

10 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.65-1.79 (6H, m), 1.85-1.91 (6H, m), 2.00-2.07 (3H, m), 4.32 (2H, s), 6.54 (1H, d, $J=16$ Hz), 7.25 (1H, s), 7.40 (2H, d, $J=8$ Hz), 7.57 (1H, d, $J=16$ Hz), 7.70 (2H, d, $J=8$ Hz).

MASS (ES+): m/z 363 (M+1).

15 Preparation 203

To a stirred solution of Compound (199) (316 mg) in N,N -dimethylformamide (5 mL) were added O -(tetrahydro-2H-pyran-2-yl)hydroxylamine (206 mg), 1-hydroxybenzotriazole (238 mg) and 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (338 mg),

20 and the resulting mixture was stirred at ambient temperature for 5 hrs. The reaction mixture was diluted with ethyl acetate and washed successively with water, saturated sodium hydrogen carbonate solution and brine. The organic phase was dried over sodium

25 preparative thin-layer chromatography (chloroform: methanol = 10: 1) to give Compound (203) (341 mg) as a pale yellow powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.45-1.75 (6H, m), 3.53 (1H, m), 3.95 (1H, m), 4.03 (2H, m), 4.90 (1H, m), 6.46 (1H, d, $J=16$ Hz), 7.14 (2x1H, dd, $J=8.8, 8.8$ Hz), 7.32 (2x1H, d, $J=8.5$ Hz), 7.42-7.57 (4H, 30 m), 7.76 (2x1H, dd, $J=8.8, 6.0$ Hz), 11.22 (1H, s), 11.98 (1H, br-s);

MASS (ES+): m/z 422.

Preparation 204

To a suspension of p -iodo-phenylalanine (1.20 g) in n -BuOH 35 (15 mL) was added 1-phenyl-1,2,3-butanetrione-2-oxime (788 mg). The resulting mixture was refluxed for 2 days and cooled, then

concentrated in vacuo. The residue was triturated with ethyl acetate to give Compound (204) (870 mg) as a pale yellow solid.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.48 (3H, s), 3.96 (2H, s), 7.08 (2x1H, d, $J=8.5$ Hz), 7.44-7.60 (3H, m), 7.67 (2x1H, d, $J=8.5$ Hz), 8.18

5 (2x1H, d, $J=8$ Hz), 12.44 (1H, s);

MASS (ES+): m/z 403.

Preparation 205

Compound (205) was obtained from p-iodophenylalanine according to a manner similar to Preparation 204 (402 mg).

10 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.35 (3H, s), 2.38 (3H, s), 3.91 (2H, s), 7.07 (2x1H, d, $J=8$ Hz), 7.67 (2x1H, d, $J=8$ Hz);

MASS (ES+): m/z 341.

Preparation 206

To a stirred solution of Compound (204) (661 mg) in DMF (10 mL) were added tert-butyl acrylate (0.72 mL), palladium(II) acetate (18 mg), tris(2-methylphenyl)phosphine (100 mg) and N,N-diisopropylethylamine (0.86 mL). The mixture was stirred at 90°C for 4 hrs. The resulting mixture was allowed to cool to ambient temperature, poured into water and extracted with ethyl acetate.

20 The organic phase was washed with brine, dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by silica gel column chromatography eluting with a mixture of hexane and ethyl acetate (1:1 v/v) to give Compound (206) (707 mg) as a pale yellow foam.

25 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.47 (3x3H, s), 2.48 (3H, s), 4.02 (2H, s), 6.48 (1H, d, $J=16$ Hz), 7.30 (2x1H, d, $J=8.5$ Hz), 7.44-7.58 (4H, m), 7.64 (2x1H, d, $J=8.5$ Hz), 8.19 (2x1H, d, $J=7.5$ Hz), 12.49 (1H, s);

MASS (ES+): m/z 403.

30 Preparation 207

Compound (207) was obtained from Compound (205) according to a manner similar to Preparation 206 (370 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (3x3H, s), 2.45 (3H, s), 2.54 (3H, s), 4.08 (2H, s), 6.36 (1H, d, $J=16$ Hz), 7.25 (2x1H, d, $J=8$ Hz),

35 7.48 (2x1H, d, $J=8$ Hz), 7.56 (1H, d, $J=16$ Hz);

MASS (ES+): m/z 341.

Preparation 208

Compound (208) was obtained from Compound (201) according to a manner similar to Preparation 188 (461 mg). The Compound (208) was used in Example 66.

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.46-1.76 (6H, m), 3.53 (1H, m), 3.95 (1H, m), 4.02 (2H, s), 4.90 (1H, m), 6.47 (1H, d, $J=16$ Hz), 7.31 (2x1H, d, $J=8.5$ Hz), 7.42-7.68 (6H, m), 8.19 (2x1H, d, $J=7$ Hz), 11.22 (1H, s), 11.48 (1H, s);
MASS (ES+): m/z 446.

10 Preparation 209

Compound (209) was obtained from Compound (200) according to a manner similar to Preparation 188 (268 mg). The Compound (209) was used in Example 65.

- 15 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.46-1.78 (6H, m), 2.36 (3H, s), 2.39 (3H, s), 3.53 (1H, m), 3.96 (1H, m), 3.97 (2H, s), 4.90 (1H, m), 6.47 (1H, d, $J=16$ Hz), 7.29 (2x1H, d, $J=8$ Hz), 7.47 (1H, d, $J=16$ Hz), 7.53 (2x1H, d, $J=8$ Hz), 11.22 (1H, s), 12.31 (1H, br-s);
MASS (ES+): m/z 384.

Preparation 210

- 20 Compound (210) was obtained from Compound (340) described later according to a manner similar to Preparation 188 (107 mg). The Compound (210) was used in Example 96.

- 25 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.48-1.73 (6H, m), 1.92-2.00 (4H, m), 3.18-3.24 (4H, m), 3.48-3.57 (1H, m), 3.88-4.01 (1H, m), 4.10 (2H, s), 4.87-4.93 (1H, m), 6.42-6.52 (3H, m), 7.25-7.37 (3H, m), 7.46 (1H, d, $J=16$ Hz), 7.52 (2H, d, $J=8$ Hz);
MASS (ES+): m/z 447 (M+1).

Preparation 211

- To a mixture of methyl (2E)-3-[4-(hydroxymethyl)phenyl]acrylate (1.0 g) and triethylamine (1.26 g) in dichloromethane (10 mL) was dropwise added methanesulfonyl chloride (715 mg) in dichloromethane (5 mL) in an ice bath. The resulting mixture was stirred at the same temperature for 1 hr and then allowed to warm to ambient temperature. After stirring at ambient temperature for 3 hrs, the mixture was concentrated in vacuo. The residue was extracted with ethyl acetate (100 mL), and
- 35

washed with 10% aqueous citric acid (25 mL), saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic phase was separated, dried over anhydrous sodium sulfate and evaporated in vacuo to give Compound (211) (655 mg) as a colorless amorphous solid, which was used in the next step without further purification. ¹H-NMR (300 MHz, CDCl₃, δ): 3.82 (3H, s), 4.59 (2H, s), 6.45 (1H, d, J=15.8 Hz), 7.41 (2H, d, J=8.4 Hz), 7.52 (2H, d, J=8.4 Hz), 7.69 (1H, d, J=15.8 Hz).

Preparation 212

To a stirred solution of 4-phenyl-1H-imidazole (115 mg) in N,N-dimethylformamide (0.5 mL) was added sodium hydride (60% oil dispersion, 31 mg) in an ice bath. The mixture was stirred at the same temperature for 15 min and then allowed to warm to ambient temperature. After stirring at ambient temperature for 15 min, the mixture was cooled in an ice bath. To this mixture was dropwise added Compound (211) (180 mg) in N,N-dimethylformamide (1.5 mL), and the mixture was stirred at the same temperature for 2 hrs. The mixture was extracted with ethyl acetate (50 mL), and washed with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The residue was purified by preparative thin-layer chromatography eluting with a mixture of ethyl acetate and hexane (3:1 v/v) to afford Compound (212) (135 mg) as a colorless oil. ¹H-NMR (300 MHz, CDCl₃, δ): 3.81 (3H, s), 5.17 (2H, s), 6.44 (1H, d, J=15.8 Hz), 7.17-7.29 (4H, m), 7.33-7.41 (2H, m), 7.53 (2H, d, J=8.4 Hz), 7.63 (1H, d, J=15.8 Hz), 7.68 (1H, s), 7.76 (2H, d, J=8.4 Hz); MASS (ES⁺): m/z 319 (M+1).

Preparation 213

Compound (213) was obtained from Compound (211) according to a manner similar to Preparation 212 (457 mg). ¹H-NMR (300 MHz, CDCl₃, δ): 2.40 (3H, s), 3.81 (3H, s), 5.10 (2H, s), 6.43 (1H, d, J=16.1 Hz), 7.13 (2H, d, J=6.7 Hz), 7.14 (1H, s), 7.22 (1H, t, J=7.0 Hz), 7.36 (2H, t, J=8.1 Hz), 7.51 (2H, d, J=8.1 Hz), 7.67 (1H, d, J=16.1 Hz), 7.74 (2H, d, J=7.0 Hz);

MASS (ES+): m/e 333 (M+1).

Preparation 214

Compound (214) was obtained from Compound (211) according to a manner similar to Preparation 212 (723 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.25 (3H, s), 3.72 (3H, s), 5.28 (2H, s), 6.64 (1H, d, $J=16.1$ Hz), 7.20 (2H, d, $J=8.4$ Hz), 7.42 (2H, d, $J=8.4$ Hz), 7.64 (2H, d, $J=8.4$ Hz), 7.65 (1H, d, $J=16.1$ Hz), 7.74 (2H, d, $J=8.1$ Hz), 7.84 (1H, s);

MASS (ES+): m/z 367 (M+1).

10 Preparation 215

Compound (215) was obtained from tert-butyl (2E)-3-[3-chloromethylphenyl]acrylate according to a manner similar to Preparation 212 (308 mg).

15 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 5.16 (2H, s), 6.36 (1H, d, $J=15.8$ Hz), 7.18 (1H, d, $J=1.1$ Hz), 7.20 (1H, m), 7.23 (1H, t, $J=7.4$ Hz), 7.34-7.41 (4H, m), 7.48 (1H, d, $J=7.7$ Hz), 7.54 (1H, d, $J=16.1$ Hz), 7.61 (1H, d, $J=1.1$ Hz), 7.76 (2H, dd, $J=8.4, 1.5$ Hz);

MASS (ES+): m/z 361 (M+1).

Preparation 216

20 Compound (216) was obtained from tert-butyl (2E)-3-[3-chloromethylphenyl]acrylate according to a manner similar to Preparation 212 (418 mg).

25 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 2.27 (3H, s), 5.12 (2H, s), 6.34 (1H, d, $J=16.1$ Hz), 7.08 (1H, br.d, $J=7.7$ Hz), 7.24 (1H, s), 7.37 (2H, d, $J=8.38$ Hz), 7.38 (1H, t, $J=7.8$ Hz), 7.47 (1H, d, $J=8.1$ Hz), 7.54 (1H, d, $J=15.8$ Hz), 7.60 (2H, d, $J=8.8$ Hz);

MASS (ES+): m/z 409 (M+1).

Preparation 217

30 Compound (217) was obtained from Compound (211) according to a manner similar to Preparation 212 (354 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 3.81 (3H, s), 5.24 (2H, s), 6.43 (1H, d, $J=16.1$ Hz), 6.98 (1H, s), 7.06-7.14 (3H, m), 7.20-7.23 (1H, m), 7.36-7.46 (3H, m), 7.47-7.57 (3H, m), 7.67 (1H, d, $J=16.1$ Hz);

MASS (ES+): m/z 319 (M+1).

35 Preparation 218

Compound (218) was obtained as a mixture of two regioisomers

from Compound (211) according to a manner similar to Preparation 212 (291 mg).

¹H-NMR (300 MHz, CDCl₃, δ): (for a mixture of two regioisomers) 2.09 (1.2H, s), 2.22 (1.8H, s), 3.81 (3H, s), 5.06 (1.2H, s), 5.08 (0.8H, s), 6.43 (1H, d, J=16.1 Hz), 6.60 (0.6H, s), 6.85 (0.4H, s), 7.05 (0.8H, d, J=8.1 Hz), 7.16 (1.2H, d, J=8.1 Hz), 7.44-7.55 (3H, m), 7.66 (1H, d, J=16.1 Hz);

MASS (ES+): (for a mixture of two regioisomers) m/z 257 (M+1).

Preparation 219

10 To a stirred solution of Compound (212) (185 mg) in a mixture of methanol (1 mL) and 1,4-dioxane (1 mL) was added 1N sodium hydroxide (0.87 mL) at ambient temperature. The resulting solution was stirred at the same temperature for 2 hrs. The mixture was cooled in an ice bath, and to this mixture was added
15 concentrated hydrochloric acid to acidify the mixture. The precipitate was filtered and washed with 50% aqueous methanol (1 mL) to afford Compound (219) (130 mg) as a white amorphous solid.

¹H-NMR (300 MHz, DMSO-d₆, δ): 5.47 (2H, s), 6.58 (1H, d, J=16.1 Hz), 7.38-7.55 (5H, m), 7.60 (1H, d, J=16.1 Hz), 7.77 (2H, d, J=8.4 Hz),
20 7.84 (2H, d, J=7.0 Hz), 8.25 (1H, s), 9.35 (1H, s);

MASS (ES+): m/z 305 (M+1).

Preparation 220

Compound (220) was obtained from Compound (213) according to a manner similar to Preparation 219 (357 mg).

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 2.68 (3H, s), 5.42 (2H, s), 6.58 (1H, d, J=16.1 Hz), 7.42 (1H, t, J=7.7 Hz), 7.43 (2H, d, J=8.4 Hz), 7.51 (2H, t, J=7.5 Hz), 7.60 (1H, d, J=16.1 Hz), 7.75 (2H, d, J=8.4 Hz), 7.83 (2H, d, J=7.3 Hz), 8.14 (1H, s);

MASS (EI+) 319 (M+1).

30 Preparation 221

Compound (221) was obtained from Compound (214) according to a manner similar to Preparation 219 (564 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.25 (3H, s), 5.30 (2H, s), 6.52 (1H, d, J=16.1 Hz), 7.21 (2H, d, J=8.1 Hz), 7.43 (2H, d, J=8.4 Hz), 7.57
35 (1H, d, J=15.8 Hz), 7.64 (2H, d, J=8.1 Hz), 7.70 (2H, d, J=8.1 Hz), 7.94 (1H, s);

MASS (ES+): m/z 353 (M+1).

Preparation 222

To a stirred solution of Compound (219) (105 mg) in N,N-dimethylformamide (1.5 mL) were successively added 1-
5 hydroxybenzotriazole (61 mg), O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (49 mg) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (86 mg) in an ice bath, and the resulting mixture was stirred at ambient temperature for three days. The mixture was extracted with ethyl acetate (50 mL), and washed
10 with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by preparative thin-layer chromatography eluting with a mixture of methanol and chloroform (10: 90 v/v) to afford Compound (222) (120 mg) as a
15 colorless viscous oil. The Compound (222) was used in Example 67.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.50-1.97 (6H, m), 3.60-3.72 (1H, m), 3.90-4.05 (1H, m), 5.02 (1H, br.s), 5.17 (2H, s), 7.14-7.30 (6H, m), 7.32-7.42 (2H, m), 7.44-7.56 (2H, m), 7.62-7.82 (3H, m);

MASS (ES+): m/z 404 (M+1).

20 Preparation 223

Compound (223) was obtained from Compound (220) according to a manner similar to Preparation 203 (431 mg). The Compound (223) was used in Example 78.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.54 (3H, br.), 1.69 (3H, br.), 2.29
25 (3H, s), 3.52 (1H, m), 3.94 (1H, m), 4.90 (1H, s), 5.21 (2H, s), 6.49 (1H, d, $J=16.8$ Hz), 7.16 (1H, t, $J=7.5$ Hz), 7.24 (2H, d, $J=8.1$ Hz), 7.32 (2H, t, $J=7.9$ Hz), 7.47 (1H, d, $J=16.8$ Hz), 7.58 (2H, d, $J=8.1$ Hz), 7.63 (1H, s), 7.71 (2H, d, $J=7.0$ Hz);

MASS (ES+): m/e 418 (M+1).

30 Preparation 224

Compound (224) was obtained from Compound (221) according to a manner similar to Preparation 203 (569 mg). The Compound (224) was used in Example 79.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.54 (3H, br.), 1.69 (3H, br.), 2.25
35 (3H, s), 3.53 (1H, m), 3.95 (1H, m), 4.90 (1H, s), 5.27 (2H, s), 6.48 (1H, d, $J=15.8$ Hz), 7.21 (2H, d, $J=8.1$ Hz), 7.42 (2H, d, $J=8.4$

Hz), 7.47 (1H, d, J=15.8 Hz), 7.58 (2H, d, J=8.1 Hz), 7.65 (2H, d, J=8.4 Hz), 7.84 (1H, s);

MASS (ES+): m/z 452 (M+1).

Preparation 225

5 Compound (225) was obtained from (2E)-3-[4-(1H-imidazol-1-ylmethyl)phenyl]acrylic acid according to a manner similar to Preparation 203 (388 mg). The Compound (225) was used in Example 95.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.42-1.78 (6H, m), 3.53 (1H, m), 3.95
10 (1H, m), 4.90 (1H, m), 5.23 (2H, s), 6.49 (1H, d, J=16 Hz), 6.93 (1H, s), 7.21 (1H, s), 7.27 (2x1H, d, J=8 Hz), 7.47 (1H, d, J=16 Hz), 7.56 (2x1H, d, J=8 Hz), 7.80 (1H, s), 11.25 (1H, s);

MASS (ES+): m/e 328.

Preparation 226

15 To a stirred solution of Compound (250) (125 mg, described later in Preparation 250) in acetonitrile (2 mL) was added N-chlorosuccinimide (70 mg) in an ice bath. The mixture was stirred at ambient temperature for 2 days. The mixture was concentrated in vacuo and the residue was purified by preparative thin-layer
20 chromatography eluting with a mixture of methanol and chloroform (4:96 v/v) to afford Compound (226) (100 mg) as a yellow oil.

¹H-NMR (300 MHz, CDCl₃, δ): 1.52 (9H, s), 4.09 (2H, s), 6.33 (1H, d, J=16.1 Hz), 7.21-7.44 (7H, m), 7.51 (1H, d, J=16.1 Hz), 7.59 (2H, d, J=7.3 Hz);

25 MASS (ES+): m/z 395 (M+1).

Preparation 227

Compound (227) was obtained from Compound (245) described later according to a manner similar to Preparation 226 (230 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 4.09 (2H, s), 6.32 (1H, d, J=15.8 Hz), 7.27 (2H, d, J=7.7 Hz), 7.29 (1H, t, J=7.7 Hz), 7.40
30 (2H, t, J=7.4 Hz), 7.45 (2H, d, J=8.1 Hz), 7.53 (1H, d, J=16.1 Hz), 7.57 (2H, d, J=8.1 Hz);

MASS (ES+): m/e 395 (M+1).

Preparation 228

35 Compound (228) was obtained from Compound (251) described later according to a manner similar to Preparation 226 (157 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 3.80 (3H, s), 3.97 (2H, s), 6.25 (1H, d, $J=16.1$ Hz), 6.90 (2H, d, $J=8.8$ Hz), 7.15 (2H, d, $J=8.1$ Hz), 7.32 (2H, d, $J=8.1$ Hz), 7.44 (1H, d, $J=16.1$ Hz), 7.54 (2H, d, $J=8.8$ Hz);
MASS (ES+): m/e 425 (M+1).

5 Preparation 229

To a solution of Compound (6) (4.1 g) in DMF (41 mL) were added HOBt (2.75 g) and 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide (EDCI, 3.15 g) at 5°C. To the mixture was slowly added 2-amino-1-phenylethanone hydrochloride (2.82 g) at 5°C. The mixture was allowed to warm to ambient temperature and stirred for 2 hrs at the same temperature. The resulting mixture was poured into water (100 mL) and extracted with AcOEt. The organic phase was sequentially washed with NH_4Cl , NaHCO_3 and brine, and dried over NaSO_4 . The solvent was removed in vacuo, and the residue was purified by silica gel flash column chromatography eluting with CHCl_3 : MeOH = 95: 5 to give pale yellow solid. The resulting solid was triturated with isopropyl ether to give Compound (229) (5.425 g) as pale yellow powder.

15
 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.54 (9H, s), 3.69 (2H, s), 4.75 (2H, d, $J=4.0$ Hz), 6.38 (1H, d, $J=16.1$ Hz), 7.35 (2H, d, $J=8.1$ Hz), 7.47-7.56 (1+1+2+2H, m), 7.61 (1H, m), 7.95 (2H, d, $J=8.4$ Hz);
MASS (ES+): m/e 380 (M+1).

Preparation 230

Compound (230) was obtained from Compound (26) according to a manner similar to Preparation 229 (1.73 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 3.67 (2H, s), 3.70 (3H, s), 6.19 (1H, d, $J=7.3$ Hz), 6.38 (1H, d, $J=16.1$ Hz), 6.85 (1H, br.d, $J=7.3$ Hz), 7.25-7.59 (7H, m), 7.60-7.68 (1H, m), 8.06-8.13 (2H, m);
MASS (ES-): m/z 436 (M-1).

30 Preparation 231

Compound (231) was obtained from Compound (6) according to a manner similar to Preparation 229 (5.69 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (9H, s), 3.67 (2H, s), 3.70 (3H, s), 6.12 (1H, d, $J=7.0$ Hz), 6.36 (1H, d, $J=16.1$ Hz), 6.82 (1H, br.d, $J=7.0$ Hz), 7.30 (2H, d, $J=8.1$ Hz), 7.44-7.53 (4H, m), 7.57 (1H, d, $J=16.1$ Hz), 8.03 (2H, d, $J=8.8$ Hz);

MASS (ES+): m/z 472 (M+1).

Preparation 232

Compound (232) was obtained from Compound (6) according to a manner similar to Preparation 229 (3.22 g).

5 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48 (3x3H, s), 3.57 (2H, s), 4.61 (2H, d, J=5.5 Hz), 6.49 (1H, d, J=16 Hz), 7.30-7.42 (4H, m), 7.53 (1H, d, J=16 Hz), 7.63 (2x1H, d, J=8.5 Hz), 8.02-8.11 (2H, m), 8.48 (1H, t, J=5.5 Hz);

MASS (ES-): m/z 442 (M+HCO₂H-1).

10 Preparation 233

Compound (233) was obtained from Compound (6) according to a manner similar to Preparation 229 (1.74 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.68 (2H, s), 4.71 (2H, d, J=5.5 Hz), 6.37 (2H, d, J=16.1 Hz), 6.50 (1H, br.s), 7.34 (2H, d, J=8.1 Hz), 7.47 (2H, d, J=8.4 Hz), 7.52 (2H, d, J=8.1 Hz), 7.58 (1H, d, J=16.1 Hz), 7.89 (2H, d, J=8.4 Hz);

MASS (ES+): m/z 414 (M+1).

Preparation 234

20 Compound (234) was obtained from Compound (26) according to a manner similar to Preparation 229 (1.95 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 3.68 (2H, s), 4.75 (2H, d, J=4.4 Hz), 6.40 (1H, d, J=16.1 Hz), 6.56 (1H, br.s), 7.25-7.70 (8H, m), 7.91-8.02 (2H, m);

MASS (ES+): m/z 380 (M+1).

25 Preparation 235

Compound (235) was obtained from Compound (6) according to a manner similar to Preparation 229 (2.74 g).

30 ¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.68 (2H, s), 3.88 (3H, s), 4.68 (2H, d, J=4.4 Hz), 6.37 (1H, d, J=15.8 Hz), 6.95 (2H, d, J=9.1 Hz), 7.34 (2H, d, J=8.1 Hz), 7.52 (2H, d, J=8.1 Hz), 7.58 (1H, d, J=16.1 Hz), 7.92 (2H, d, J=8.8 Hz);

MASS (ES+): m/e 410 (M+1).

Preparation 236

35 Compound (236) was obtained from Compound (6) according to a manner similar to Preparation 229 (2.86 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.69 (2H, s), 3.85 (3H, s),

4.73 (2H, d, J=4.4 Hz), 6.73 (2H, d, J=15.8 Hz), 7.16 (1H, dd, J=8.1, 2.6 Hz), 7.34 (2H, d, J=8.1 Hz), 7.39 (1H, t, J=8.1 Hz), 7.45 (1H, m), 7.52 (2H, d, J=8.4 Hz), 7.58 (1H, d, J=16.1 Hz);
MASS (ES+): m/e 410 (M+1).

5 Preparation 237

Compound (237) was obtained from Compound (6) according to a manner similar to Preparation 229 (1.82 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.66 (2H, s), 3.93 (3H, s), 4.69 (2H, d, J=4.4 Hz), 6.37 (1H, d, J=16.1 Hz), 6.63 (1H, br),
10 7.00 (1H, t, J=8.4 Hz), 7.03 (1H, t, J=7.7 Hz), 7.35 (2H, d, J=8.4 Hz), 7.51 (2H, d, J=8.1 Hz), 7.58 (1H, d, J=16.1 Hz), 7.90 (1H, dd, J=7.7, 1.8 Hz);
MASS (ES+): m/e 410 (M+1).

Preparation 238

15 Compound (238) was obtained from 4-iodophenylacetic acid according to a manner similar to Preparation 229 (4.20 g).

¹H-NMR (300 MHz, CDCl₃, δ): 3.61 (2H, s), 4.74 (2H, d, J=4.4 Hz), 7.09 (2H, d, J=8.4 Hz), 7.49 (2H, t, J=7.4 Hz), 7.63 (1H, t, J=7.4 Hz), 7.70 (2H, d, J=8.4 Hz), 7.95 (2H, d, J=7.4 Hz);
20 MASS (ES+): m/e 380 (M+1).

Preparation 239

Compound (239) was obtained from Compound (6) according to a manner similar to Preparation 229 (600 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (9H, s), 1.58-1.84 (12H, m),
25 1.99-2.10 (3H, m), 3.61 (2H, s), 4.21 (2H, d, J=4.4 Hz), 6.35 (1H, d, J=16.1 Hz), 7.30 (2H, d, J=8.1 Hz), 7.50 (2H, d, J=8.1 Hz), 7.56 (1H, d, J=16.1 Hz);
MASS (ES+): m/z 438 (M+1).

Preparation 240

30 Compound (240) was obtained from Compound (6) according to a manner similar to Preparation 229 (231 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.68 (2H, s), 3.69 (3H, s), 6.18 (1H, d, J=7.3 Hz), 6.36 (1H, d, J=15.8 Hz), 7.30 (2H, d, J=8.1 Hz), 7.51 (4H, m), 7.62 (2H, m), 8.08 (2H, d, J=7.0 Hz);
35 MASS (ES+): m/e 438 (M+1).

Preparation 241

Compound (241) was obtained according to a manner similar to Preparation 229 (1.33 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.17 (9H, s), 1.54 (9H, s), 3.62 (2H, s), 4.25 (2H, d, $J=4.4$ Hz), 6.36 (1H, d, $J=15.8$ Hz), 7.31 (2H, d, $J=8.4$ Hz), 7.51 (2H, d, $J=8.1$ Hz), 7.57 (1H, d, $J=16.1$ Hz);
MASS (ES+): m/e 360 (M+1).

Preparation 242

Compound (242) was obtained from Compound (6) according to a manner similar to Preparation 229 (1.58 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.12 (3H, t, $J=7.0$ Hz), 1.53 (9H, s), 3.67 (2H, s), 4.13 (2H, q, $J=7.4$ Hz), 6.15 (1H, d, $J=7.3$ Hz), 6.36 (1H, d, $J=15.8$ Hz), 6.86 (1H, br.d, $J=6.6$ Hz), 7.31 (2H, d, $J=8.1$ Hz), 7.48 (2H, t, $J=7.4$ Hz), 7.50 (2H, d, $J=8.1$ Hz), 7.57 (1H, d, $J=16.1$ Hz), 7.63 (1H, t, $J=7.4$ Hz), 8.08 (2H, d, $J=8.1$ Hz);
MASS (ES+): m/z 452 (M+1).

Preparation 243

Compound (243) was obtained from Compound (6) according to a manner similar to Preparation 229 (200 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.30 (4H, m), 1.53 (9H, s), 1.64 (2H, m), 1.80 (4H, m), 2.36 (1H, m), 3.62 (2H, s), 4.17 (2H, d, $J=4.4$ Hz), 6.35 (1H, d, $J=16.1$ Hz), 7.30 (2H, d, $J=8.4$ Hz), 7.49 (2H, d, $J=8.1$ Hz), 7.56 (1H, d, $J=15.8$ Hz);
MASS (ES+): m/z 386 (M+1).

Preparation 244

Compound (244) was obtained from Compound (6) according to a manner similar to Preparation 229 (866 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.34 (3H, t, $J=7$ Hz), 3.60 (2H, s), 3.71 (2H, s), 4.16 (2H, d, $J=5$ Hz), 4.27 (2H, q, $J=7$ Hz), 6.42 (1H, d, $J=16$ Hz), 7.16-7.37 (7H, m), 7.50 (2H, d, $J=8$ Hz), 7.66 (1H, d, $J=16$ Hz);
MASS (ES+): m/z 366 (M+1).

Preparation 245

To a suspension of Compound (229) (5.42 g) in xylene (50 mL) were added AcONH_4 and AcOH , and the mixture was refluxed for 3 hrs with azeotropic removal of water. The resulting solution was cooled to ambient temperature and poured into saturated aqueous NaHCO_3 .

solution (100 mL) and extracted with AcOEt. The organic layer was washed with saturated aqueous NaHCO₃, water and brine, and dried over Na₂SO₄. The solvent was removed in vacuo, and the residual solid was triturated with isopropyl ether to give Compound (245)

5 (4.56 g) as pale yellow powder.

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 4.19 (2H, s), 6.35 (1H, d, J=15.7 Hz), 7.21 (1H, s), 7.25 (1H, t, J=7.0 Hz), 7.29 (2H, d, J=8.4 Hz), 7.38 (2H, t, J=7.0 Hz), 7.48 (2H, d, J=7.7 Hz), 7.56 (1H, d, J=15.4 Hz), 7.69 (2H, br.d);

10 MASS (ES+): m/e 361 (M+1).

Preparation 246

Compound (246) was obtained from Compound (230) according to a manner similar to Preparation 245 (1.10 g).

15 ¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 3.82 (3H, s), 4.17 (2H, s), 6.36 (1H, d, J=16.1 Hz), 7.22-7.48 (9H, m), 7.54 (1H, d, J=16.1 Hz), 7.74-7.87 (1H, m);

MASS (ES+): m/z 419 (M+1).

Preparation 247

20 Compound (247) was obtained from Compound (231) according to a manner similar to Preparation 245 (3.8 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.82 (3H, s), 4.17 (2H, s), 6.36 (1H, d, J=16.1 Hz), 7.29 (2H, d, J=8.1 Hz), 7.38 (2H, d, J=8.1 Hz), 7.49 (2H, d, J=8.1 Hz), 7.55 (1H, d, J=16.1 Hz), 7.78-7.90 (2H, m);

25 MASS (ES+): m/z 453 (M+1).

Preparation 248

Compound (248) was obtained from Compound (232) according to a manner similar to Preparation 245 (2.08 g).

30 ¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (3x3H, s), 4.16 (2H, s), 6.36 (1H, d, J=16.2 Hz), 7.06 (2x1H, dd, J=8.8, 8.7 Hz), 7.14 (1H, s), 7.27 (2x1H, d, J=8 Hz), 7.46 (2x1H, d, J=8 Hz), 7.54 (1H, d, J=16.2 Hz), 7.73 (2H, m), 8.84 (1H, br);

MASS (ES+): m/z 379.

Preparation 249

35 Compound (249) was obtained from Compound (233) according to a manner similar to Preparation 245 (1.12 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (9H, s), 4.17 (2H, s), 6.33 (1H, d, $J=16.1$ Hz), 7.17 (1H, s), 7.27 (2H, d, $J=8.1$ Hz), 7.33 (2H, d, $J=8.4$ Hz), 7.46 (2H, d, $J=8.1$ Hz), 7.53 (2H, d, $J=16.1$ Hz), 7.64 (2H, d, $J=8.4$ Hz);

5 MASS (ES+): m/z 395 (M+1).

Preparation 250

Compound (250) was obtained from Compound (234) according to a manner similar to Preparation 245 (1.28 g).

10 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.52 (9H, s), 4.16 (2H, s), 6.33 (1H, d, $J=16.1$ Hz), 7.17-7.44 (8H, m), 7.52 (1H, d, $J=16.1$ Hz), 7.64-7.73 (2H, m);

MASS (ES+): m/z 361 (M+1).

Preparation 251

15 Compound (251) was obtained from Compound (235) according to a manner similar to Preparation 245 (2.00 g).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (9H, s), 3.82 (3H, s), 4.16 (2H, s), 6.34 (1H, d, $J=15.8$ Hz), 6.91 (2H, d, $J=8.8$ Hz), 7.10 (1H, s), 7.27 (2H, d, $J=8.4$ Hz), 7.47 (2H, d, $J=8.4$ Hz), 7.55 (1H, d, $J=16.1$ Hz), 7.60 (2H, br.d);

20 MASS (ES+): m/z 391 (M+1).

Preparation 252

Compound (252) was obtained from Compound (236) according to a manner similar to Preparation 245 (2.72 g).

25 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 3.78 (3H, s), 4.05 (2H, s), 6.28 (1H, d, $J=16.1$ Hz), 6.78 (1H, m), 7.17 (2H, d, $J=8.1$ Hz), 7.18 (1H, s), 7.247 (2H, d, $J=5.17$ Hz), 7.37 (2H, d, $J=8.4$ Hz), 7.49 (1H, d, $J=15.7$ Hz);

MASS (ES+): m/e 391 (M+1).

Preparation 253

30 Compound (253) was obtained from Compound (237) according to a manner similar to Preparation 245 (1.56 g).

35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (9H, s), 3.85 (3H, s), 4.19 (2H, s, $J=16.1$ Hz), 6.34 (1H, d, $J=8.4$ Hz), 6.76 (1H, d, $J=7.5$ Hz), 7.00 (1H, t, $J=7.5$ Hz), 7.22 (1H, t, $J=8.1$ Hz), 7.30 (2H, d), 7.41 (1H, s, $J=8.4$ Hz), 7.48 (2H, d, $J=16.1$ Hz), 7.72 (1H, d, $J=8.4$ Hz), 7.76 (1H, br.d);

MASS (ES+): m/e 391 (M+1).

Preparation 254

Compound (254) was obtained from Compound (238) according to a manner similar to Preparation 245 (4.00 g).

5 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 4.08 (2H, s), 7.03 (2H, d, $J=8.4$ Hz), 7.24 (1H, t, $J=7.4$ Hz), 7.37 (2H, t, $J=7.7$ Hz), 7.65 (2+2H, d, $J=8.4$ Hz);

MASS (ES+): m/e 361 (M+1).

Preparation 255

10 Compound (255) was obtained from N-(2-oxo-2-phenylethyl)acetamide according to a manner similar to Preparation 245 (803 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.48 (3H, s), 7.20 (1H, s), 7.23 (1H, t, $J=8.1$ Hz), 7.37 (2H, t, $J=7.7$ Hz), 7.69 (2H, d, $J=8.1$ Hz).

15 Preparation 256

Compound (256) was obtained from Compound (239) according to a manner similar to Preparation 245 (460 mg).

20 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 1.69-1.81 (6H, m), 1.83-1.91 (6H, m), 1.99-2.08 (3H, m), 4.09 (2H, s), 6.34 (1H, d, $J=16.1$ Hz), 6.57 (1H, s), 7.23 (2H, d, $J=8.2$ Hz), 7.45 (2H, d, $J=8.2$ Hz), 7.55 (1H, d, $J=16.1$ Hz);

MASS (ES+): m/z 419 (M+1).

Preparation 257

25 Compound (257) was obtained from Compound (240) according to a manner similar to Preparation 245 (58 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.54 (9H, s), 3.82 (3H, s), 4.17 (2H, s), 6.35 (1H, d, $J=16.1$ Hz), 7.28 (2H, d, $J=7.7$ Hz), 7.39 (3H, m), 7.48 (2H, d, $J=7.7$ Hz), 7.55 (1H, d, $J=16.1$ Hz), 7.87 (2H, br.);

MASS (ES+): m/e 419 (M+1).

30 Preparation 258

Compound (258) was obtained from Compound (241) according to a manner similar to Preparation 245 (1.06 g).

35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.28 (9H, s), 1.53 (9H, s), 4.07 (2H, s), 6.31 (1H, d, $J=16.1$ Hz), 6.60 (1H, s), 7.21 (2H, d, $J=8.1$ Hz), 7.43 (2H, d, $J=8.4$ Hz), 7.52 (1H, d, $J=15.8$ Hz);

MASS (ES+): m/e 341 (M+1).

Preparation 259

Compound (259) was obtained from Compound (243) according to a manner similar to Preparation 245 (80 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.32 (4H, m), 1.53 (9H, s), 1.73 (4H, m),
5 1.98 (2H, m), 2.50 (1H, m), 4.04 (2H, s), 6.31 (1H, d, J=16.1 Hz),
6.58 (1H, s), 7.20 (2H, d, J=8.1 Hz), 7.42 (2H, d, J=8.1 Hz), 7.52
(1H, d, J=16.1 Hz);

MASS (ES+): m/z 367 (M+1).

Preparation 260

10 Compound (260) was obtained from Compound (244) according to a manner similar to Preparation 245 (232 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.39 (3H, t, J=7 Hz), 3.92 (2H, s), 4.09
(2H, s), 4.26 (2H, q, J=7 Hz), 6.41 (1H, d, J=16 Hz), 7.21-7.33 (7H,
m), 7.48 (2H, d, J=8 Hz), 7.65 (1H, d, J=16 Hz);

15 MASS (ES+): m/z 347 (M+1).

Preparation 261

To a solution of Compound (297) (2.56 g, described later in Preparation 297 in dichloromethane) (17 mL) was added
trifluoroacetic acid (2.74 mL), and the mixture was stirred for 1
20 hr. The solvent was removed in vacuo, and the residue was
trituated with isopropyl ether to give Compound (261) (2.83 g) as
colorless powder.

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.37 (2H, s), 6.54 (1H, d, J=15.8 Hz),
7.39 (2H, d, J=8.1 Hz), 7.43 (1H, t, J=7.7 Hz), 7.53 (2H, t, J=8.1
25 Hz), 7.58 (1H, d, J=15.8 Hz), 7.72 (2H, d, J=8.4 Hz), 7.77 (2H, d,
J=8.1 Hz), 8.05 (1H, s);

MASS (ES+): m/e 305 (M+1).

Preparation 262

30 Compound (262) was obtained from Compound (246) according to a manner similar to Preparation 261 (1.55 g).

¹H-NMR (300 MHz, CDCl₃-CD₃OD (10:1), δ): 3.87 (3H, s), 4.35 (2H, s),
6.46 (1H, d, J=16.2 Hz), 7.10-7.76 (10H, m);

MASS (ES+): m/z 369 (M+1, free).

Preparation 263

35 Compound (263) was obtained from Compound (251) according to a manner similar to Preparation 261 (2.32 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.81 (3H, s), 4.37 (2H, s), 6.55 (1H, d, J=16.1 Hz), 7.10 (2H, d, J=8.8 Hz), 7.40 (2H, d, J=8.4 Hz), 7.59 (1H, d, J=16.1 Hz), 7.71 (2H, d, J=8.8 Hz), 7.72 (2H, d, J=8.4 Hz), 7.95 (1H, s);

5 MASS (ES+): m/z 335 (M+1).

Preparation 264

Compound (264) was obtained from Compound (252) according to a manner similar to Preparation 261 (1.27 g).

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.83 (3H, s), 4.39 (2H, s), 6.55 (1H, d, J=15.8 Hz), 7.02 (1H, d, J=8.1 Hz), 7.33-7.47 (5H, m), 7.60 (1H, d, J=15.8 Hz), 7.73 (2H, d, J=8.1 Hz), 8.11 (1H, s);

MASS: Not Detected.

Preparation 265

15 Compound (265) was obtained from Compound (253) according to a manner similar to Preparation 261 (1.24 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.93 (3H, s), 4.40 (2H, s), 6.54 (1H, d, J=16.1 Hz), 7.12 (1H, t, J=7.3 Hz), 7.22 (1H, d, J=7.7 Hz), 7.39 (2H, d, J=8.1 Hz), 7.46 (1H, t, J=7.9 Hz), 7.58 (1H, d, J=16.1 Hz), 7.72 (2H, d, J=8.1 Hz), 7.89 (1H, s);

20 MASS (ES+): m/e 335 (M+1).

Preparation 266

Compound (266) was obtained from Compound (293) described later according to a manner similar to Preparation 261 (208 mg).

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.08 (2H, s), 6.05 (1H, d, J=16.1 Hz), 7.35 (2H, d, J=8.4 Hz), 7.37 (1H, t, J=7.4 Hz), 7.49 (2H, d, J=7.4 Hz), 7.57 (1H, d, J=16.1 Hz), 7.66 (2H, d, J=8.1 Hz), 7.71 (1H, d, J=7.0 Hz);

MASS (ES+): m/e 383 (M+1).

Preparation 267

30 Compound (267) was obtained from Compound (227) according to a manner similar to Preparation 261 (234 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.03 (2H, s), 6.49 (1H, d, J=15.8 Hz), 7.32 (1H, t, J=7.3 Hz), 7.34 (2H, d, J=8.4 Hz), 7.47 (2H, t, J=7.3 Hz), 7.57 (1H, d, J=16.1 Hz), 7.65 (2H, d, J=8.4 Hz), 7.70 (2H, d, J=7.3 Hz);

35 MASS (ES+): m/e 339 (M+1).

Preparation 268

Compound (268) was obtained from Compound (294) described later according to a manner similar to Preparation 261 (1.32 g).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.72 (6H, s), 4.21 (2H, s), 4.39 (2H, s), 6.52 (1H, d, J=15.8 Hz), 7.38 (2H, d, J=8.1 Hz), 7.47 (1H, m), 7.54 (4H, m), 7.58 (1H, d, J=15.4 Hz), 7.68 (2H, d, J=8.4 Hz);
5 MASS (ES+): m/e 362 (M+1).

Preparation 269

Compound (269) was obtained from Compound (295) described later according to a manner similar to Preparation 261 (412 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48 (2H, br.), 1.69 (4H, br.), 2.91 (2H, br.), 3.27 (2H, br.), 4.24 (2H, s), 4.38 (2H, s), 6.53 (1H, d, J=16.1 Hz), 7.38 (2H, d, J=8.1 Hz), 7.47-7.61 (6H, m), 7.69 (2H, d, J=8.1 Hz);
15 MASS (ES+): m/e 402 (M+1).

Preparation 270

Compound (270) was obtained from Compound (296) described later according to a manner similar to Preparation 261 (407 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.07 (4H, br.), 3.75 (4H, br.), 4.22 (2H, s), 4.29 (2H, s), 6.52 (1H, d, J=15.7 Hz), 7.39 (2H, d, J=8.1 Hz), 7.46 (1H, t, J=7.0 Hz), 7.53 (2H, t, J=7.0 Hz), 7.58 (1H, d, J=16.1 Hz), 7.59 (2H, d, J=7.3 Hz), 7.68 (2H, d, J=8.1 Hz);
20 MASS (ES+): m/e 404 (M+1).

Preparation 271

Compound (271) was obtained from Compound (228) according to a manner similar to Preparation 261 (171 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.79 (3H, s), 4.03 (2H, s), 6.50 (1H, d, J=16.1 Hz), 7.05 (2H, d, J=8.8 Hz), 7.34 (2H, d, J=8.1 Hz), 7.57 (1H, d, J=15.8 Hz), 7.62 (2H, d, J=8.8 Hz), 7.65 (2H, d, J=8.1 Hz);
30 MASS (ES+): m/e 369 (M+1).

Preparation 272

Compound (272) was obtained from Compound (215) according to a manner similar to Preparation 261 (339 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 5.46 (2H, s), 6.60 (1H, d, J=16.1 Hz), 7.43 (1H, t, J=7.0 Hz), 7.46-7.54 (4H, m), 7.61 (1H, d, J=16.1 Hz), 7.72 (1H, m), 7.78 (2H, d, J=7.0 Hz), 7.84 (1H, s), 8.25 (1H, s),
35

9.29 (1H, s);

MASS (ES+): m/e 305 (M+1).

Preparation 273

5 Compound (273) was obtained from Compound (218) according to a manner similar to Preparation 261 (491 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.30 (3H, s), 5.48 (2H, s), 6.57 (1H, d, $J=16.1$ Hz), 7.34-7.52 (3H, m), 7.58 (1H, d, $J=15.8$ Hz), 7.60 (4H, s), 7.71 (1H, s), 9.26 (1H, s);

MASS (ES+): m/e 353 (M+1).

10 Preparation 274

Compound (274) was obtained from Compound (257) according to a manner similar to Preparation 261 (65 mg).

15 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.78 (3H, s), 4.20 (2H, s), 6.51 (1H, d, $J=15.8$ Hz), 7.38 (2H, d, $J=7.7$ Hz), 7.47 (3H, m), 7.57 (1H, d, $J=15.8$ Hz), 7.67 (2H, d, $J=8.4$ Hz), 7.76 (2H, m);

MASS (ES+): m/e 363 (M+1).

Preparation 275

Compound (275) was obtained from Compound (258) according to a manner similar to Preparation 261 (454 mg).

20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.21 (9H, s), 4.32 (2H, s), 6.55 (1H, d, $J=16.1$ Hz), 7.34 (2H, d, $J=8.4$ Hz), 7.58 (1H, d, $J=16.1$ Hz), 7.71 (2H, d, $J=8.1$ Hz);

MASS (ES+): m/e 285 (M+1).

Preparation 276

25 Compound (276) was obtained from Compound (259) according to a manner similar to Preparation 261 (98.5 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.32 (4H, m), 1.72 (4H, m), 1.94 (2H, m), 2.47 (1H, m), 4.28 (2H, s), 6.54 (1H, d, $J=15.8$ Hz), 7.31 (1H, s), 7.33 (2H, d, $J=8.1$ Hz), 7.58 (1H, d, $J=15.4$ Hz), 7.71 (2H, d, $J=8.1$ Hz);

30 MASS (ES+): m/z 311 (M+1).

Preparation 277

To a solution of Compound (261) (2.46 g) in DMF (30 mL) were added O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (1.66 g), HOBt (2.74 g) and EDCI (3.15 g) at ambient temperature, and the mixture
35 was stirred for 3 hrs. The resulting mixture was diluted with

saturated aqueous NaHCO₃ solution and the precipitate was collected by filtration. The obtained powder was washed with water and dried to give Compound (277) (2.46 g) as pale yellow powder. The obtained Compound (277) was used in Example 58.

- 5 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br.), 1.70 (3H, br.), 3.51 (1H, m), 3.96 (1H, m), 4.04 (2H, s), 4.90 (1H, s), 6.46 (1H, d, J=16.1 Hz), 7.16 (1H, m), 7.31 (4H, m), 7.46 (1H, d, J=15.7 Hz), 7.51 (3H, m), 7.73 (2H, d, J=7.3 Hz);
MASS (ES+): m/e 304 (M+1).

10 Preparation 278

Compound (278) was obtained from Compound (262) according to a manner similar to Preparation 277 (896 mg). The obtained Compound (278) was used in Example 59.

- 15 ¹H-NMR (300 MHz, CDCl₃, δ): 1.48-1.86 (6H, m), 3.58-3.69 (1H, m), 3.81 (3H, s), 3.91-4.09 (3H, m), 4.95 (1H, br.s), 6.09-6.30 (1H, m), 7.17-7.31 (6H, m), 7.33-7.43 (4H, m), 7.74-7.84 (2H, m);
MASS (ES+): m/z 462 (M+1).

Preparation 279

- 20 Compound (279) was obtained from Compound (263) according to a manner similar to Preparation 277 (1.67 g). The obtained Compound (278) was used in Example 70.

- 25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.53 (1H, m), 3.76 (3H, s), 3.94 (1H, m), 4.08 (2H, s), 4.91 (1H, s), 6.47 (1H, d, J=15.8 Hz), 6.94 (2H, d, J=8.8 Hz), 7.33 (2H, d, J=8.1 Hz), 7.43 (1H, s), 7.46 (1H, d, J=15.8 Hz), 7.54 (2H, d, J=7.7 Hz), 7.64 (2H, d, J=8.8 Hz);
MASS (ES+): m/z 434 (M+1).

Preparation 280

- 30 Compound (280) was obtained from Compound (264) according to a manner similar to Preparation 277 (758 mg). The obtained Compound (280) was used in Example 71.

- 35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.52 (1H, m), 3.78 (3H, s), 3.95 (1H, m), 4.04 (2H, s), 4.90 (1H, s), 6.46 (1H, d, J=16.1 Hz), 6.74 (1H, d, J=8.1 Hz), 7.20-7.33 (5H, m), 7.46 (1H, d, J=15.8 Hz), 7.51 (1H, s), 7.52 (2H, d, J=8.1 Hz);
MASS (ES+): m/e 434 (M+1).

Preparation 281

Compound (281) was obtained from Compound (265) according to a manner similar to Preparation 277 (729 mg). The obtained Compound (281) was used in Example 72.

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.53 (3H, br.s), 1.69 (3H, br.s), 3.53 (1H, m), 3.88 (3H, s), 3.95 (1H, m), 4.05 (2H, s), 4.90 (1H, s), 6.46 (1H, d, $J=16.3$ Hz), 6.96 (1H, t, $J=7.3$ Hz), 7.03 (1H, d, $J=8.1$ Hz), 7.17 (1H, t, $J=8.1$ Hz), 7.32 (2H, d, $J=8.1$ Hz), 7.42 (1H, br.s), 7.46 (1H, d, $J=16.1$ Hz), 7.52 (2H, d, $J=8.1$ Hz);
- 10 MASS (ES+): m/z 434 (M+1).

Preparation 282

Compound (282) was obtained from Compound (266) according to a manner similar to Preparation 277 (148 mg). The obtained Compound (282) was used in Example 73.

- 15 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.55 (1H, m), 3.95 (1H, m), 4.02 (2H, s), 4.91 (1H, s), 6.48 (1H, d, $J=16.1$ Hz), 7.32 (1H, t, $J=7.0$ Hz), 7.34 (2H, d, $J=8.1$ Hz), 7.46 (2H, t, $J=7.3$ Hz), 7.46 (1H, d, $J=15.4$ Hz), 7.54 (2H, d, $J=8.1$ Hz), 7.71 (2H, d, $J=7.3$ Hz);
- 20 MASS (ES+): m/e 482 (M+1).

Preparation 283

Compound (283) was obtained from Compound (267) according to a manner similar to Preparation 277 (166 mg). The obtained Compound (283) was used in Example 74.

- 25 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.53 (1H, m), 3.69 (1H, m), 4.02 (2H, s), 4.90 (1H, s), 6.47 (1H, d, $J=16.1$ Hz), 7.32 (1H, t, $J=8.1$ Hz), 7.33 (2H, d, $J=8.1$ Hz), 7.47 (2H, t, $J=15.8$ Hz), 7.47 (1H, d, $J=8.1$ Hz), 7.54 (2H, d, $J=8.4$ Hz), 7.70 (2H, d, $J=8.1$ Hz);
- 30 MASS (ES+): m/e 438 (M+1).

Preparation 284

Compound (284) was obtained from Compound (268) according to a manner similar to Preparation 277 (811 mg). The obtained Compound (283) was used in Examples 90 and 111.

- 35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.51-1.67 (6H, br), 2.28 (6H, s), 3.60 (1H, m), 3.94 (1H, m), 4.12 (2H, s), 4.71 (1H, s), 7.39 (3H, m),

7.59 (2H, d, J=6.6 Hz), 7.61 (1H, d, J=16.8 Hz);

MASS (ES+): m/e 461 (M+1).

Preparation 285

Compound (285) was obtained from Compound (269) according to
5 a manner similar to Preparation 277 (105 mg). The obtained
Compound (285) was used in Example 75.

¹H-NMR (300 MHz, CDCl₃, δ): 1.45 (3H, br.), 1.59 (3+4H, br.), 1.84
(2H, br.), 2.48 (4H, br.), 3.64 (1H, m), 3.71 (2H, s), 3.97 (1H, m),
4.12 (2H, s), 5.01 (1H, s), 6.32 (1H, d, J=16.1 Hz), 7.26 (1+2H, m),
10 7.39 (2+2H, m), 7.49 (1H, d, J=16.5 Hz), 7.59 (2H, d, J=7.0 Hz);

MASS (ES+): m/e 501 (M+1).

Preparation 286

Compound (286) was obtained from Compound (270) according to
a manner similar to Preparation 277 (129 mg). The obtained
15 Compound (286) was used in Example 76.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br.), 1.69 (3H, br.), 2.40
(4H, br.), 3.52 (1H, m), 3.57 (4H, br.), 3.95 (1H, m), 4.00 (2H, s),
4.90 (1H, s), 6.46 (1H, d, J=15.8 Hz), 7.23 (1H, br.), 7.33 (2H, d,
J=8.4 Hz), 7.36 (2H, br.), 7.46 (1H, d, J=15.8 Hz), 7.51 (2H, d,
20 J=8.1 Hz), 7.70 (2H, br.);

MASS (ES+): m/e 503 (M+1).

Preparation 287

Compound (287) was obtained from Compound (271) according to
a manner similar to Preparation 277 (108 mg). The obtained
25 Compound (287) was used in Example 77.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br.), 1.69 (3H, br.), 3.54
(1H, m), 3.79 (3H, s), 3.94 (1H, m), 3.98 (2H, s), 4.90 (1H, s),
6.47 (1H, d, J=16.1 Hz), 7.04 (2H, d, J=8.8 Hz), 7.33 (2H, d, J=7.7
Hz), 7.47 (1H, d, J=15.4 Hz), 7.53 (2H, d, J=7.7 Hz), 7.61 (2H, d,
30 J=8.8 Hz);

MASS (ES+): m/e 468 (M+1).

Preparation 288

Compound (288) was obtained from Compound (272) according to
a manner similar to Preparation 277 (215 mg). The obtained
35 Compound (288) was used in Example 80.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br.), 1.69 (3H, br.), 3.53

(1H, m), 3.94 (1H, m), 4.90 (1H, s), 5.25 (2H, s), 6.50 (1H, d, J=16.1 Hz), 7.18 (1H, t, J=7.3 Hz), 7.32 (3H, m), 7.38 (1H, d, J=15.4 Hz), 7.44 (1H, d, J=7.3 Hz), 7.52 (1H, s), 7.56 (1H, s), 7.73 (3H, m), 7.86 (1H, s);

5 MASS (ES+): m/e 404 (M+1).

Preparation 289

Compound (289) was obtained from Compound (273) according to a manner similar to Preparation 277 (419 mg). The obtained Compound (289) was used in Example 81.

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br.), 1.69 (3H, br.), 3.52 (1H, m), 3.95 (1H, m), 4.91 (1H, s), 5.26 (2H, s), 6.48 (1H, d, J=15.8 Hz), 7.19 (1H, d, J=7.7 Hz), 7.43 (5H, m), 7.51 (1H, s), 7.65 (2H, d, J=8.8 Hz), 7.86 (1H, s);

MASS (ES+): m/e 452 (M+1).

15 Preparation 290

Compound (290) was obtained from Compound (274) according to a manner similar to Preparation 277 (43 mg). The obtained Compound (290) was used in Example 84.

20 ¹H-NMR (300 MHz, CDCl₃, δ): 1.60 (3H, br.), 1.81 (3H, br.), 6.64 (1H, m), 3.82 (3H, s), 3.97 (1H, s), 4.03 (2H, s), 5.00 (1H, br.), 6.28 (1H, br.), 7.12 (2H, br.), 7.32 (2H, br.), 7.39 (3H, m), 7.54 (1H, br.), 7.79 (2H, br.);

MASS (ES+): m/e 462 (M+1).

Preparation 291

25 Compound (291) was obtained from Compound (275) according to a manner similar to Preparation 277 (85 mg). The obtained Compound (291) was used in Example 85.

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.53 (1H, m), 3.94 (2H, s), 3.97 (1H, m), 4.91 (1H, s), 6.45 (1H, d, J=15.3 Hz), 6.56 (1H, s), 7.27 (2H, d, J=8.1 Hz), 7.47 (1H, d, J=16.1 Hz), 7.50 (2H, d, J=8.8 Hz);

MASS (ES+): m/e 384 (M+1).

Preparation 292

35 Compound (292) was obtained from Compound (276) according to a manner similar to Preparation 277 (42 mg). The obtained Compound (292) was used in Example 110.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.30 (6H, m), 1.62-2.00 (10H, m), 2.51 (1H, m), 3.67 (1H, m), 4.02 (1H, m), 4.02 (2H, s), 5.01 (1H, s), 6.37 (1H, br.), 6.58 (1H, s, $J=8.0$ Hz), 7.23 (2H, d, $J=7.7$ Hz), 7.43 (2H, d, $J=16.1$ Hz), 7.62 (1H, d);

5 MASS (ES+): m/z 410 (M+1).

Preparation 293

To a suspension of Compound (245) (2.0 g) in MeCN (20 mL) was added N-bromosuccinimide (988 mg) at 5°C and the mixture was stirred for 0.5 hr at 5°C. The mixture was poured into 5% aqueous
10 NaHSO_3 solution (100 mL), and the mixture was extracted with AcOEt. The organic layer was washed with saturated aqueous NaHCO_3 solution and brine, and dried over Na_2SO_4 . The solvent was removed in vacuo and the residual brown oil was purified by Silica gel flash column chromatography eluting with Hexane: AcOEt = 1:1 to give Compound
15 (293) (1.68 g) as orange form.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 4.13 (2H, s), 6.34 (1H, d, $J=16.1$ Hz), 7.28 (2H, d, $J=8.1$ Hz), 7.31 (1H, t, $J=7.4$ Hz), 7.40 (2H, t, $J=7.4$ Hz), 7.47 (2H, d, $J=8.1$ Hz), 7.54 (1H, d, $J=16.1$ Hz), 7.57 (2H, d, $J=7.4$ Hz);

20 MASS (ES+): m/e 439 (M+1).

Preparation 294

To a solution of Compound (245) (1.0 g) were added N,N-dimethylamine hydrochloride (339 mg) and paraformaldehyde (125 mg), and the mixture was heated at 90°C for 1 hr. The reaction mixture
25 was poured into saturated aqueous NaHCO_3 solution and extracted with AcOEt. The organic phase was washed with saturated aqueous NaHCO_3 solution, water and brine, and dried over Na_2SO_4 . The solvent was removed in vacuo to give Compound (294) (742 mg) as yellow form.

30 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.53 (9H, s), 2.23 (6H, s), 3.58 (2H, s), 4.16 (2H, s), 6.35 (1H, d, $J=15.8$ Hz), 7.29 (1H, t, $J=8.1$ Hz), 7.30 (2H, d, $J=8.1$ Hz), 7.39 (2H, t, $J=8.1$ Hz), 7.48 (2H, t, $J=8.1$ Hz), 7.57 (1H, d, $J=16.1$ Hz), 7.58 (1H, br.);

MASS (ES+): m/e 418 (M+1).

35 Preparation 295

Compound (295) was obtained from Compound (245) according to

a manner similar to Preparation 294 (243 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.44 (2H, br.), 1.53 (9+4H, br.), 2.42 (4H, br.), 3.63 (2H, s), 4.11 (2H, s), 6.34 (1H, d, J=15.8 Hz), 7.27 (1H, t, J=8.1 Hz), 7.28 (2H, d, J=8.1 Hz), 7.37 (2H, t, J=8.1 Hz), 7.45 (2H, d, J=8.4 Hz), 7.55 (1H, d, J=15.7 Hz), 7.58 (2H, d, J=8.1 Hz);

MASS (ES+): m/e 458 (M+1).

Preparation 296

Compound (296) was obtained from Compound (245) according to a manner similar to Preparation 294 (297 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 2.48 (4H, br.), 3.64 (2H, s), 3.68 (4H, br.), 4.15 (2H, s), 6.35 (1H, d, J=16.1 Hz), 7.28 (1H, t, J=8.1 Hz), 7.29 (2H, d, J=8.1 Hz), 7.39 (2H, t, J=7.5 Hz), 7.48 (2H, d, J=8.4 Hz), 7.56 (2H, br.), 7.56 (1H, d, J=16.1 Hz);

MASS (ES+): m/e 460 (M+1).

Preparation 297

To a stirred solution of Compound (254) (4.0 g) in DMF (40 mL) were added acrylic acid t-butyl ester (8.13 mL), palladium(II) acetate (125 mg), triphenylphosphine (583 mg) and N,N-diisopropylethylamine (3.2 mL). The mixture was stirred at 100°C for 1 hr. The resulting mixture was poured into saturated aqueous NaHCO₃ solution and extracted with AcOEt. The organic phase was washed with saturated aqueous NaHCO₃ solution and brine, and dried over Na₂SO₄. The solvent was removed in vacuo and purified by silica gel column chromatography eluted with AcOEt to give Compound (297) (2.56 g) as colorless solid.

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 4.18 (2H, s), 6.34 (1H, d, J=16.1 Hz), 7.21 (1H, s), 7.25 (1H, m), 7.28 (2H, d, J=8.4 Hz), 7.37 (2H, t, J=7.4 Hz), 7.47 (2H, d, J=8.4 Hz), 7.55 (1H, d, J=16.1 Hz);

MASS (ES+): m/e 361 (M+1).

Preparation 298

To a solution of 4-iodophenol (1.0 g) in DMF (10 mL) was added K₂CO₃ (powder 325 mesh, manufactured by Aldrich, 691 mg) was added, and the mixture was stirred for 15 min. To the mixture was added ethyl 4-bromobutanoate (0.722 mL), and the mixture was

stirred for 65 hrs at ambient temperature. The resulting mixture was poured into saturated aqueous ammonium chloride solution and extracted with AcOEt. The organic phase was sequentially washed with water, saturated aqueous ammonium chloride solution, water and brine, and dried over Na₂SO₄. The solvent was removed in vacuo, and residual colorless oil was purified by silica gel column chromatography eluting with AcOEt and hexane (1:4) to give Compound (298) (1.32 g) as colorless oil.

¹H-NMR (300 MHz, CDCl₃, δ): 1.26 (3H, t, J=7.1 Hz), 2.10 (2H, quint., J=6.7 Hz), 2.50 (2H, t, J=7.3 Hz), 3.97 (2H, t, J=6.0 Hz), 4.14 (2H, q, J=7.1 Hz), 6.66 (2H, d, J=9.2 Hz), 7.54 (2H, d, J=9.2 Hz);

MASS: Not Detected.

Preparation 299

Compound (299) was obtained from 4-iodophenol according to a manner similar to Preparation 298 (1.70 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.26 (3H, t, J=7.1 Hz), 1.81 (4H, m), 2.38 (2H, m), 3.94 (2H, m), 4.14 (2H, q, J=7.2 Hz), 6.66 (2H, d, J=9.2 Hz), 7.54 (2H, d, J=9.2 Hz);

MASS: Not Detected.

20 Preparation 300

To a solution of Compound (298) (1.32 g) in dioxane (13 mL) was added 1N NaOH aq solution (11.9 mL) and the mixture was stirred for 15 hrs. The organic solvent was removed in vacuo, and the pH value of residual aqueous phase was adjusted to 3 with 1N HCl. The precipitate was collected by filtration and dried in vacuo to give Compound (300) (1.047 g) as colorless powder.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.91 (2H, quint., J=6.8 Hz), 2.36 (2H, t, J=7.3 Hz), 3.95 (2H, t, J=6.4 Hz), 6.78 (2H, d, J=8.9 Hz), 7.58 (2H, d, J=9.2 Hz);

30 MASS (ES+): Not Detected.

Preparation 301

Compound (301) was obtained from Compound (299) according to a manner similar to Preparation 300 (883 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.56-1.76 (4H, m), 2.26 (2H, t, J=7.1 Hz), 3.94 (2H, t, J=6.2 Hz), 6.78 (2H, d, J=8.8 Hz), 7.57 (2H, d, J=8.8 Hz);

MASS: Not Detected.

Preparation 302

Compound (302) was obtained from Compound (300) according to a manner similar to Preparation 297 (1.026 g).

5 ¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 2.13 (2H, quint., J=6.6 Hz), 2.58 (2H, t, J=7.2 Hz), 4.05 (2H, t, J=6.0 Hz), 6.24 (1H, d, J=15.8 Hz), 6.87 (2H, d, J=8.8 Hz), 7.44 (2H, d, J=8.8 Hz), 7.53 (2H, d, J=16.1 Hz);

MASS Not Detected.

10 Preparation 303

Compound (303) was obtained from Compound (301) according to a manner similar to Preparation 297 (635 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 1.85 (4H, m), 2.45 (2H, t, J=5.9 Hz), 4.00 (2H, t, J=5.9 Hz), 6.24 (1H, d, J=16.1 Hz), 6.87 (2H, d, J=8.8 Hz), 7.44 (2H, d, J=8.8 Hz), 7.54 (1H, d, J=15.8 Hz);

15 MASS: Not Detected.

Preparation 304

Compound (304) was obtained from 4-(4-iodophenyl)butanoic acid according to a manner similar to Preparation 297 (1.68 mg).

20 ¹H-NMR (300 MHz, CDCl₃, δ): 1.53 (9H, s), 1.97 (2H, quint., J=7.5 Hz), 2.38 (2H, t, J=7.3 Hz), 2.69 (2H, t, J=7.5 Hz), 6.33 (1H, d, J=15.8 Hz), 7.19 (2H, d, J=8.1 Hz), 7.44 (2H, d, J=8.1 Hz), 7.56 (2H, d, J=15.8 Hz);

MASS: Not Detected.

25 Preparation 305

Compound (305) was obtained from Compound (318) described later according to a manner similar to Preparation 297 (2.45 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.43 (1H, s), 1.53 (9H, s), 1.69 (1H, s), 1.93 (1H, s), 2.59 (1H, s), 6.33 (1H, d, J=16.4 Hz), 7.10 (2H, d, J=8.1 Hz), 7.43 (2H, d, J=8.1 Hz), 7.55 (1H, d, J=15.8 Hz);

30 MASS (ES⁻): m/e 287 (M-1).

Preparation 306

To a solution of Compound (302) in DMF (3 mL) were added tert-butyl 2-aminophenylcarbamate (224 mg), HOBt (172 mg) and N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDCI·HCl, 224 mg), and the mixture was stirred at ambient

35

temperature for 2 hrs. The mixture was poured into saturated aqueous NaHCO₃ solution and extracted with ethyl acetate. The organic phase was sequentially washed with saturated NH₄Cl solution, saturated NaHCO₃ solution and brine, and dried over Na₂SO₄. The solvent was removed in vacuo and the residue was purified by preparative thin layer chromatography (chloroform: methanol = 10:1) to give Compound (306) (379 mg) as pale yellow form.

¹H-NMR (300 MHz, CDCl₃, δ): 1.50 (9H, s), 1.53 (9H, s), 2.23 (2H, quint., J=6.6 Hz), 2.60 (2H, t, J=7.0 Hz), 4.09 (2H, t, J=6.0 Hz), 6.24 (1H, d, J=15.8 Hz), 6.88 (2H, d, J=8.8 Hz), 7.16 (2H, m), 7.36 (1H, m), 7.44 (2H, d, J=8.8 Hz), 7.50 (1H, m), 7.53 (1H, d, J=15.8 Hz);

MASS (ES+): m/e 497 (M+1).

Preparation 307

Compound (307) was obtained from Compound (303) according to a manner similar to Preparation 306 (395 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.50 (9H, s), 1.53 (9H, s), 1.92 (4H, m), 2.47 (2H, m), 4.03 (2H, m), 6.24 (1H, d, J=16.1 Hz), 6.87 (2H, d, J=8.8 Hz), 7.17 (2H, m), 7.35 (1H, m), 7.44 (2H, d, J=8.8 Hz), 7.50 (1H, m), 7.53 (1H, d, J=16.1 Hz);

MASS (ES+): m/e 511 (M+1).

Preparation 308

Compound (308) was obtained from Compound (304) according to a manner similar to Preparation 306 (294 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.48 (9H, s), 1.53 (9H, s), 2.07 (2H, quint., J=7.7 Hz), 2.40 (2H, t, J=7.9 Hz), 2.73 (2H, t, J=7.9 Hz), 6.33 (1H, d, J=7.9 Hz), 7.18 (2H, m, J=16.1 Hz), 7.21 (2H, d), 7.35 (1H, m, J=8.1 Hz), 7.44 (2H, d), 7.50 (1H, m, J=8.1 Hz), 7.56 (1H, d, J=16.1 Hz);

MASS (ES+): m/e 481 (M+1).

Preparation 309

To a stirred solution of Compound (306) (379 mg) in AcOH (4 mL) was added 1N HCl in AcOH (3.82 mL), and the mixture was heated at 120°C for 2 hrs. The reaction mixture was cooled to ambient temperature and diluted with AcOEt. The precipitate was collected by filtration to give Compound (309) (199 mg) as pale yellow powder.

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.36 (2H, quint., J=6.5 Hz), 3.30 (2H, t, J=7.5 Hz), 4.16 (2H, t, J=5.9 Hz), 6.37 (1H, d, J=16.1 Hz), 6.79 (2H, d, J=8.4 Hz), 7.52 (1H, d, J=16.5 Hz), 7.55 (2H, m), 7.59 (2H, d, J=8.8 Hz), 7.79 (2H, m);

5 MASS (ES+): m/e 323 (M+1).

Preparation 310

Compound (310) was obtained from Compound (307) according to a manner similar to Preparation 309 (205 mg).

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.84 (2H, m), 2.03 (2H, m), 3.21 (2H, t, J=7.6 Hz), 4.08 (2H, t, J=5.9 Hz), 6.38 (1H, d, J=16.1 Hz), 6.96 (2H, d, J=8.8 Hz), 7.53 (2H, m), 7.54 (1H, d, J=15.8 Hz), 7.63 (2H, d, J=9.1 Hz), 7.78 (2H, m);

MASS (ES+): m/e 337 (M+1).

Preparation 311

15 Compound (311) was obtained from Compound (308) according to a manner similar to Preparation 309 (174 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.19 (2H, quint., J=7.1 Hz), 2.75 (2H, t, J=7.3 Hz), 3.12 (2H, t, J=7.5 Hz), 6.48 (1H, d, J=16.1 Hz), 7.30 (2H, d, J=7.7 Hz), 7.52 (2H, m), 7.56 (1H, d, J=15.0 Hz), 7.62 (2H, 20 d, J=8.4 Hz), 7.76 (2H, m);

MASS (ES+): m/e 307 (M+1).

Preparation 312

To a stirred solution of Compound (309) (234 mg) in DMF (2 mL) was added O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (71.5 mg), 25 HOBT (97.4 mg) and EDCI (112 mg), and the resulting mixture was stirred at ambient temperature for 16 hrs. To the reaction mixture was added saturated aqueous NaHCO₃ solution, and the precipitate was collected by filtration and washed with water. The obtained powder was dried in vacuo to give Compound (312) (161 mg) as 30 colorless powder. The obtained compound (312) was used in Example 103.

¹H-NMR (300 MHz, CDCl₃, δ): 1.62 (3H, br.), 1.84 (3H, br.), 2.38 (2H, quint., J=7.1 Hz), 3.13 (2H, t, J=7.1 Hz), 3.65 (1H, m), 3.98 (1H, m), 4.05 (2H, t, J=6.0 Hz), 5.02 (1H, s), 6.78 (2H, d, J=8.8 Hz), 35 7.23 (2H, m), 7.37 (2H, d, J=8.4 Hz), 7.56 (2H, br.), 7.63 (1H, d, J=16.1 Hz);

MASS (ES+): m/e 422 (M+1).

Preparation 313

Compound (313) was obtained from Compound (310) according to a manner similar to Preparation 312 (197 mg). The obtained

5. compound (313) was used in Example 104.

¹H-NMR (300 MHz, CDCl₃, δ): 1.64 (3H, br.), 1.84 (3H, br.), 1.91 (2H, m), 2.07 (2H, m), 3.01 (2H, t, J=7.7 Hz), 3.65 (1H, m), 3.97 (1H, m), 3.99 (2H, t, J=6.3 Hz), 5.02 (1H, s), 6.81 (2H, d, J=8.8 Hz), 7.22 (2H, m), 7.40 (2H, d, J=8.1 Hz), 7.53 (2H, br.m), 7.67 (2H, d, J=15.0 Hz);

10 MASS (ES+): m/e 436 (M+1).

Preparation 314

Compound (314) was obtained from Compound (311) according to a manner similar to Preparation 312 (177 mg). The obtained

15 compound (314) was used in Example 105.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.53 (3H, br.), 1.69 (3H, br.), 2.09 (2H, quint., J=7.7 Hz), 2.69 (2H, t, J=7.7 Hz), 2.81 (2H, t, J=7.4 Hz), 3.55 (1H, m), 3.96 (1H, m), 4.91 (1H, s), 6.47 (1H, d, J=16.1 Hz), 7.10 (2H, m), 7.29 (2H, d, J=8.1 Hz), 7.46 (1H, d, J=15.8 Hz), 7.47 (2H, m), 7.51 (2H, d, J=7.7 Hz);

20 MASS (ES+): m/e 406 (M+1).

Preparation 315

Compound (315) was obtained from Compound (320) described later according to a manner similar to Preparation 312 (287 mg).

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.52 (3H, br.), 1.66 (1H, m), 1.69 (3H, br.), 1.82 (1H, m), 2.40 (1H, m), 2.56 (1H, m), 3.50 (1H, m), 3.95 (1H, m), 4.90 (1H, s), 6.47 (1H, d, J=15.8 Hz), 7.11 (2H, m), 7.25 (2H, d, J=8.1 Hz), 7.41 (1H, d, J=16.1 Hz), 7.44 (2H, m), 7.50 (2H, d, J=8.8 Hz);

30 MASS (ES+): m/e 404 (M+1).

Preparation 316

Compound (316) was obtained from Compound (335) described later according to a manner similar to Preparation 312 (55 mg). The obtained compound (316) was used in Example 100.

35 ¹H-NMR (300 MHz, CDCl₃, δ): 1.60-1.90 (6H, m), 3.64-3.73 (1H, m), 3.80 (3H, s), 3.94-4.04 (1H, m), 4.13 (2H, s), 4.82 (2H, s), 4.97-

5.07 (1H, m), 6.81-7.31 (8H, m), 7.45 (2H, d, J=8 Hz), 7.71 (1H, d, J=16 Hz);

MASS (ES+): m/z 516 (M+1).

Preparation 317

5 Compound (317) was obtained from Compound (202) according to a manner similar to Preparation 312 (177 mg). The obtained compound (317) was used in Example 101.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.49-2.02 (21H, m), 3.49-3.58 (1H, m), 3.88-4.01 (3H, m), 4.87-4.93 (1H, m), 6.45 (1H, d, J=16 Hz), 7.18
10 (1H, d, J=8 Hz), 7.26 (2H, d, J=8 Hz), 7.41-7.53 (3H, m).

MASS (ES+): m/z 462 (M+1).

Preparation 318

To a stirred solution of trans-2-phenylcyclopropanecarboxylic acid (5.0 g) in acetic acid (30 mL)
15 were added periodic acid (3.13 g), iodine (3.13 g) and concentrated sulfuric acid (3.62 mL), and the mixture was stirred at 75°C for 5 hrs. The resulting mixture was poured into 5% aqueous NaHSO₃ solution (100 mL), and the precipitate was collected by filtration. The filtrate was extracted with AcOEt, and the organic phase was
20 sequentially washed with 5% aqueous NaHSO₃ solution, saturated aqueous NH₄Cl solution and brine, and dried over Na₂SO₄. The solvent was removed in vacuo, and the residual solid was combined with the powder obtained by filtration, and dissolved in AcOEt. The precipitate was filtered off, and the filtrate was concentrated
25 and crystallized from ethyl acetate and hexane (20 mL: 40 mL) to give Compound (318) (3.88 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.36 (1H, m), 1.66 (1H, m), 1.87 (1H, m), 2.53 (1H, m), 6.86 (2H, d, J=8.4 Hz), 7.60 (2H, d, J=8.4 Hz);

MASS (ES-): m/e 287 (M-1).

30 Preparation 319

To a solution of Compound (305) (500 mg) in DMF (5 mL) were added 1,2-benzenediamine (206 mg), HOBt (305 mg) and EDCI·HCl (432 mg), and the mixture was stirred for 2 hrs at ambient temperature. The mixture was poured into saturated aqueous NaHCO₃ solution and
35 extracted with AcOEt. The organic phase was sequentially washed with saturated aqueous NaHCO₃ solution, water and brine, and dried

over Na₂SO₄. The solvent was removed in vacuo, and the residue was purified by preparative thin layer chromatography (chloroform: methanol = 10: 1). The obtained amorphous solid was triturated with isopropyl ether to give Compound (319) (341 mg) as colorless powder.

¹H-NMR (300 MHz, CDCl₃, δ): 1.40 (1H, m), 1.53 (9H, s), 1.77 (1H, m), 1.84 (1H, m), 2.62 (1H, m), 3.85 (1H, br.), 6.34 (1H, d, J=15.8 Hz), 6.80 (2H, m), 7.07 (2H, m), 7.13 (2H, d, J=8.4 Hz), 7.34 (2H, s), 7.45 (2H, d, J=8.4 Hz), 7.56 (1H, d, J=16.1 Hz);

10 MASS (ES+): m/e 379 (M+1).

Preparation 320

A solution of Compound (319) (341 mg) in AcOH (3.5 mL) was added 1N-hydrogen chloride in AcOH (3.6 mL) was heated at 110°C for 2 hr. The resulting mixture was allowed to cool to ambient temperature, diluted with ethyl acetate. The precipitate was collected by filtration and dried to give Compound (320) (242 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.03 (1H, m), 2.14 (1H, m), 2.70 (1H, m), 2.92 (1H, m), 6.53 (1H, d, J=15.8 Hz), 7.35 (2H, d, J=8.4 Hz), 7.49 (2H, m), 7.58 (1H, d, J=16.1 Hz), 7.68 (2H, d, J=8.1 Hz), 7.73 (2H, m);

20 MASS (ES+): m/e 305 (M+1).

Preparation 321

To a stirred solution of Compound (217) (350 mg) in methanol (3 mL) was added 1N sodium hydroxide (1.65 mL) in an ice bath. After 1 hr, the mixture was allowed to warm to ambient temperature and stirred at the same temperature for 16 hrs. To this mixture was added concentrated hydrochloric acid to acidify the mixture, and the resulting mixture was concentrated in vacuo. The residue was dissolved in N,N-dimethylformamide (5 mL), and to this solution were successively added O-(tetrahydro-2H-pyran-2-yl)hydroxylamine (167 mg), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide (222 mg) and 1-hydroxybenzotriazole (193 mg) at ambient temperature. The resulting mixture was stirred at the same temperature for two days, extracted with ethyl acetate (100 mL), and successively washed with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous sodium sulfate and

concentrated in vacuo. The residue was purified by silica gel flash chromatography eluting with a mixture of methanol and ethyl acetate (5:95 v/v) to afford Compound (321) (387 mg) as a colorless amorphous solid. The obtained Compound (321) was used in Example 82.

$^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.43–1.82 (6H, m), 3.43–3.59 (1H, m), 3.84–4.07 (1H, m), 4.89 (1H, br.s), 5.36 (2H, s), 6.47 (1H, d, $J=16.1$ Hz), 7.00–7.09 (3H, m), 7.33–7.59 (10H, m);
MASS (ES+): m/z 404 (M+1).

10 Preparation 322

Compound (322) was obtained as a mixture of two regioisomers from Compound (218) according to a manner similar to Preparation 321 (316 mg in total). The obtained Compound (322) was used in Example 115.

15 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): (for a mixture of two regioisomers) 1.34–1.78 (6H, m), 2.03 (1.2H, s), 2.06 (1.8H, s), 3.47–3.59 (1H, m), 3.85–4.04 (1H, m), 4.83–4.95 (1H, m), 5.13 (1.2H, s), 5.19 (0.8H, s), 6.49 (1H, d, $J=16.4$ Hz), 6.67 (0.4H, s), 6.85 (0.6H, s), 7.13 (0.8H, d, $J=8.1$ Hz), 7.26 (1.2H, d, $J=8.1$ Hz), 7.39–7.72 (4H, 20 m);
MASS (ES+): (for a mixture of two regioisomers) m/z 342 (M+1).

Preparation 323

To a stirred solution of Compound (326) (100 mg, described later in Preparation 326) in N,N -dimethylformamide (2 mL) were 25 added methylamine hydrochloride (23 mg), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide (56 mg) and 1-hydroxybenzotriazole (48 mg) at ambient temperature, and the resulting mixture was stirred at the same temperature for 20 hrs. The mixture was extracted with ethyl acetate (50 mL), and 30 successively washed with saturated aqueous sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated in vacuo to afford crude Compound (324) (86 mg) as a colorless amorphous solid. The Compound (324) was used in Example 83.

35 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 1.42–1.97 (6H, m), 2.85 (3H, br.d, $J=4.4$ Hz), 3.54–3.68 (1H, m), 3.89–4.07 (2H, m), 4.98 (2H, br.s), 6.05–

6.34 (1H, m), 6.95-7.73 (10H, m);

MASS (ES+): m/z 461 (M+1).

Preparation 324

To a solution of Compound (326) (60 mg, described later in
5 Preparation 326) in DMF (1 mL) were added piperidine (0.015 mL),
HOBt (23.6 mg) and EDCI HCl (33.4 mg), and the mixture was stirred
for 2 hrs at ambient temperature. The mixture was poured into
water and extracted with AcOEt, and the organic phase was
sequentially washed with saturated aqueous NH₄Cl solution,
10 saturated aqueous NaHCO₃ solution and brine, and dried over Na₂SO₄.
The solvent was removed in vacuo, and the residue was purified by
preparative thin layer chromatography (chloroform: methanol = 10:
1) to give Compound (324) (44 mg) as colorless amorphous solid.
The obtained Compound (324) was used in Example 88.

15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.25 (2H, br.), 1.53 (7H, br.), 1.69
(3H, br.), 3.25 (4H, br.), 3.53 (1H, m), 3.94 (1H, m), 4.03 (2H, s),
4.90 (1H, s), 6.46 (1H, d, J=15.4 Hz), 7.26 (1H, t, J=7.3 Hz), 7.36
(4H, m), 7.43 (1H, d, J=15.8 Hz), 7.53 (4H, m);
MASS (ES+): m/e 515 (M+1).

20 Preparation 325

Compound (325) was obtained from Compound (326) described
later according to a manner similar to Preparation 325 (57 g). The
obtained compound (325) was used in Example 89.

¹H-NMR (300 MHz, CDCl₃, δ): 1.18 (6H, br.), 1.60 (3H, br.), 1.84 (3H,
25 br.), 3.65 (1H, m), 3.98 (1H, m), 4.09 (2H, s), 4.18 (1H, m), 5.00
(1H, s), 7.22 (1H, m), 7.39 (6H, m), 7.67 (3H, m);
MASS (ES+): m/e 489 (M+1).

Preparation 326

To a solution of Compound (290) (200 mg) in dioxane (2 mL)
30 was added 1N NaOH aqueous solution (1.3 mL), and the mixture was
heated at 70°C for 6 hrs. The organic solvent was removed in vacuo
and the pH value of aqueous base adjusted to 3 with 1N HCl. The
precipitate was collected by filtration and dried to give Compound
(326) (145 mg) as pale yellow powder. The obtained Compound (326)
35 was used in Example 91.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.53

(1H, m), 3.95 (1H, m), 4.05 (2H, s), 4.90 (1H, s), 6.46 (1H, d, J=16.4 Hz), 7.32-7.40 (5H, m), 7.46 (1H, d, J=16.1 Hz), 7.53 (2H, d, J=7.7 Hz), 7.86 (2H, br);
MASS (ES+): m/e 448 (M+1).

5 Preparation 327

To a suspension of lithium aluminum hydride in THF (5 mL) was added a solution of Compound (290) (500 mg) in THF (5 mL) and the mixture was refluxed for 6 hrs. To the resulting solution was added water (50 mL) at ambient temperature, and the mixture was
10 stirred for 1 hr. The precipitate was collected by filtration, and the obtained powder was purified by silica gel column chromatography eluting with a mixture of chloroform and methanol (10: 1) to give Compound (327) (110 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.54 (9H, s), 3.68 (2H, s), 3.88 (3H, s),
15 4.69 (2H, d, J=4.0 Hz), 6.37 (1H, d, J=15.8 Hz), 6.60 (1H, br), 6.95 (2H, d, J=8.8 Hz), 7.33 (2H, d, J=8.1 Hz), 7.52 (2H, d, J=8.1 Hz), 7.58 (1H, d, J=16.1 Hz), 7.91 (2H, d, J=9.2 Hz);
MASS (ES+): m/e 434 (M+1).

Preparation 328

20 Compound (328) was obtained from Compound (50) according to a manner similar to Preparation 327 (493 mg). The obtained compound (328) was used in Example 97.

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.48-1.72 (6H, m), 3.49-3.57 (1H, m),
3.90-4.01 (1H, m), 4.18 (2H, s), 4.51-4.58 (2H, m), 4.87-4.92 (1H,
25 m), 6.47 (1H, d, J=16 Hz), 7.33-7.57 (8H, m);
MASS (ES+): m/z 408 (M+1).

Preparation 329

To a suspension of Compound (327) (400 mg) in dichloromethane (3 mL) was added triethylamine (0.154 mL) and
30 methanesulfonyl chloride (0.079 mL) at 5°C, and the mixture was stirred for 1 hr. The solvent was removed in vacuo, and the obtained benzyl chloride was dissolved in MeOH. To the solution was added sodium methoxide in MeOH (5 equivalent) at ambient temperature, and the mixture was stirred for 0.5 hr. The resulting
35 mixture was poured into saturated aqueous NH₄Cl solution and extracted with AcOEt. The organic phase was washed with water and

brine, and dried over Na_2SO_4 . The solvent was removed in vacuo, and the residue was purified by preparative thin layer chromatography (chloroform: methanol = 10: 1) to give Compound (329) (110 mg).

The obtained compound (329) was used in Example 93.

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.54 (3H, br.), 1.69 (3H, br.), 3.29 (3H, s), 3.53 (1H, m), 3.97 (1H, m), 4.05 (2H, s), 4.39 (2H, s), 4.90 (1H, s), 6.47 (1H, d, $J=15.8$ Hz), 7.25-7.61 (10H, m);
MASS (ES+): m/e 448 (M+1).

Preparation 330

- 10 To a solution of Compound (323) (1 g) in DMF (5 mL) were added N,O-dimethylhydroxylamine hydrochloride (240 mg) and HOBT (393 mg) and EDCI (451 mg), and the mixture was stirred at ambient temperature. To the resulting mixture was added saturated aqueous NaHCO_3 solution, and the precipitate was collected by filtration.

- 15 The obtained powder was washed with water and Et_2O to give Compound (330) (817 mg) as pale yellow powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.53 (3H, br.), 1.69 (3H, br.), 3.21 (3H, s), 3.53 (1H, m), 3.63 (3H, s), 3.95 (1H, m), 4.05 (2H, s), 4.90 (1H, s), 6.46 (1H, d, $J=16.1$ Hz), 7.30-7.60 (10H, m);

- 20 MASS (ES+): m/z 491 (M+1).

Preparation 331

- To a solution of Compound (330) (300 mg) in THF (5 mL) was added methyl magnesium iodide in Et_2O (3.64 mL, 0.84 mol/L, Kanto Chemical, Co., Inc.) and the mixture was heated at 70°C for 3 hrs.
- 25 Methylmagnesium iodide in Et_2O (0.84 mol/L, 3.64 mL) was then added thereto, and the mixture was additionally heated at 70°C for 2 hrs. The mixture was cooled, poured into water and extracted with AcOEt . The organic phase was successively washed with saturated aqueous NaHCO_3 solution, water and brine, and dried over Na_2SO_4 . The
- 30 solvent was removed in vacuo, and the residue was purified by preparative thin layer chromatography (chloroform: methanol = 10: 1) to give Compound (331) (67 mg) as orange form. The obtained Compound (331) was used in Example 94.

- 35 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.54 (3H, br.), 1.69 (3H, br.), 2.36 (3H, s), 3.55 (1H, m), 3.95 (1H, m), 4.22 (2H, s), 4.90 (1H, s), 6.48 (1H, d, $J=15.8$ Hz), 7.31-7.57 (10H, m);

MASS (ES+): m/z 446 (M+1).

Preparation 332

Compound (332) was obtained from Compound (326) according to a manner similar to Preparation 330 (124 mg). The obtained
5 compound (332) was used in Example 87.

Preparation 333

To a solution of Compound (51) (120 mg), pyrrolidine (0.029 mL) and 1-hydroxybenzotriazole (46.2 mg) in N,N-dimethylformamide (2.6 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
10 (65.5 mg) at 4°C. The mixture was warmed to ambient temperature and stirred for 8 hrs. The reaction mixture was added saturated NaHCO₃ (3 mL) and water (12 mL). A resulting precipitate was collected by filtration, and washed with water to give Compound (333) (95 mg).
15 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.47-1.93 (10H, m), 3.39-3.57 (5H, m), 3.87-4.01 (1H, m), 4.22 (2H, s), 4.88-4.93 (1H, m), 6.47 (1H, d, J=16 Hz), 7.28-7.70 (8H, m);
MASS (ES+): m/z 475 (M+1).

Preparation 334

20 To a mixture of Compound (328) (170 mg), N,N-diisopropylethylamine (0.16 mL) and THF (8.4 mL) was added methanesulfonyl chloride (0.068 mL) at 4°C. The reaction mixture was stirred for 3 hrs and diethylamine (0.432 mL) was added thereto. After stirring for 15 hrs at room temperature, the resulting
25 mixture was partitioned between ethyl acetate and H₂O. The organic layer was washed with saturated NaHCO₃ and brine, dried over MgSO₄, filtered and evaporated in vacuo to give Compound (334) (38 mg).
¹H-NMR (300 MHz, DMSO-d₆, δ): 1.03-1.22 (6H, m), 1.48-1.72 (6H, m), 3.31-3.44 (4H, m), 3.49-3.57 (1H, m), 3.88-4.03 (1H, m), 4.10-4.41
30 (4H, m), 4.87-4.95 (1H, m), 6.48 (1H, d, J=16 Hz), 7.22-7.87 (8H, m).
MASS (ES+): m/z 463 (M+1).

Preparation 335

Compound (335) was obtained from Compound (6) according to
35 manners similar to Preparations 199 and 229 (2.89 g).

¹H-NMR (300 MHz, CDCl₃, δ): 1.34 (3H, t, J=7 Hz), 3.64-4.08 (7H, m),

4.27 (2H, q, J=7 Hz), 4.53-4.64 (2H, m), 6.42 (1H, d, J=16 Hz),
6.81-6.92 (2H, m), 7.01 (1H, d, J=8 Hz), 7.27-7.35 (3H, m), 7.45-
7.53 (2H, m), 7.66 (1H, d, J=16 Hz);

MASS (ES-): m/z 410 (M-1).

5 Preparation 336

To a mixture of Compound (335) (1.44 g) in chloroform (12 mL) was added trifluoroacetic anhydride (1.48 mL) at 4°C. After stirring for 3 hrs at room temperature, the reaction mixture was evaporated in vacuo. To a mixture of the above product in N,N-
10 dimethylformamide (18 mL) was added ammonium acetate (405 mg) at 4°C. After stirring at 70°C for 2 hrs, the resulting mixture was partitioned between ethyl acetate and H₂O. The organic layer was washed with H₂O and brine, dried over MgSO₄, filtered and evaporated in vacuo. To a mixture of the above product in pyridine
15 (1 mL) was added phosphorus oxychloride (0.3 mL) at 4°C. After stirring at 90°C for 2 hrs, the resulting mixture was partitioned between ethyl acetate and H₂O. The organic layer was washed with H₂O and brine, dried over MgSO₄, filtered and evaporated in vacuo. The residue was purified by column chromatography on silica gel to
20 give Compound (336) (313 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.34 (3H, t, J=7 Hz), 3.79 (3H, s), 4.12 (2H, s), 4.19 (2H, q, J=7 Hz), 4.81 (2H, s), 6.41 (1H, d, J=16 Hz), 6.81-6.90 (4H, m), 7.11-7.19 (3H, m), 7.45 (2H, d, J=8 Hz), 7.65 (1H, d, J=16 Hz);

25 MASS (ES+): m/z 445 (M+1).

Preparation 337

To a mixture of Compound (336) (175 mg) in dioxane (4 mL) was added 1N sodium hydroxide (1.18 mL). After stirring at 80°C for 1 hr, the reaction mixture was added H₂O (20 mL) and acidified
30 with 1N hydrochloric acid (to pH 3-4). The resulting precipitate was collected by filtration and washed with H₂O to give Compound (337) (153 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 3.79 (3H, s), 4.14 (2H, s), 4.82 (2H, s), 6.42 (1H, d, J=16 Hz), 6.81-6.91 (4H, m), 7.11-7.21 (3H, m), 7.48
35 (2H, d, J=8 Hz), 7.73 (1H, d, J=16 Hz);

MASS (ES-): m/z 415 (M-1).

Preparation 338

To a mixture of (1Z)-2-(4-iodophenyl)-N'-hydroxyethaneimidamide (2.17 g) in EtOH (40 mL) was added ethyl propiolate (0.803 mL), and the mixture was heated under reflux for 6 hrs. The reaction mixture was evaporated in vacuo. The residue was added diphenylether (20 mL) and stirred at 200°C for 2 hrs. After cooling, the resulting mixture was purified by column chromatography on silica gel (eluting with CHCl₃: MeOH = 50: 1) to give Compound (338) (845 mg).

¹H-NMR (300 MHz, CDCl₃, δ): 1.30-1.43 (3H, m), 4.06-4.13 (2H, m), 4.26-4.43 (2H, m), 6.97-7.06 (2H, m), 7.61 (1H, s), 7.65-7.72 (2H, m);

MASS (ES+): m/z 357 (M+1).

Preparation 339

To a solution of Compound (338) (400 mg), palladium(II) acetate (12.6 mg) and triphenylphosphine (29.5 mg) in DMF (11 mL) were added acrylic acid (0.154 mL) and N,N-diisopropylethylamine (0.49 mL), and the mixture was stirred at 70°C for 6 hrs. The reaction mixture was partitioned between ethyl acetate and H₂O, and the inorganic layer was evaporated in vacuo. To a mixture of the obtained reaction product, O-tetrahydro-2H-pyran-2-ylhydroxylamine (1.5 eq.) and 1-hydroxybenzotriazole (1.5 eq.) in N,N-dimethylformamide (6 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.5 eq.) at 4°C. The mixture was warmed to ambient temperature and stirred for 6 hrs. To the reaction mixture were added saturated NaHCO₃ (6 mL) and water (24 mL), and the mixture was extracted with ethyl acetate. The organic layer was washed with H₂O and brine, dried over MgSO₄, filtered and evaporated in vacuo. The residue was purified by column chromatography on silica gel to give Compound (339) (98 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.25 (3H, t, J=7 Hz), 1.44-1.75 (6H, m), 3.45-3.59 (1H, m), 3.88-4.01 (1H, m) 4.01 (2H, s), 4.18 (2H, q, J=7 Hz), 4.87-4.93 (1H, m), 6.46 (1H, d, J=16 Hz), 7.24-7.78 (5H, m), 8.32 (1H, s);

MASS (ES+): m/z 400 (M+1).

Preparation 340

Compound (340) was obtained according to manners similar to Preparations 319 and 320 (184 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.95-2.08 (4H, m), 3.22-3.34 (4H, m), 4.50 (2H, s), 6.55 (1H, d, J=16 Hz), 6.62 (1H, s), 6.84 (1H, dd, J=2, 8 Hz), 7.48-7.75 (6H, m);

MASS (ES+): m/z 348 (M+1).

Example 1

To a stirred solution of Compound (5) (125 mg) in methanol (5 mL) was added hydrogen chloride methanol reagent 10 (0.5 mL, manufactured by Tokyo Kasei Kogyo Co., Ltd.), and the mixture was stirred at ambient temperature for 30 minutes. The solvent was evaporated in vacuo and the residue was triturated with the mixture of methanol and ethyl acetate (1:2) to give Compound E1 as a white solid (81 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.57 (2H, s), 6.50 (1H, d, J=15.7 Hz), 7.41-7.56 (5H, m), 7.60 (2x1H, d, J=8 Hz), 7.73-7.81 (2H, m), 10.84 (1H, br);

MASS (ES+): m/e 294.

Example 2

Compound E2 was obtained from Compound (13) according to a manner similar to Example 1 (79 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.76 (2H, s), 5.82 (2H, s), 6.49 (1H, d, J=16 Hz), 7.12-7.21 (2H, m), 7.26-7.34 (3H, m), 7.38-7.62 (7H, m), 7.73 (1H, dd, J=7, 1.5 Hz), 7.83 (1H, dd, J=7, 1.5 Hz);

MASS (ES+): m/e 384.

Example 3

Compound E3 was obtained from Compound (19) according to a manner similar to Example 1 (1.74 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.28 (2x1H, t, J=7.5 Hz), 3.47 (2x1H, t, J=7.5 Hz), 6.45 (1H, d, J=16 Hz), 7.32 (2x1H, d, J=8 Hz), 7.41 (1H, d, J=16 Hz), 7.46-7.60 (4H, m), 7.73-7.83 (2H, m), 10.80 (1H, s), 15.10 (1H, br);

MASS (ES+): m/e 308.

Example 4

Compound E4 was obtained from Compound (22) according to a manner similar to Example 1 (377 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 5.79 (2H, s), 6.50 (1H, d, J=16 Hz), 7.43 (1H, d, J=16 Hz), 7.48-7.64 (6H, m), 7.86-7.94 (2H, m), 9.84 (1H, s);

MASS (ES+): m/e 294.

5 Example 5

Compound E5 was obtained from Compound (25) according to a manner similar to Example 1 (102 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.31 (1H, m), 3.70 (1H, dd, J=13, 8 Hz), 4.74 (1H, br-t, J=8 Hz), 5.34 (1H, d, J=17 Hz), 5.41 (1H, d, J=17 Hz), 6.37 (1H, d, J=15.5 Hz), 6.73 (2x1H, d, J=6.5 Hz), 7.07-7.45 (16H, m), 7.71 (1H, d, J=7.5 Hz), 9.02 (1H, brs), 10.73 (1H, brs);

MASS (ES+): m/e 474.

Example 6

15 Compound E6 was obtained from Compound (29) according to a manner similar to Example 1 (1.88 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.57 (2H, s), 6.51 (1H, d, J=16 Hz), 7.40-7.56 (6H, m), 7.68 (1H, s), 7.73-7.81 (2H, m), 10.88 (1H, s);

MASS (ES+): m/e 294.

20 Example 7

Compound E7 was obtained from Compound (32) according to a manner similar to Example 1 (162 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.58 (2H, s), 6.50 (1H, d, J=15.7 Hz), 7.38-7.56 (6H, m), 7.61 (2x1H, d, J=8 Hz), 7.73 (2x1H, d, J=7 Hz), 7.78-7.88 (2H, m), 7.96 (1H, s), 10.84 (1H, s);

MASS (ES+): m/e 370.

Example 8

Compound E8 was obtained from Compound (34) according to a manner similar to Example 1 (75 mg).

30 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.50 (2H, s), 6.48 (1H, d, J=16 Hz), 7.42-7.48 (3H, m), 7.56-7.71 (4H, m), 7.97 (1H, d, J=2 Hz);

MASS (ESI): m/z 372 (M+1).

Example 9

35 Compound E9 was obtained from Compound (35) according to manners similar to Preparation 9 and Example 1 (152 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.63 (3H, s), 4.58 (2H, s), 6.49 (1H,

d, J=16 Hz), 7.46 (1H, d, J=16 Hz), 7.50 (2H, d, J=8 Hz), 7.61 (2H, d, J=8 Hz), 7.85-7.93 (4H, m), 8.04-8.10 (3H, m);

MASS (ESI): m/z 412 (M+1).

Example 10

5 Compound E10 was obtained from Compound (36) according to manners similar to Preparation 9 and Example 1 (138 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.58 (2H, s), 6.50 (1H, d, J=16 Hz), 7.17-7.21 (1H, m), 7.46 (1H, d, J=16 Hz), 7.51 (2H, d, J=8 Hz), 7.59-7.65 (4H, m), 7.80-7.83 (2H, m), 7.95 (1H, s);

10 MASS (ESI): m/z 376 (M+1).

Example 11

Compound E11 was obtained from Compound (37) according to manners similar to Preparation 9 and Example 1 (120 mg).

15 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.58 (2H, s), 6.49 (1H, d, J=16 Hz), 7.46 (1H, d, J=16 Hz), 7.51 (2H, d, J=8 Hz), 7.61 (2H, d, J=8 Hz), 7.64-7.73 (2H, m), 7.79 (1H, d, J=8 Hz), 7.90 (1H, dd, J=2, 8 Hz), 8.00-8.05 (2H, m);

MASS (ESI): m/z 376 (M+1).

Example 12

20 Compound E12 was obtained from Compound (39) according to a manner similar to Example 1 (142 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.42 (2H, s), 6.46 (1H, d, J=16 Hz), 7.41-7.47 (3H, m), 7.56 (2H, d, J=8 Hz), 7.73 (1H, d, J=8 Hz), 7.80 (1H, d, J=8 Hz), 8.22 (1H, s);

25 MASS (ESI): m/z 317 (M-1).

Example 13

Compound E13 was obtained from Compound (40) according to manners similar to Preparation 9 and Example 1 (710 mg).

30 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.55 (2H, s), 6.50 (1H, d, J=16 Hz), 7.36-7.44 (1H, m), 7.45 (1H, d, J=16 Hz), 7.50 (2H, d, J=8 Hz), 7.59 (2H, d, J=8 Hz), 7.66 (1H, dd, J=2, 8 Hz), 7.77-7.83 (1H, m);

MASS (ESI): m/z 312 (M+1).

Example 14

35 Compound E14 was obtained from Compound (42) according to a manner similar to Example 1 (504 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.51 (2H, s), 6.48 (1H, d, J=16 Hz),

7.45 (1H, d, J=16 Hz), 7.47 (2H, d, J=8 Hz), 7.52 (1H, dd, J=2, 8 Hz), 7.59 (2H, d, J=8 Hz), 7.76 (1H, d, J=8 Hz), 7.86 (1H, J=2 Hz);
MASS (ESI): m/z 328 (M+1).

Example 15

5 Compound E15 was obtained from Compound (44) according to a manner similar to Example 1 (14 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.79-2.85 (4H, m), 3.11-3.22 (4H, m), 4.52 (2H, s), 6.48 (1H, d, J=16 Hz), 7.15-7.67 (8H, m);

MASS (ESI): m/z 392 (M+1).

10 Example 16

Compound E16 was obtained from Compound (46) according to a manner similar to Example 1 (45 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.15-3.21 (4H, m), 3.75-3.81 (4H, m), 4.52 (2H, s), 6.49 (1H, d, J=16 Hz), 7.10 (1H, d, J=2 Hz), 7.29 (1H,
15 dd, J=2, 8 Hz), 7.42-7.50 (3H, m), 7.57-7.64 (3H, m);

MASS (ESI): m/z 379 (M+1).

Example 17

Compound E17 was obtained from Compound (48) according to a manner similar to Example 1 (27 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.60-1.70 (2H, m), 1.83-2.00 (4H, m), 3.39-3.54 (4H, m), 4.54 (2H, s), 6.49 (1H, d, J=16 Hz), 7.40-7.86 (8H, m);

MASS (ESI): m/z 377 (M+1).

Example 18

25 A mixture of Compound (39), sodium azide (485 mg) and triethylamine hydrochloride (1.54 g) in N,N-dimethylformamide (7.5 mL) was heated at 130°C for 6 hours. After cooling, the reaction mixture was partitioned between ethyl acetate (20 mL) and water (40 mL). The organic layer was washed with brine, dried over MgSO₄,
30 filtered, and evaporated in vacuo. The crude product was triturated with ethyl acetate-water. The product was treated according to a manner similar to Example 1 to give Compound E18 (18 mg).

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.30 (2H, s), 6.44 (1H, d, J=16 Hz), 7.40 (2H, d, J=8 Hz), 7.44 (1H, d, J=16 Hz), 7.55 (2H, d, J=8 Hz), 7.72 (1H, d, J=8 Hz), 7.90 (1H, d, J=8 Hz), 8.21 (1H, s);

MASS (ESI): m/z 362 (M+1).

Example 19

Compound E19 was obtained from Compound (50) according to a manner similar to Example 1 (101 mg).

5 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.91 (3H, s), 4.55 (2H, s), 6.49 (1H, d, J=16 Hz), 7.46 (1H, d, J=16 Hz), 7.50 (2H, d, J=8 Hz), 7.60 (2H, d, J=8 Hz), 7.84 (1H, d, J=8 Hz), 8.05 (1H, d, J=8 Hz), 8.27 (1H, s);

MASS (ESI): m/z 352 (M+1).

10 Example 20

Compound E20 was obtained from Compound (51) according to a manner similar to Example 1 (72 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.55 (2H, s), 6.49 (1H, d, J=16 Hz), 7.45 (1H, d, J=16 Hz), 7.49 (2H, d, J=8 Hz), 7.60 (2H, d, J=8 Hz),
15 7.81 (1H, d, J=8 Hz), 8.04 (1H, dd, J=2, 8 Hz), 8.25 (1H, s);

MASS (ESI): m/z 338 (M+1).

Example 21

Compound E21 was obtained from Compound (53) according to a manner similar to Example 1 (55 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 2.68 (3H, s), 4.56 (2H, s), 6.49 (1H, d, J=16 Hz), 7.45 (1H, d, J=16 Hz), 7.50 (2H, d, J=8 Hz), 7.59 (2H, d, J=8 Hz), 7.83 (1H, d, J=8 Hz), 8.06 (1H, dd, J=2, 8 Hz), 8.28 (1H, s);

MASS (ESI): m/z 336 (M+1).

25 Example 22

Compound E22 was obtained from Compound (55) according to a manner similar to Example 1 (279 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.48 (2H, s), 6.47 (1H, d, J=16 Hz), 7.33-7.49 (4H, m), 7.57 (2H, d, J=8 Hz), 7.65-7.72 (4H, m);

30 MASS (ESI): m/z 372 (M+1).

Example 23

Compound E23 was obtained from Compound (57) according to a manner similar to Example 1 (50 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.25 (3H, s), 1.27 (3H, s), 3.04-3.16
35 (1H, m), 4.55 (2H, s), 6.49 (1H, d, J=16 Hz), 7.42-7.52 (4H, m), 7.56-7.62 (3H, m), 7.68 (1H, d, J=8 Hz);

MASS (ESI): m/z 336 (M+1).

Example 24

Compound E24 was obtained from Compound (59) according to manners similar to Preparation 9 and Example 1 (249 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.42 (2H, s), 5.35 (2H, s), 6.45 (1H, d, $J=16$ Hz), 7.10-7.61 (13H, m);

MASS (ESI): m/z 400 (M+1).

Example 25

10 Compound E25 was obtained from Compound (62) according to a manner similar to Example 1 (417 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.58 (2H, s), 6.50 (1H, d, $J=15.7$ Hz), 7.38-7.56 (6H, m), 7.61 (2x1H, d, $J=8$ Hz), 7.73 (2x1H, d, $J=7$ Hz), 7.78-7.88 (2H, m), 7.96 (1H, s), 10.84 (1H, s);

MASS (ES+): m/e 370.

15 Example 26

Compound E26 was obtained from Compound (72) according to a manner similar to Example 1 (207 mg).

20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.57 (2H, s), 6.50 (1H, d, $J=15.7$ Hz), 7.35 (2x1H, dd, $J=8.8$, 8.8 Hz), 7.46 (1H, d, $J=15.7$ Hz), 7.50 (2x1H, d, $J=8$ Hz), 7.61 (2x1H, d, $J=8$ Hz), 7.74-7.86 (4H, m), 7.94 (1H, s);

MASS (ES+): m/e 388.

Example 27

25 Compound E27 was obtained from Compound (75) according to a manner similar to Example 1 (123 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.58 (2H, s), 6.51 (1H, d, $J=15.8$ Hz), 7.38-7.59 (7H, m), 7.65-7.87 (5H, m), 7.96 (1H, s), 10.86 (1H, br);

MASS (ES+): m/e 370.

Example 28

30 Compound E28 was obtained from Compound (80) according to a manner similar to Example 1 (103 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.28 (3H, s), 4.59 (2H, s), 6.49 (1H, d, $J=16$ Hz), 7.42-7.65 (5H, m), 7.87-7.92 (2H, m), 7.98-8.10 (5H, m);

35 MASS (ES+): m/e 447.

Example 29

Compound E29 was obtained from Compound (83) according to a manner similar to Example 1 (200 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.49 (2H, s), 6.49 (1H, d, J=16 Hz), 7.44 (1H, d, J=16 Hz), 7.47 (2x1H, d, J=8 Hz), 7.57 (2x1H, d, J=8 Hz), 7.57 (1H, dd, J=8, 5 Hz), 8.34 (1H, d, J=8 Hz), 8.58 (1H, d, J=5 Hz);

MASS (ES+): m/e 295.

Example 30

Compound E30 was obtained from Compound (90) according to a manner similar to Example 1 (175 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.52 (2H, s), 6.50 (1H, d, J=16 Hz), 7.23 (1H, br-d, J=8.5 Hz), 7.41-7.53 (4H, m), 7.59 (2x1H, d, J=8 Hz), 7.69 (1H, d, J=8.5 Hz);

MASS (ES+): m/e 309.

Example 31

Compound E31 was obtained from Compound (93) according to a manner similar to Example 1 (14 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 0.92 (3H, t, J=7.5 Hz), 1.63 (2H, tq, J=7.5, 7.5 Hz), 2.34 (2H, t, J=7.5 Hz), 4.54 (2H, s), 6.49 (1H, d, J=15.7 Hz), 7.41-7.51 (3H, m), 7.55 (1H, dd, J=9, 2 Hz), 7.60 (2x1H, d, J=8.5 Hz), 7.69 (1H, d, J=9 Hz), 8.31 (1H, d, J=2 Hz);

MASS (ES+): m/e 379.

Example 32

Compound E32 was obtained from Compound (97) according to a manner similar to Example 1 (73 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.45 (2H, s), 6.45 (1H, d, J=16 Hz), 7.43 (1H, d, J=16 Hz), 7.45 (2x1H, d, J=8 Hz), 7.50 (2x1H, dd, J=8, 8 Hz), 7.54 (2x1H, d, J=8 Hz), 8.11 (1H, d, J=8 Hz), 8.21 (1H, d, J=8 Hz);

MASS (ES+): m/e 339.

Example 33

Compound E33 was obtained from Compound (100) according to a manner similar to Example 1 (15 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.43 (2H, s), 6.48 (1H, d, J=16 Hz), 7.39-7.60 (5H, m), 8.08 (1H, d, J=6.5 Hz), 8.55 (1H, d, J=6.5 Hz), 9.35 (1H, s);

MASS (ES+): m/e 295.

Example 34

Compound E34 was obtained from Compound (103) according to a manner similar to Example 1 (240 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.22 (3H, s), 3.40 (3H, s), 4.08 (2H, s), 6.44 (1H, d, $J=15.8$ Hz), 7.32 (2x1H, d, $J=8$ Hz), 7.42 (1H, d, $J=15.8$ Hz), 7.51 (2x1H, d, $J=8$ Hz);

MASS (ES+): m/e 356.

Example 35

10 Compound E35 was obtained from Compound (105) according to a manner similar to Example 1 (160 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.42 (2H, s), 6.47 (1H, d, $J=16$ Hz), 7.44 (1H, d, $J=16$ Hz), 7.44 (2x1H, d, $J=8$ Hz), 7.56 (2x1H, d, $J=8$ Hz), 7.80 (1H, d, $J=9$ Hz), 8.19 (1H, dd, $J=9, 2.2$ Hz), 8.50 (1H, d, $J=2.2$ Hz);

15 MASS (ES+): m/e 339.

Example 36

Compound E36 was obtained from Compound (112) according to a manner similar to Example 1 (375 mg).

20 $^1\text{H-NMR}$ (300 MHz, CDCl_3 , δ): 5.66 (2H, s), 6.38 (1H, d, $J=15.7$ Hz), 7.19 (2x1H, d, $J=8.8$ Hz), 7.43 (1H, d, $J=15.7$ Hz), 7.48-7.56 (2H, m), 7.58 (2x1H, d, $J=8.8$ Hz), 7.76-7.84 (2H, m), 10.75 (1H, br-s);

MASS (ES+): m/e 310.

Example 37

25 Compound E37 was obtained from Compound (115) according to a manner similar to Example 1 (40 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.40-1.94 (8H, m), 3.85 (1H, m), 4.54 (2H, s), 6.51 (1H, d, $J=15.8$ Hz), 7.28-7.80 (8H, m);

MASS (ES+): m/e 377.

30 Example 38

Compound E38 was obtained from Compound (122) according to a manner similar to Example 1 (1.19 g).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 5.67 (2H, s), 6.53 (1H, d, $J=16$ Hz), 7.14 (1H, m), 7.27 (1H, m), 7.35 (1H, s), 7.41 (1H, dd, $J=8, 8$ Hz), 7.46 (1H, d, $J=16$ Hz), 7.47-7.79 (2H, m), 7.77-7.87 (2H, m), 10.90 (1H, br);

MASS (ES+): m/e 310.

Example 39

Compound E39 was obtained from Compound (125) according to a manner similar to Example 1 (110 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.05 (2x3H, t, J=7 Hz), 3.52 (4H, m), 4.51 (2H, s), 6.49 (1H, d, J=15.8 Hz), 7.41-7.53 (5H, m), 7.56-7.64 (3H, m);

MASS (ES+): m/e 365.

Example 40

10 Compound E40 was obtained from Compound (132) according to a manner similar to Example 1 (1472 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.84 (3H, s), 5.62 (2H, s), 6.43 (1H, d, J=15.8 Hz), 7.14 (1H, d, J=8 Hz), 7.21 (1H, d, J=8 Hz), 7.28 (1H, s), 7.42 (1H, d, J=15.8 Hz), 7.50-7.58 (2H, m), 7.78-7.86 (2H, m);

15 MASS (ES+): m/e 340.

Example 41

Compound E41 was obtained from Compound (134) according to manners similar to Preparation 9 and Example 1 (115 mg).

20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.02 (2x3H, s), 4.52 (2H, s), 6.50 (1H, d, J=16 Hz), 6.90-7.70 (8H, m);

MASS (ES+) m/e 337.

Example 42

Compound E42 was obtained from Compound (141) according to a manner similar to Example 1 (450 mg).

25 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.82 (3H, s), 5.61 (2H, s), 6.40 (1H, d, J=15.8 Hz), 7.11 (1H, d, J=8.8 Hz), 7.28 (1H, dd, J=8.8, 1.7 Hz), 7.40 (1H, d, J=15.8 Hz), 7.43 (1H, d, J=1.7 Hz), 7.50-7.58 (2H, m), 7.78-7.86 (2H, m);

MASS (ES+): m/e 340.

30 Example 43

Compound E43 was obtained from Compound (144) according to a manner similar to Example 1 (160.8 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.45 (2H, s), 6.16 (2H, s), 6.47 (1H, d, J=15.8 Hz), 7.43 (1H, d, J=8.1 Hz), 7.46 (1H, d, J=16.0 Hz), 7.59 (2H, d, J=8.1 Hz);

MASS (ES+): m/e 338 (M+1).

Example 44

Compound E44 was obtained from Compound (147) according to a manner similar to Example 1 (160.8 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.00 (3H, s), 4.42 (2H, s), 6.44 (1H, d, J=16.2 Hz), 7.04 (1H, d, J=8.0 Hz), 7.27 (1H, d, J=8.4 Hz), 7.41 (1H, t, J=8.2 Hz), 7.42 (2H, d, J=8.0 Hz), 7.45 (1H, d, J=16.0 Hz), 7.57 (2H, d, J=8.0 Hz);

MASS (ES⁺): m/e 324 (M+1).

Example 45

10 Compound E45 was obtained from Compound (151) according to a manner similar to Example 1 (481.2 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.87 (3H, d, J=7.4 Hz), 4.85 (1H, q, J=7.4 Hz), 6.49 (1H, d, J=16.1 Hz), 7.43 (1H, d, J=15.7 Hz), 7.49-7.52 (2H, m), 7.51 (2H, d, J=8.5 Hz), 7.58 (2H, d, J=8.5 Hz), 7.74-7.77 (2H, m);

MASS (ES⁺): m/e 308 (M+1).

Example 46

Compound E46 was obtained from Compound (155) according to a manner similar to Example 1 (576.3 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 1.88 (3H, d, J=7.3 Hz), 4.83 (1H, q, J=7.4 Hz), 6.50 (1H, d, J=15.7 Hz), 7.43 (1H, d, J=15.4 Hz), 7.48-7.51 (2H, m), 7.53 (2H, d, J=8.2 Hz), 7.58 (2H, d, J=8.4 Hz), 7.74-7.77 (2H, m);

MASS (ES⁺): m/e 308 (M+1).

25 Example 47

Compound E47 was obtained from Compound (158) according to a manner similar to Example 1 (274.9 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.36 (3H, t, J=7.0 Hz), 4.08 (2H, q, J=7.0 Hz), 4.58 (2H, s), 6.49 (1H, d, J=15.7 Hz), 7.05 (2H, d, J=8.8 Hz), 7.46 (1H, d, J=17.0 Hz), 7.50 (2H, d, J=8.4 Hz), 7.61 (2H, d, J=8.4 Hz), 7.66 (2H, d, J=8.8 Hz), 7.77 (1H, dd, J=8.8, 1.5 Hz), 7.81 (1H, d, J=8.4 Hz), 7.90 (1H, s);

MASS (ES⁺): m/e 414 (M+1).

Example 48

35 Compound E48 was obtained from Compound (161) according to a manner similar to Example 1 (231.1 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.60 (2H, s), 6.49 (1H, d, J=15.8 Hz), 7.46 (1H, d, J=16.1 Hz), 7.52 (2H, d, J=8.4 Hz), 7.60 (2H, d, J=8.4 Hz), 7.93 (2H, s), 8.19 (1H, s), 8.66 (1H, d, J=9.0 Hz), 8.81 (1H, dd, J=5.5, 1.5 Hz), 9.21 (1H, d, J=1.8 Hz);

5 MASS (ES+): m/e 371 (M+1).

Example 49

Compound E49 was obtained from Compound (164) according to a manner similar to Example 1 (130.9 mg).

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.54 (2H, s), 6.48 (1H, d, J=15.8 Hz), 7.46 (1H, d, J=15.8 Hz), 7.47 (2H, d, J=8.1 Hz), 7.58 (2H, t, J=7.0 Hz), 7.60 (2H, d, J=8.1 Hz), 7.71 (1H, t, J=7.7 Hz), 7.76 (2H, d, J=7.4 Hz), 7.83 (1H, dd, J=8.8, 1.5 Hz), 7.87 (2H, d, J=8.4 Hz), 8.01 (1H, s);

MASS (ES+): m/e 398 (M+1).

15 Example 50

Compound E50 was obtained from Compound (166) according to a manner similar to Example 1 (69.3 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.51 (2H, s), 5.89 (1H, s), 6.46 (1H, d, J=45.8 Hz), 7.20 (1H, t, J=7.0 Hz), 7.30 (2H, t, J=7.3 Hz), 7.39 (2H, d, J=7.3 Hz), 7.46 (4H, m), 7.59 (2H, d, J=8.4 Hz), 7.65 (1H, d, J=8.4 Hz), 7.74 (1H, s);

MASS (ES+): m/e 398 (M+1).

Example 51

Compound E51 was obtained from Compound (169) according to a manner similar to Example 1 (103.9 mg).

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 3.00 (6H, s), 4.57 (2H, s), 6.48 (1H, d, J=15.8 Hz), 7.47 (1H, d, J=15.0 Hz), 7.50 (1H, d, J=8.1 Hz), 7.61 (2H, d, J=8.4 Hz), 7.65 (2H, d, J=9.1 Hz), 7.79 (2H, s), 7.87 (1H, s);

30 MASS (ES+): m/e 413 (M+1).

Example 52

Compound E52 was obtained from Compound (172) according to a manner similar to Example 1 (203.9 mg).

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.46 (2H, s), 6.46 (1H, d, J=16.1 Hz), 7.44 (2H, d, J=8.5 Hz), 7.45 (1H, d, J=16.0 Hz), 7.57 (2H, d, J=8.5 Hz), 7.69 (1H, d, J=8.5 Hz), 7.85 (1H, d, J=7.8 Hz), 8.03 (1H, s);

MASS (ES+): m/e 361 (M+1).

Example 53

Compound E53 was obtained from Compound (175) according to a manner similar to Example 1 (419.6 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.35 (2H, s), 6.46 (1H, d, J=15.7 Hz), 7.31-7.40 (2H, m); 7.42 (2H, d, J=8.0 Hz), 7.43 (1H, d, J=16.0 Hz), 7.56 (2H, d, J=8.0 Hz);

MASS (ES+): m/e 330 (M+1).

Example 54

10 Compound E54 was obtained from Compound (178) according to a manner similar to Example 1 (90.7 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 0.94 (6H, d, J=6.6 Hz), 1.64 (2H, dt, J=6.6, 6.6 Hz), 1.80 (1H, hept, J=6.6 Hz), 4.07 (2H, t, J=6.6 Hz), 4.51 (2H, s), 6.48 (1H, d, J=16.2 Hz), 7.11 (1H, dd, J=2.1, 9.0 Hz),
15 7.21 (1H, d, J=2.1 Hz), 7.44 (2H, d, J=8.4 Hz), 7.45 (1H, d, J=16.2 Hz), 7.59 (2H, d, J=8.5 Hz), 7.63 (1H, d, J=9.0 Hz);

MASS (ES+): m/e 380 (M+1).

Example 55

20 Compound E55 was obtained from Compound (181) according to a manner similar to Example 1 (215.8 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.29 (6H, d, J=6.1 Hz), 4.51 (2H, s), 4.70 (1H, hept, J=6.1 Hz), 6.48 (1H, d, J=16.0 Hz), 7.09 (1H, dd, J=2.2, 8.9 Hz), 7.20 (1H, d, J=2.2 Hz), 7.45 (1H, d, J=16.0 Hz), 7.47 (2H, d, J=8.5 Hz), 7.59 (2H, d, J=8.1 Hz), 7.63 (1H, d, J=8.9
25 Hz);

MASS (ES+): m/e 352 (M+1).

Example 56

Compound E56 was obtained from Compound (184) according to a manner similar to Example 1 (387.8 mg).

30 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.52 (2H, s), 6.48 (1H, d, J=16.1 Hz), 7.04 (2H, d, J=7.7 Hz), 7.18 (1H, t, J=7.7 Hz), 7.22 (1H, dd, J=8.8, 2.2 Hz), 7.30 (1H, d, J=2.2 Hz), 7.41 (2H, t, J=8.0 Hz), 7.46 (1H, d, J=16.0 Hz), 7.47 (2H, d, J=8.1 Hz), 7.57 (2H, d, J=7.7 Hz), 7.77 (1H, d, J=9.2 Hz);

35 MASS (ES+): m/e 386 (M+1).

Example 57

Compound E57 was obtained from Compound (187) according to a manner similar to Example 1 (78.2 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.85 (3H, s), 4.52 (2H, s), 6.48 (1H, d, $J=15.7$ Hz), 7.12 (1H, dd, $J=8.8, 2.2$ Hz), 7.20 (1H, d, $J=2.2$ Hz),
5 7.45 (1H, d, $J=15.0$ Hz), 7.46 (2H, d, $J=8.4$ Hz), 7.60 (2H, d, $J=8.4$ Hz), 7.65 (1H, d, $J=9.2$ Hz);
MASS (ES+): m/e 324 (M+1).

Example 58

To a solution of Compound (277) (2.46 g) in MeOH (12 mL) was
10 added hydrogen chloride methanol reagent 10 (15 mL, Tokyo Kasei Kogyo Co., Ltd.), and the mixture was stirred at ambient temperature for 2 hrs. The reaction mixture was diluted with isopropyl ether (50 mL) and the precipitate was collected by filtration. The obtained pale yellow powder was crystallized with
15 a mixed solvent (EtOH: $\text{H}_2\text{O} = 6: 4$, 100 mL) to give Compound E58 (1.23 g) as colorless powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.39 (2H, s), 6.46 (1H, d, $J=15.7$ Hz),
7.43 (3H, m), 7.50 (1H, t, $J=6.6$ Hz), 7.52 (2H, t, $J=7.0$ Hz), 7.58
(2H, d, $J=8.0$ Hz), 7.82 (2H, d, $J=7.4$ Hz), 8.06 (1H, s);
20 MASS (ES+): m/e 320 (M+1).

Example 59

Compound E59 was obtained from Compound (278) according to a manner similar to Example 58 (274 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.81 (3H, s), 4.33 (2H, s), 6.49 (1H,
25 d, $J=16.1$ Hz), 7.36-7.65 (8H, m), 7.74-7.85 (2H, m);
MASS (ES+): m/z 378 (M+1, free).

Example 60

Compound E60 was obtained from Compound (188) according to a manner similar to Example 58 (1.36 g).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.80 (3H, s), 4.26 (2H, s), 6.45 (1H,
30 d, $J=15.8$ Hz), 7.38-7.48 (3H, m), 7.52-7.61 (4H, m), 7.82 (2H, d);
MASS (ES+): m/z 412 (M+1, free).

Example 61

Compound E61 was obtained from Compound (193) according to a
35 manner similar to Example 58 (68 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.32 (2H, s), 6.47 (1H, d, $J=16.1$ Hz),

7.39-7.61 (6H, m), 7.62-7.89 (4H, m);

MASS (ES+): m/z 363 (M+1, free).

Example 62

Compound E61 was obtained from Compound (194) according to a
5 manner similar to Example 58 (40 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.10 (2H, s), 6.44 (1H, d, J=15.8 Hz),
7.31-7.64 (8H, m), 7.76-7.86 (2H, m);

MASS (ES+): m/z 345 (M+1, free).

Example 63

10 Compound E63 was obtained from Compound (203) according to a
manner similar to Example 58 (212 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.40 (2H, s), 6.47 (1H, d, J=16 Hz),
7.34-7.50 (5H, m), 7.58 (2x1H, d, J=8 Hz), 7.87-7.96 (2H, m), 8.05
(1H, s), 10.81 (1H, s);

15 MASS (ES+): m/z 338.

Example 64

Compound E64 was obtained from Compound (189) according to a
manner similar to Example 58 (81 mg).

20 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.40 (2H, s), 6.47 (1H, d, J=15.2 Hz),
7.44 (1H, d, J=15.2 Hz), 7.45 (2H, d, J=8.4 Hz), 7.55-7.66 (4H, m,
J=8.4 Hz), 7.89 (2H, d, J=8.8 Hz), 8.11 (1H, s);

MASS (ES+): m/z 354 (M+1, free).

Example 65

25 Compound E65 was obtained from Compound (209) according to a
manner similar to Example 58 (145 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 2.52 (3H, s), 2.53 (3H, s), 4.27 (2H,
s), 6.47 (1H, d, J=16 Hz), 7.41 (2x1H, d, J=8 Hz), 7.44 (1H, d,
J=16 Hz), 7.56 (2x1H, d, J=8 Hz), 10.82 (1H, br-s);

MASS: not detected.

30 Example 66

Compound E66 was obtained from Compound (208) according to a
manner similar to Example 58 (364 mg).

35 ¹H-NMR (300 MHz, DMSO-d₆, δ): 2.25 (3H, s), 4.29 (2H, s), 6.48 (1H,
d, J=16 Hz), 7.39-7.50 (3H, m), 7.53-7.63 (4H, m), 7.71 (1H, m),
7.87 (2x1H, d, J=7.5 Hz), 10.81 (1H, br-s); MASS (ES+): m/z 362.

Example 67

Compound E67 was obtained from Compound (222) according to a manner similar to Example 58 (90 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 6.52 (1H, d, J=16.1 Hz), 7.40-7.57 (6H, m), 7.63 (2H, d, J=8.1 Hz), 7.86 (2H, d, J=7.0 Hz), 8.29 (1H, s),
5 9.47 (1H, s);

MASS (ES+): m/z 320 (M+1, free).

Example 68

Compound E68 was obtained from Compound (190) according to a manner similar to Example 58 (80 mg).

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.43 (2H, s), 6.49 (1H, d, J=16.1 Hz), 7.38-7.57 (7H, m), 7.64 (1H, s), 7.87 (2H, d, J=7.0 Hz), 8.09 (1H, s);

MASS (ES+): m/z 320 (M+1, free).

Example 69

15 Compound E69 was obtained from Compound (191) according to a manner similar to Example 58 (47 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.09 (2H, s), 6.47 (1H, d, J=16.1 Hz), 7.29-7.59 (8H, m), 7.70-7.77 (2H, m);

MASS (ES+): m/z 354 (M+1, free).

20 Example 70

Compound E70 was obtained from Compound (279) according to a manner similar to Example 58 (1.34 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.81 (3H, s), 4.38 (2H, s), 6.47 (1H, d, J=16.1 Hz), 7.08 (2H, d, J=8.8 Hz), 7.44 (2H, d, J=8.4 Hz), 7.45
25 (1H, d, J=15.1 Hz), 7.58 (2H, d, J=8.4 Hz), 7.77 (2H, d, J=8.8 Hz), 7.94 (1H, s);

MASS (ES+): m/e 350 (M+1).

Example 71

30 Compound E71 was obtained from Compound (280) according to a manner similar to Example 58 (482 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.83 (3H, s), 4.40 (2H, s), 6.47 (1H, d, J=15.8 Hz), 7.00 (1H, m), 7.39-7.47 (6H, m), 7.58 (2H, d, J=8.1 Hz), 8.09 (1H, s);

MASS (ES+): m/e 350 (M+1).

35 Example 72

Compound E72 was obtained from Compound (281) according to a

manner similar to Example 58 (498 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.93 (3H, s), 4.40 (2H, s), 6.46 (1H, d, J=15.7 Hz), 7.11 (1H, t, J=8.1 Hz), 7.21 (1H, d, J=8.1 Hz), 7.43 (2+1+1H, d, J=8.1 Hz), 7.58 (2H, d, J=8.1 Hz), 7.80 (1H, dd, J=8.1, 1.6 Hz), 7.88 (1H, s);
5 MASS (ES+): m/z 350 (M+1).

Example 73

Compound E73 was obtained from Compound (282) according to a manner similar to Example 58 (94.8 mg).

10 ¹H-NMR (300 MHz, DMSO-d₆, δ): 4.13 (2H, s), 6.44 (1H, d, J=15.8 Hz), 7.37 (2H, d, J=8.4 Hz), 7.38 (2H, t, J=8.1 Hz), 7.43 (1H, d, J=15.4 Hz), 7.50 (2H, t, J=8.1 Hz), 7.54 (2H, d, J=8.4 Hz), 7.74 (2H, d, J=8.1 Hz);
MASS (ES+): m/e 398 (M+1).

15 Example 74

Compound E74 was obtained from Compound (283) according to a manner similar to Example 58 (76 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.03 (2H, s), 6.43 (1H, d, J=16.1 Hz), 7.32 (1H, t, J=7.4 Hz), 7.34 (2H, d, J=7.4 Hz), 7.43 (1H, d, J=15.4 Hz), 7.47 (2H, t, J=7.7 Hz), 7.52 (2H, d, J=8.4 Hz), 7.70 (2H, d, J=8.1 Hz);
20 MASS (ES+): m/e 354 (M+1).

Example 75

Compound E75 was obtained from Compound (285) according to a manner similar to Example 58 (74.3 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 1.64 (2H, br.), 1.72 (4H, br.), 2.88 (2H, br.), 3.27 (2H, br.), 4.35 (2H, s), 4.43 (2H, s), 6.47 (1H, d, J=15.7 Hz), 7.45 (1H, d, J=15.7 Hz), 7.49 (2+1H, m), 7.56 (2+2H, m), 7.67 (2H, d, J=7.4 Hz);
30 MASS (ES+): m/e 417 (M+1).

Example 76

Compound E76 was obtained from Compound (286) according to a manner similar to Example 58 (76.8 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 3.15 (4H, br.), 3.81 (4H, br.), 4.36 (2H, s), 4.46 (2H, s), 6.48 (1H, d, J=15.8 Hz), 7.45 (1H, d, J=16.1 Hz), 7.49 (2+1H, m), 7.55 (2+2H, m), 7.71 (2H, d, J=7.3 Hz);

MASS (ES+): m/e 419 (M+1).

Example 77

Compound E77 was obtained from Compound (287) according to a manner similar to Example 58 (76 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.79 (3H, s), 4.05 (2H, s), 6.44 (1H, d, J=15.8 Hz), 7.05 (2H, d, J=8.1 Hz), 7.34 (2H, d, J=8.4 Hz), 7.43 (1H, d, J=16.1 Hz), 7.52 (2H, d, J=8.1 Hz), 7.64 (2H, d, J=8.1 Hz);
MASS (ES+): m/e 384 (M+1).

Example 78

10 Compound E78 was obtained from Compound (223) according to a manner similar to Example 58 (280 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.69 (3H, s), 5.43 (2H, s), 6.49 (1H, d, J=15.8 Hz), 7.44 (3H, m), 7.47 (1H, d, J=15.8 Hz), 7.53 (2H, t, J=7.3 Hz), 7.62 (2H, d, J=8.4 Hz), 7.81 (2H, d, J=7.0 Hz), 8.16 (1H,
15 s);

MASS (ES+): m/e 334 (M+1).

Example 79

Compound E79 was obtained from Compound (224) according to a manner similar to Example 58 (432.4 mg).

20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.31 (3H, s), 5.53 (2H, s), 6.50 (1H, d, J=15.8 Hz), 7.40 (2H, d, J=8.1 Hz), 7.47 (1H, d, J=15.8 Hz), 7.63 (2H, d, J=8.1 Hz), 7.64 (4H, s), 9.36 (1H, s);

MASS (ES+): m/z 368 (M+1).

Example 80

25 Compound E80 was obtained from Compound (288) according to a manner similar to Example 58 (176 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 5.47 (2H, s), 6.52 (1H, d, J=15.8 Hz), 7.41-7.53 (6H, m), 7.60 (1H, d, J=6.2 Hz), 7.68 (1H, s), 7.84 (2H, d, J=7.0 Hz), 8.26 (1H, s), 9.34 (1H, s);

30 MASS (ES+): m/e 319 (M+1).

Example 81

Compound E81 was obtained from Compound (289) according to a manner similar to Example 58 (268 mg).

35 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.33 (3H, s), 5.54 (2H, s), 6.53 (1H, d, J=16.1 Hz), 7.369-7.54 (4H, m), 7.58 (1H, s), 7.63 (2H, d, J=8.4 Hz), 7.68 (2H, d, J=8.4 Hz), 9.42 (1H, s);

MASS (ES+): m/e 368 (M+1).

Example 82

Compound E82 was obtained from Compound (321) according to a manner similar to Example 58 (220 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 5.50 (2H, s), 6.48 (1H, d, J=16.1 Hz), 7.15 (2H, d, J=8.1 Hz), 7.41 (1H, d, J=16.1 Hz), 7.53 (2H, d, J=8.1 Hz), 7.59-7.78 (5H, m), 7.88-7.97 (2H, m);

MASS (ES+): m/z 320 (M+1, free).

Example 83

10 Compound E83 was obtained from Compound (323) according to a manner similar to Example 58 (60 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.73 (3H, d, J=4.8 Hz), 4.28 (2H, s), 6.46 (1H, d, J=15.8 Hz), 7.35-7.61 (7H, m), 7.70-7.80 (2H, m), 8.24-8.37 (1H, m);

15 MASS (ES+): m/z 377 (M+1, free).

Example 84

Compound E84 was obtained from Compound (290) according to a manner similar to Example 58 (22.2 mg).

20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.78 (3H, s), 4.22 (2H, s), 6.44 (1H, d, J=15.8 Hz), 7.39 (2H, d, J=8.1 Hz), 7.43 (1H, d, J=15.8 Hz), 7.48 (3H, m), 7.54 (2H, d, J=8.4 Hz), 7.76 (2H, m);

MASS (ES+): m/e 378 (M+1).

Example 85

25 Compound E85 was obtained from Compound (291) according to a manner similar to Example 58 (13.5 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.29 (9H, s), 4.32 (2H, s), 6.46 (1H, d, J=15.8 Hz), 7.29 (1H, s), 7.39 (2H, d, J=8.4 Hz), 7.44 (1H, d, J=16.1 Hz), 7.57 (2H, d, J=8.1 Hz);

MASS (ES+): m/z 300 (M+1).

30 Example 86

Compound E86 was obtained from Compound (198) according to a manner similar to Example 58 (491 mg).

35 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.23 (3H, t, J=7.0 Hz), 4.24 (2H, s), 4.26 (2H, q, J=7.0 Hz), 6.44 (1H, d, J=15.8 Hz), 7.41 (2H, d, J=8.0 Hz), 7.43 (1H, d, J=15.4 Hz), 7.48 (3H, m), 7.55 (2H, d, J=8.1 Hz), 7.75 (2H, m);

MASS (ES+): m/e 392 (M+1).

Example 87

Compound E87 was obtained from Compound (332) according to a manner similar to Example 58 (86.7 mg).

5 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.71 (3H, s), 2.98 (3H, s), 4.32 (2H, s), 6.46 (1H, d, $J=16.1$ Hz), 7.41-7.60 (10H, m);

MASS: Not Detected.

Example 88

10 Compound E88 was obtained from Compound (325) according to a manner similar to Example 58 (35.8 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.09 (2H, br.), 1.49 (4H, br.), 3.12 (4H, br.), 4.34 (2H, s), 6.47 (1H, d, $J=15.8$ Hz), 7.42-7.60 (10H, m);

MASS (ES+): m/e 431 (M+1).

15 Example 89

Compound E89 was obtained from Compound (326) according to a manner similar to Example 58 (40.5 mg).

20 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.11 (6H, d, $J=6.6$ Hz), 4.02 (1H, m), 4.23 (2H, s), 6.44 (1H, d, $J=15.8$ Hz), 7.40 (2H, d, $J=7.7$ Hz), 7.48 (4H, m); 7.56 (2H, d, $J=8.1$ Hz), 7.73 (2H, d, $J=8.1$ Hz);

MASS (ES+): m/e 405 (M+1).

Example 90

Compound E90 was obtained from Compound (284) according to a manner similar to Example 58 (11.1 mg).

25 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.71 (6H, s), 4.21 (2H, s), 4.38 (2H, s), 6.44 (1H, d, $J=16.1$ Hz), 7.40 (3H, m), 7.48 (2H, m), 7.55 (5H, m);

MASS (ES+): m/e 377 (M+1).

Example 91

30 Compound E91 was obtained from Compound (323) according to a manner similar to Example 58 (40.1 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.27 (2H, s), 6.45 (1H, d, $J=15.8$ Hz), 7.41-7.58 (8H, m), 7.78 (2H, m);

MASS: Not Detected.

35 Example 92

Compound E92 was obtained from Compound (327) according to a

manner similar to Example 58 (14.8 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.37 (2H, s), 4.60 (2H, s), 6.47 (1H, d, $J=15.0$ Hz), 7.42-7.60 (8H, m), 7.67 (2H, d, $J=7.0$ Hz);

MASS (ES+): m/e 350 (M+1).

5 Example 93

Compound E93 was obtained from Compound (329) according to a manner similar to Example 58 (145 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.34 (3H, s), 4.37 (2H, s), 4.51 (2H, s), 6.46 (1H, d, $J=15.8$ Hz), 7.43-7.65 (10H, m);

10 MASS (ES+): m/e 364 (M+1).

Example 94

Compound E94 was obtained from Compound (330) according to a manner similar to Example 58 (42.4 mg).

15 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 2.17 (3H, s), 4.41 (2H, s), 6.46 (1H, d, $J=16.1$ Hz), 7.37-7.44 (2H, m), 7.45 (1H, d, $J=15.8$ Hz), 7.52-7.60 (6H, m), 7.71 (1H, m);

MASS (ES+): m/z 362 (M+1).

Example 95

20 Compound E95 was obtained from Compound (225) according to a manner similar to Example 58 (176 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 5.47 (2H, s), 6.53 (1H, d, $J=16$ Hz), 7.44 (1H, d, $J=16$ Hz), 7.45 (2x1H, d, $J=8.5$ Hz), 7.60 (2x1H, d, $J=8.5$ Hz), 7.72 (1H, dd, $J=1.7, 1.5$ Hz), 7.81 (1H, dd, $J=1.7, 1.3$ Hz), 9.34 (1H, dd, $J=1.5, 1.3$ Hz), 10.89 (1H, br-s), 14.73 (1H, br-s);

25 MASS (ES+): m/e 244.

Example 96

Compound E96 was obtained from Compound (210) according to a manner similar to Example 58 (21 mg).

30 $^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.96-2.03 (4H, m), 3.24-3.31 (4H, m), 4.49 (2H, s), 6.49 (1H, d, $J=16$ Hz), 6.60 (1H, s), 6.83 (1H, dd, $J=2, 8$ Hz), 7.42-7.62 (6H, m);

MASS (ES+): m/z 363 (M+1).

Example 97

35 Compound E97 was obtained from Compound (328) according to a manner similar to Example 58 (7.9 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 4.55 (2H, s), 4.66 (2H, s), 6.48 (1H, d, $J=16$ Hz), 7.43-7.51 (4H, m), 7.60 (2H, d, $J=8$ Hz), 7.71 (2H, d, $J=8$ Hz);

MASS (ES+): m/z 324 (M+1).

5 Example 98

Compound E98 was obtained from Compound (333) according to a manner similar to Example 58 (72 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.77-1.92 (4H, m), 3.33-3.54 (4H, m), 4.57 (2H, s), 6.50 (1H, d, $J=16$ Hz), 7.45 (1H, d, $J=16$ Hz), 7.50 (2H, d, $J=8$ Hz), 7.60 (2H, d, $J=8$ Hz), 7.64 (1H, dd, $J=2, 8$ Hz), 7.80 (1H, d, $J=8$ Hz), 7.88 (1H, s);

MASS (ES+): m/z 391 (M+1).

Example 99

Compound E99 was obtained from Compound (334) according to a manner similar to Example 58 (12 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.24 (6H, t, $J=7$ Hz), 2.93-3.09 (4H, m), 4.25 (2H, s), 4.32-4.39 (2H, m), 6.78 (1H, d, $J=16$ Hz), 7.38-7.65 (7H, m), 7.82 (1H, s);

MASS (ES+): m/z 379 (M+1).

20 Example 100

Compound E100 was obtained from Compound (316) according to a manner similar to Example 58 (40 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 3.71 (3H, s), 4.10 (2H, s), 5.11 (2H, s), 6.41 (1H, d, $J=16$ Hz), 6.86 (2H, d, $J=8$ Hz), 7.09 (2H, d, $J=8$ Hz), 7.16 (2H, d, $J=8$ Hz), 7.36-7.52 (3H, m), 7.80 (1H, s);

MASS (ES-): m/z 430 (M-1).

Example 101

Compound E101 was obtained from Compound (317) according to a manner similar to Example 58 (164 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6 , δ): 1.65-1.79 (6H, m), 1.84-1.93 (6H, m), 2.00-2.07 (3H, m), 4.32 (2H, s), 6.47 (1H, d, $J=16$ Hz), 7.25 (1H, s), 7.38-7.48 (3H, m), 7.56 (2H, d, $J=8$ Hz);

MASS (ES+): m/z 378 (M+1).

Example 102

35 Compound E102 was obtained from Compound (338) according to a manner similar to Example 58 (35 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.31 (3H, t, $J=7$ Hz), 4.28 (2H, s), 4.33 (2H, q, $J=7$ Hz), 6.47 (1H, d, $J=16$ Hz), 7.37 (2H, d, $J=8$ Hz), 7.43 (1H, d, $J=16$ Hz), 7.55 (2H, d, $J=8$ Hz), 8.23 (1H, s);
MASS (ES+): m/z 316 (M+1).

5 Example 103

Compound E103 was obtained from Compound (312) according to a manner similar to Example 58 (138 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.36 (2H, m), 3.30 (2H, t, $J=7.4$ Hz), 4.15 (2H, t, $J=5.8$ Hz), 6.30 (1H, d, $J=16.1$ Hz), 6.79 (2H, d, $J=8.8$ Hz), 7.38 (1H, d, $J=15.8$ Hz), 7.45 (2H, d, $J=8.8$ Hz), 7.54 (2H, m), 7.78 (2H, m);
MASS (ES+): m/e 338 (M+1).

Example 104

Compound E104 was obtained from Compound (313) according to a manner similar to Example 58 (174 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.84 (2H, m), 2.03 (2H, m), 3.21 (2H, t, $J=7.4$ Hz), 4.07 (2H, t, $J=6.4$ Hz), 6.31 (1H, d, $J=15.4$ Hz), 6.96 (2H, d, $J=8.8$ Hz), 7.40 (1H, d, $J=15.8$ Hz), 7.49 (2H, d, $J=8.1$ Hz), 7.53 (2H, m), 7.78 (2H, m);
MASS (ES+): m/e 352 (M+1).

Example 105

Compound E105 was obtained from Compound (314) according to a manner similar to Example 58 (153 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.21 (2H, quint., $J=7.3$ Hz), 2.74 (2H, t, $J=7.3$ Hz), 3.14 (2H, t, $J=7.7$ Hz), 6.42 (1H, d, $J=16.1$ Hz), 7.29 (2H, d, $J=8.0$ Hz), 7.41 (1H, d, $J=15.8$ Hz), 7.48 (2H, d, $J=7.7$ Hz), 7.52 (2H, m), 7.76 (2H, m);
MASS (ES+): m/e 322 (M+1).

Example 106

30 A suspension of Compound E116 (1.23 g, described later in Example 116) and benzenesulfonic acid (732 mg) in 80% aqueous ethanol (20 mL) was dissolved at 90°C. The insoluble materials were removed by filtration and the filtrate was evaporated in vacuo to give a crude solid. This solid was recrystallized from 80%
35 aqueous ethanol to give Compound E106 as a pale yellow crystal (820 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.53 (2H, s), 6.47 (1H, d, $J=16$ Hz), 7.28-7.36 (3H, m), 7.41-7.56 (5H, m), 7.57-7.66 (4H, m), 7.72-7.81 (2H, m), 10.78 (1H, s);

MASS (ES+): m/z 294.

5 Example 107

A suspension of Compound E116 (1.80 g, described later in Example 116) and (1R)-(-)-10-camphorsulfonic acid (855 mg) in 80% aqueous ethanol (20 mL) was dissolved at 90°C. The insoluble materials were removed by filtration, and the filtrate was
10 evaporated in vacuo to give a crude solid. This solid was recrystallized from 80% aqueous ethanol to give Compound E107 as a white crystal (1.30 g).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 0.74 (3H, s), 1.05 (3H, s), 1.20-1.35 (2H, m), 1.74-1.96 (3H, m), 2.24 (1H, m), 2.39 (1H, d, $J=14.7$ Hz),
15 2.69 (1H, m), 2.89 (1H, d, $J=14.7$ Hz), 4.55 (2H, s), 6.48 (1H, d, $J=15.7$ Hz), 7.42-7.58 (5H, m), 7.61 (2x1H, d, $J=8$ Hz), 7.74-7.82 (2H, m), 10.79 (1H, s);

MASS (ES+): m/z 294.

Example 108

20 To a stirred suspension of Compound E116 (1.0 g, described later in Example 116) in ethanol (20 mL) was added 4-methylbenzenesulfonic acid monohydrate (713 mg) at ambient temperature. The resulting suspension was stirred at 70°C for 1 hr, during which time water (0.1 mL) was added to the mixture in order
25 to dissolve some insoluble materials. The mixture was allowed to cool to ambient temperature and the stirring was continued for additional 1 hr. The precipitate was filtered, washed with ethanol (5 mL, 2 times) and dried to afford crude Compound E108 (1.58 g) as a pale tan solid. The Compound E110 was used without further
30 purification.

Example 109

A suspension of crude Compound E108 (salt, 710 mg) in 80% aqueous acetonitrile (20 mL) was heated at 90°C for 15 min. The insoluble materials were removed by filtration and the resulting
35 filtrate was allowed to cool to ambient temperature. The solution was stirred at ambient temperature for 1 hr and then in an ice bath

for 0.5 hr. The precipitate formed was filtered, washed with acetonitrile (5 mL, 2 times) and dried to afford Compound E111 (370 mg) as a pale tan solid.

m.p. 228.5–230.5°C

5 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.29 (3H, s), 4.54 (2H, s), 6.47 (1H, d, $J=15.8$ Hz), 7.11 (2H, d, $J=7.7$ Hz), 7.42–7.57 (7H, m), 7.61 (2H, d, $J=8.4$ Hz), 7.72–7.82 (2H, m);

MASS (ES+): m/z 294 (M+1, free).

Example 110

10 To a solution of Compound (292) (42 mg) in MeOH (1 mL) was added hydrogen chloride methanol reagent 10 (0.8 mL, manufactured by Tokyo Kasei Kogyo Co., Ltd.), and the mixture was stirred at ambient temperature for 2 hrs. To the reaction mixture was added saturated aqueous NaHCO_3 solution and extracted with AcOEt. The
15 organic phase was sequentially washed with saturated aqueous NaHCO_3 solution, water and brine, and the solvent was removed in vacuo. The obtained colorless solid was triturated with MeCN to give Compound E110 (4.8 mg) as colorless powder.

20 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 1.28 (4H, m), 1.69 (4H, m), 1.90 (2H, m), 2.37 (1H, m), 3.91 (2H, s), 6.41 (1H, d, $J=15.8$ Hz), 6.62 (1H, s), 7.26 (2H, d, $J=8.1$ Hz), 7.41 (1H, d, $J=15.8$ Hz), 7.47 (2H, d, $J=7.7$ Hz);

MASS (ES+): m/z 326 (M+1).

Example 111

25 Compound E111 was obtained from Compound (284) according to a manner similar to Example 110 (289 mg).

30 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.16 (6H, s), 3.44 (2H, s), 4.00 (2H, s), 6.41 (1H, d, $J=16.5$ Hz), 7.18 (1H, t, $J=7.0$ Hz), 7.32 (2H, d, $J=7.3$ Hz), 7.33 (2H, t, $J=7.3$ Hz), 7.41 (1H, d, $J=15.8$ Hz), 7.50 (2H, d, $J=7.7$ Hz), 7.70 (2H, d, $J=7.3$ Hz);

MASS (ES+): m/z 377 (M+1).

Example 112

35 Compound E112 was obtained from Compound (197) according to a manner similar to Example 58 (22.5 mg).

m.p. 235–239°C

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.34 (2H, s), 6.47 (1H, d, $J=15.8$ Hz),

7.37 (2H, d, J=8.1 Hz), 7.44 (1H, d, J=15.8 Hz), 7.57 (2H, d, J=8.1 Hz), 7.60 (2H, s);

MASS (ES+) m/z 243 (M+1, free).

Example 113

5 To a stirred solution of [(1S,4R)-7,7-dimethyl-2-oxobicyclo[2.2.1]hept-1-yl]methanesulfonic acid (832 mg, described later in Example 116) in ethanol (10 mL) was added Compound E116 (1.0 g, described later in Example 118) at ambient temperature. The resulting suspension was heated at 60°C for 1 hr and then
10 allowed to cool to ambient temperature. The suspension was further stirred at ambient temperature for 1 hr. The precipitate was filtered, washed with ethanol (2 mL, 3 times) and dried to afford Compound E113 (1.73 g) as a pale tan solid. The Compound E113 was used in the following Example 114 without further purification.

15 Example 114

A suspension of crude Compound E113 (salt, 700 mg) in 90% aqueous acetonitrile (20 mL) was heated at 90°C for 5 min. The insoluble materials were removed by filtration and the resulting filtrate was heated at 90°C again. The solution was allowed to
20 cool to ambient temperature and stirred at the same temperature for 1 hr. The precipitate was filtered, washed with acetonitrile (2 mL, 2 times) and dried to afford Compound E114 (522 mg) as a pale tan solid.

m.p. 185.5-194.5°C

25 ¹H-NMR (300 MHz, DMSO-d₆, δ): 0.74 (3H, s), 1.05 (3H, s), 1.20-1.35 (2H, m), 1.79 (1H, d, J=18.0 Hz), 1.80-1.90 (1H, m), 1.93 (1H, t, J=4.2 Hz), 2.23 (1H, dt, J=18.0, 4.2 Hz), 2.39 (1H, d, J=14.7 Hz), 2.61-2.76 (1H, m), 2.89 (1H, d, J=14.7 Hz), 4.54 (2H, s), 6.48 (1H, d, J=16.1 Hz), 7.41-7.57 (5H, m), 7.61 (2H, d, J=8.1 Hz), 7.73-7.82
30 (2H, m), 10.79 (1H, br.s);

MASS (ES+): m/z 294 (M+1, free).

Example 115

To a mixture of Compound (322) (as a mixture of two regioisomers, 310 mg in total) in methanol (1 mL) was added 10%
35 hydrogen chloride in methanol (3 mL) at ambient temperature for 20 min. The mixture was concentrated in vacuo, and the residue was

purified by preparative high performance liquid chromatography
(column; Mightysil RP-18 GP 250-20 Kanto Chemical Co., Inc.)
eluting with a gradient solvent system from acetonitrile in water
(10: 90 v/v) to acetonitrile in water (50: 50 v/v). The residue

5 was lyophilized to give a mixture of Compound E115 as a mixture of
two regioisomers (85 mg in total) as a pale red viscous oil.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): (for a mixture of two regioisomers)
2.17 (1.2H, s), 2.25 (1.8H, s), 5.39 (1.2H, s), 5.47 (0.8H, s),
6.52 (1H, d, $J=15.8$ Hz), 7.18-7.37 (1.2H, m), 7.39-7.54 (2.8H, m),
10 7.60 (1.2H, d, $J=8.1$ Hz), 7.75 (0.8H, d, $J=8.1$ Hz), 9.20 (0.6H, s),
9.25 (0.4H, s);

MASS (ES+): (for a mixture of two regioisomers) m/z 258 (M+1).

Example 116

To a suspension of Compound E1 (990 mg) in ethanol (10 mL)
15 was added 1N sodium hydroxide solution (3.0 mL), and the mixture
was stirred at 70°C for 2 hrs. The resulting precipitate was
collected by filtration and washed with ethanol and H_2O to give
Compound E116 (810 mg) as a white powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.19 (2H, s), 6.41 (1H, d, $J=16$ Hz),
20 7.07-7.16 (2H, m), 7.33-7.55 (7H, m).

Example 117

To a suspension of Compound E116 (147 mg) in ethanol (5 mL)
was added 1N sodium hydroxide solution (0.501 mL), and the mixture
was stirred at room temperature for 1 hour. The resulting mixture
25 was evaporated in vacuo and triturated with isopropyl ether to give
Compound E117 (162 mg) as a pale green powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.13 (2H, s), 6.32 (1H, d, $J=16$ Hz),
6.89 (1H, d, $J=16$ Hz), 7.07-7.14 (2H, m), 7.24 (2H, d, $J=8$ Hz),
7.36 (2H, d, $J=8$ Hz), 7.43-7.50 (2H, m).

30 Example 118

To a suspension of Compound E116 (147 mg) in ethanol (5 mL)
was added 1N methanesulfonic acid solution (0.501 mL), and the
mixture was stirred at room temperature for 2 hrs. The resulting
precipitate was collected by filtration and washed with ethanol to
35 give Compound E118 (91 mg) as a white powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 2.32 (3H, s), 4.54 (2H, s), 6.47 (1H,

d, J=16 Hz), 7.46 (2H, d, J=8 Hz), 7.48-7.56 (3H, m), 7.61 (2H, d, J=8 Hz), 7.74-7.81 (2H, m).

Example 119

To a suspension of Compound E116 (147 mg) in ethanol (5 mL) was added 1N sulfuric acid (1.0 mL), and the mixture was stirred at 70°C for 2 hours. The resulting precipitate was collected by filtration and washed with ethanol to give Compound E119 (177 mg) as a white powder.

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.44 (2H, s), 6.46 (1H, d, J=16 Hz), 7.39-7.49 (5H, m), 7.59 (2H, d, J=8 Hz), 7.67-7.71 (2H, m).

Example 120

To a solution of Compound (6) (1.0 g), 3-(benzyloxy)-1,2-benzenediamine (817 mg) and 1-hydroxybenzotriazole (567 mg) in N,N-dimethylformamide (19 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (804 mg). After stirring for 6 hrs, saturated aqueous NaHCO₃ (20 mL) and water (80 mL) were added to the reaction mixture. The resulting precipitate was collected by filtration and washed with water. To a mixture of this product in acetic acid (7 mL) was added a solution of hydrogen chloride in acetic acid (13 mL). After stirring at 100°C for 1 hour, the resulting mixture was cooled to 4°C and diluted with ethyl acetate (40 mL). The resulting precipitate was collected by filtration and washed with ethyl acetate. To a solution of this product, O-tetrahydro-2H-pyran-2-ylhydroxylamine (1.5 eq.) and 1-hydroxybenzotriazole (1.5 eq.) in N,N-dimethylformamide (11 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.5 eq.) at 4°C. The mixture was warmed to ambient temperature and stirred for 8 hrs. The reaction mixture was added saturated NaHCO₃ (11 mL) and water (44 mL). A resulting precipitate was filtered and the filtrate was extracted with ethyl acetate. The organic layer was washed with H₂O and brine, dried over MgSO₄, filtered and evaporated in vacuo. The crude product was treated with trifluoroacetic acid to give Compound E120 (52 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.43 (2H, s), 6.46 (1H, d, J=16 Hz), 6.86 (1H, d, J=8 Hz), 7.12 (1H, d, J=8 Hz), 7.23-7.30 (1H, m), 7.39-7.62 (5H, m);

MASS (ES+): m/z 310 (M+1).

Example 121

To a solution of Compound (6) (1.0 g), 4-(benzyloxy)-1,2-benzenediamine (817 mg) and 1-hydroxybenzotriazole (567 mg) in N,N-dimethylformamide (19 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (804 mg). After stirring for 6 hours, the reaction mixture were added saturated NaHCO₃ (20 mL) and water (80 mL). The resulting precipitate was collected by filtration and washed with water. To a solution of this product in acetic acid (7 mL) was added hydrogen chloride in acetic acid (13 mL). After stirring at 100°C for 1 hour, the resulting mixture was cooled to 4°C and diluted with ethyl acetate (40 mL). A resulting precipitate was collected by filtration and washed with ethyl acetate. This product, O-tetrahydro-2H-pyran-2-ylhydroxylamine (1.5 eq.) and 1-hydroxybenzotriazole (1.5 eq.) were dissolved in N,N-dimethylformamide (9 mL), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.5 eq.) was added thereto at 4°C. The mixture was warmed to ambient temperature and stirred for 8 hours. To the reaction mixture were added saturated NaHCO₃ (10 mL) and water (40 mL). The resulting precipitate was collected by filtration and washed with water. The crude product was treated with hydrogen chloride in methanol to give Compound E121 (35 mg).

¹H-NMR (300 MHz, DMSO-d₆, δ): 4.52 (2H, s), 5.22 (2H, s), 6.49 (1H, d, J=16 Hz), 7.21 (1H, dd, J=2, 8 Hz), 7.30 (1H, d, J=2 Hz), 7.34-7.50 (8H, m), 7.59 (2H, d, J=8 Hz), 7.67 (1H, d, J=8 Hz);

MASS (ES+): m/z 400 (M+1).

Example 122

To a solution of Compound (51) (70 mg) in N,N-dimethylformamide (1.5 mL) was added N,N'-carbonyldiimidazole (40 mg). The mixture was stirred for 30 min, and methanesulfonamide (24 mg) and 1,8-diazabicyclo[5.4.0]-7-undecene (0.037 mL) were added thereto. The reaction mixture was stirred for 8 hrs at 80°C. The resulting mixture was acidified with 1 N hydrochloric acid at 0°C, and water (7.5 mL) was added thereto. The resulting precipitate was collected by filtration and washed with water. The crude product was treated with hydrogen chloride in methanol to

give Compound E122 (7.9 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.40 (3H, s), 4.46 (2H, s), 6.47 (1H, d, $J=16$ Hz), 7.41-7.49 (3H, m), 7.58 (2H, d, $J=8$ Hz), 7.76 (1H, d, $J=8$ Hz), 7.95 (1H, d, $J=8$ Hz), 8.29 (1H, s);

5 MASS (ES+): m/z 415 (M+1).

Example 123

To a suspension of Compound E116 (500 mg) in ethanol (5 mL) was added maleic acid (198 mg), and the mixture was stirred at 70°C for 2 hrs. The resulting precipitate was collected by filtration and washed with ethanol and water. The solid was recrystallized
10 from 50% aqueous EtOH to give Compound E123 (374 mg) as a pale yellow powder.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 4.33 (2H, s), 6.17 (2H, s), 6.44 (1H, d, $J=16$ Hz), 7.24-7.32 (2H, m), 7.38-7.48 (3H, m), 7.53-7.62 (4H,
15 m).

Example 124

Compound E124 was obtained from Compound (260) according to manners similar to Preparations 337 and 208 and Example 58 (7 mg).

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6 , δ): 3.93 (2H, s), 4.18 (2H, s), 6.45 (1H, d, $J=16$ Hz), 7.13 (1H, s), 7.23-7.38 (7H, m), 7.43 (1H, d, $J=16$ Hz),
20 7.54 (2H, d, $J=8$ Hz);

MASS (ES+): m/z 334 (M+1).

The compounds obtained in the above-mentioned Preparations
25 are shown in the following Table 2 (including Tables 2-1 to 2-44)
and the above-mentioned Examples are shown in the following Table 3
(including Tables 3-1 to 3-17).

Table 2

Table 2-1









Compound (1)	Compound (2)
	
	
	
	

Table 2-2









Compound (9)	Compound (10)
	
Compound (11)	Compound (12)
	
Compound (13)	Compound (14)
	
Compound (15)	Compound (16)
	

Table 2-3









Compound (17)	Compound (18)
	
Compound (19)	Compound (20)
	
Compound (21)	Compound (22)
	
Compound (23)	Compound (24)
	

Table 2-4









Compound (25)	Compound (26)
	
Compound (27)	Compound (28)
	
Compound (29)	Compound (30)
	
Compound (31)	Compound (32)
	

Table 2-5









Compound (33)	Compound (34)
 <p>•HCl</p>	 <p>•HCl</p>
Compound (35)	Compound (36)
 <p>•HCl</p>	
Compound (37)	Compound (38)
 <p>•HCl</p>	 <p>•HCl</p>
Compound (39)	Compound (40)
 <p>•HCl</p>	 <p>•HCl</p>

Table 2-6









Compound (41)	Compound (42)
 <p>• HCl</p>	
Compound (43)	Compound (44)
 <p>• x HCl</p>	
Compound (45)	Compound (46)
 <p>• HCl</p>	
Compound (47)	Compound (48)
 <p>• HCl</p>	

Table 2-7









Compound (49)	Compound (50)
	
Compound (51)	Compound (52)
	
Compound (53)	Compound (54)
	
Compound (55)	Compound (56)
	

Table 2-8









Compound (57)	Compound (58)
	
Compound (59)	Compound (60)
	
Compound (61)	Compound (62)
	
Compound (63)	Compound (64)
	

Table 2-9









Compound (65)	Compound (66)
	
Compound (67)	Compound (68)
	
Compound (69)	Compound (70)
	
Compound (71)	Compound (72)
	

Table 2-10









Compound (73)	Compound (74)
	
Compound (75)	Compound (76)
	
Compound (77)	Compound (78)
	
Compound (79)	Compound (80)
	

Table 2-11









Compound (81)	Compound (82)
	
Compound (83)	Compound (84)
	
Compound (85)	Compound (86)
 <p style="text-align: center;">• HCl</p>	
Compound (87)	Compound (88)
	

Table 2-12









Compound (89)	Compound (90)
	
Compound (91)	Compound (92)
	
Compound (93)	Compound (94)
	
Compound (95)	Compound (96)
	

Table 2-13









Compound (97)	Compound (98)
	
Compound (99)	Compound (100)
	
Compound (101)	Compound (102)
	
Compound (103)	Compound (104)
	

Table 2-14









Compound (105)	Compound (106)
	
Compound (107)	Compound (108)
	
Compound (109)	Compound (110)
	
Compound (111)	Compound (112)
	

Table 2-15









Compound (113)	Compound (114)
	
Compound (115)	Compound (116)
	
Compound (117)	Compound (118)
	
Compound (119)	Compound (120)
	

Table 2-16









Compound (121)	Compound (122)
	
Compound (123)	Compound (124)
	
Compound (125)	Compound (126)
	
Compound (127)	Compound (128)
	

Table 2-17









Compound (129)	Compound (130)
	
Compound (131)	Compound (132)
	
Compound (133)	Compound (134)
	
Compound (135)	Compound (136)
	

Table 2-18









Compound (137)	Compound (138)
	
Compound (139)	Compound (140)
	
Compound (141)	Compound (142)
	
Compound (143)	Compound (144)
	

Table 2-19









Compound (145)	Compound (146)
	
Compound (147)	Compound (148)
	
Compound (149)	Compound (150)
	
Compound (151)	Compound (152)
	

Table 2-20









Compound (153)	Compound (154)
	
Compound (155)	Compound (156)
	
Compound (157)	Compound (158)
	
Compound (159)	Compound (160)
	

Table 2-21









Compound (161)	Compound (162)
	
Compound (163)	Compound (164)
	
Compound (165)	Compound (166)
	
Compound (167)	Compound (168)
	

Table 2-22









Compound (169)	Compound (170)
	
Compound (171)	Compound (172)
	
Compound (173)	Compound (174)
	
Compound (175)	Compound (176)
	

Table 2-23









Compound (177)	Compound (178)
 <p>• HCl</p>	
Compound (179)	Compound (180)
	 <p>• HCl</p>
Compound (181)	Compound (182)
	
Compound (183)	Compound (184)
 <p>• HCl</p>	

Table 2-24




Compound (185)	Compound (186)
	
Compound (187)	
	

Table 2-25









Compound (188)	Compound (189)
	
Compound (190)	Compound (191)
	
Compound (192)	Compound (193)
	
Compound (194)	Compound (195)
	

Table 2-26









Compound (196)	Compound (197)
	
	
	
	

Table 2-27









Compound (204)	Compound (205)
	
Compound (206)	Compound (207)
	
Compound (208)	Compound (209)
	
Compound (210)	Compound (211)
	

Table 2-28










Compound (212)	Compound (213)
	
Compound (214)	Compound (215)
	
Compound (216)	Compound (217)
	
Compound (218)	Compound (219)
 <p>(Main product)</p>  <p>(By-product)</p>	

Table 2-29









Compound (220)	Compound (221)
	
Compound (222)	Compound (223)
	
Compound (224)	Compound (225)
	
Compound (226)	Compound (227)
	

Table 2-30









Compound (228)	Compound (229)
	
Compound (230)	Compound (231)
	
Compound (232)	Compound (233)
	
Compound (234)	Compound (235)
	

Table 2-31









Compound (236)	Compound (237)
	
Compound (238)	Compound (239)
	
Compound (240)	Compound (241)
	
Compound (242)	Compound (243)
	

Table 2-32









Compound (244)	Compound (245)
	
Compound (246)	Compound (247)
	
Compound (248)	Compound (249)
	
Compound (250)	Compound (251)
	

Table 2-33









Compound (252)	Compound (253)
	
Compound (254)	Compound (255)
	
Compound (256)	Compound (257)
	
Compound (258)	Compound (259)
	

Table 2-34









Compound (260)	Compound (261)
	
Compound (262)	Compound (263)
	
Compound (264)	Compound (265)
	
Compound (266)	Compound (267)
	

Table 2-35









Compound (268)	Compound (269)
<p style="text-align: center;">TFAOH TFAOH</p> 	
	
	
	

Table 2-36









Compound (276)	Compound (277)
	
	
	
	

Table 2-37









Compound (284)	Compound (285)
	
Compound (286)	Compound (287)
	
Compound (288)	Compound (289)
	
Compound (290)	Compound (291)
	

Table 2-38









Compound (292)	Compound (293)
	
Compound (294)	Compound (295)
	
Compound (296)	Compound (297)
	
Compound (298)	Compound (299)
	

Table 2-39









Compound (300)	Compound (301)
	
Compound (302)	Compound (303)
	
Compound (304)	Compound (305)
	
Compound (306)	Compound (307)
	

Table 2-40









Compound (308)	Compound (309)
	
Compound (310)	Compound (311)
	
Compound (312)	Compound (313)
	
Compound (314)	Compound (315)
	

Table 2-41









Compound (316)	Compound (317)
	
Compound (318)	Compound (319)
	
Compound (320)	Compound (321)
	
Compound (322)	Compound (323)
 <p>(Main product)</p> <p>(By-product)</p>	

Table 2-42









Compound (324)	Compound (325)
	
Compound (326)	Compound (327)
	
Compound (328)	Compound (329)
	
Compound (330)	Compound (331)
	

Table 2-43









Compound (332)	Compound (333)
	
Compound (334)	Compound (335)
	
Compound (336)	Compound (337)
	
Compound (338)	Compound (339)
	

Table 2-44




Table 3

Table 3-1

Compound E1	Compound E2
<p>• HCl</p>	<p>• HCl</p>
Compound E3	Compound E4
<p>• HCl</p>	<p>• HCl</p>
Compound E5	Compound E6
<p>• HCl</p>	<p>• HCl</p>
Compound E7	
<p>• HCl</p>	

Table 3-2









Compound E8	Compound E9
	
Compound E10	Compound E11
	
Compound E12	Compound E13
	
Compound E14	Compound E15
	

Table 3-3









Compound E16	Compound E17
	
Compound E18	Compound E19
	
Compound E20	Compound E21
	
Compound E22	Compound E23
	

Table 3-4









Compound E24	Compound E25
 <p>• TFA</p>	 <p>• HCl</p>
 <p>• HCl</p>	 <p>• HCl</p>
 <p>• HCl</p>	 <p>• HCl</p>
 <p>• HCl • HCl</p>	 <p>• HCl</p>

Table 3-5









Compound E32	Compound E33
 <p style="text-align: center;">• HCl</p>	 <p style="text-align: center;">• HCl</p>
Compound E34	Compound E35
 <p style="text-align: center;">• HCl</p>	 <p style="text-align: center;">• HCl</p>
Compound E36	Compound E37
 <p style="text-align: center;">• HCl</p>	 <p style="text-align: center;">• HCl</p>
Compound E38	Compound E39
 <p style="text-align: center;">• HCl</p>	 <p style="text-align: center;">• HCl</p>

Table 3-6









Compound E40	Compound E41
 <p>• HCl</p>	 <p>• HCl</p>
Compound E42	Compound E43
 <p>• HCl</p>	 <p>• HCl</p>
Compound E44	Compound E45
 <p>• HCl</p>	 <p>• HCl</p>
Compound E46	Compound E47
 <p>• HCl</p>	 <p>• HCl</p>

Table 3-7









Compound E48	Compound E49
 <p>· HCl</p>	 <p>· HCl</p>
Compound E50	Compound E51
 <p>· HCl</p>	 <p>· HCl</p>
Compound E52	Compound E53
 <p>· HCl</p>	 <p>· HCl</p>
Compound E54	Compound E55
 <p>· HCl</p>	 <p>· HCl</p>

Table 3-8



Compound E56	Compound E57
<p data-bbox="641 367 722 399">• HCl</p> 	 <p data-bbox="1079 504 1161 535">• HCl</p>

Table 3-9









Compound E58	Compound E59
 <chem>O=C(O/N)C=Cc1ccc(cc1)CN2C=NC(C2)c3ccccc3</chem> HCl	 <chem>COC(=O)C1=NC2=CC=CC=C2N1C=C(O/N)C=Cc3ccc(cc3)C4=CC=CC=C4</chem> HCl
Compound E60	Compound E61
 <chem>COC(=O)C1=NC2=CC=C(Cl)C=C2N1C=C(O/N)C=Cc3ccc(cc3)C4=CC=CC=C4</chem> HCl	 <chem>Nc1c2ccccc2n1C=C(O/N)C=Cc3ccc(cc3)C4=CC=CC=C4</chem> HCl
Compound E62	Compound E63
 <chem>N#CC1=NC2=CC=CC=C2N1C=C(O/N)C=Cc3ccc(cc3)C4=CC=CC=C4</chem> HCl	 <chem>Fc1ccc(cc1)C2=NC3=CC=CC=C3N2C=C(O/N)C=Cc4ccc(cc4)C5=CC=CC=C5</chem> HCl
Compound E64	Compound E65
 <chem>Clc1ccc(cc1)C2=NC3=CC=CC=C3N2C=C(O/N)C=Cc4ccc(cc4)C5=CC=CC=C5</chem> HCl	 <chem>CC(=O)C1=NC2=CC=CC=C2N1C=C(O/N)C=Cc3ccc(cc3)C4=CC=CC=C4</chem> HCl

Table 3-10









Compound E66	Compound E67
	
	
	
	

Table 3-11









Compound E74	Compound E75
	
Compound E76	Compound E77
	
Compound E78	Compound E79
	
Compound E80	Compound E81
	

Table 3-12









Compound E82	Compound E83
	
	
	
	

Table 3-13









Compound E90	Compound E91
	
Compound E92	Compound E93
	
Compound E94	Compound E95
	
Compound E96	Compound E97
	

Table 3-14









Compound E98	Compound E99
	
Compound E100	Compound E101
	
Compound E102	Compound E103
	
Compound E104	Compound E105
	

Table 3-15









Compound E106	Compound E107
	
Compound E108	Compound E109
	
Compound E110	Compound E111
	
Compound E112	Compound E113
	

Table 3-16












Compound E114	Compound E115
	 <p>(Main product)</p> <p>+</p> <p>(By-product)</p>
Compound E116	Compound E117
	
Compound E118	Compound E119
 <p>• MsOH</p>	 <p>• H2SO4</p>
Compound E120	Compound E121
 <p>• TFA</p>	 <p>• HCl</p>

Table 3-17

Compound E122	Compound E123
	
Compound E124	
	


INDUSTRIAL APPLICABILITY

As mentioned above, the present invention can provide a novel compound having a potent inhibitory effect on the activity of histone deacetylase, and a pharmaceutical composition comprising the same. The compound is useful as an active ingredient of an immunosuppressant and an antitumor agent, and as an active ingredient of a therapeutic or prophylactic agent for diseases such as inflammatory disorders, diabetes, diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), organ transplant rejections, autoimmune diseases, protozoal infections, tumors, etc.

This application is based on the patent applications Nos. 2003900116 and 2003905406, both were filed in Australia, and the contents of which are incorporated hereinto by reference.

CLAIMS

1. A compound having the following formula (I):




wherein

- 5 R^1 is N-containing heterocyclic ring optionally substituted with one or more suitable substituent(s),
 R^2 is hydroxyamino,
 R^3 is hydrogen or a suitable substituent,
 L^1 is $-(\text{CH}_2)_n-$ (wherein n is an integer of 0 to 6) optionally
 10 substituted with one or more suitable substituent(s),
 wherein one or more methylene(s) may be replaced with suitable heteroatom(s), and
 L^2 is lower alkenylene,
 or a salt thereof.


15

2. The compound of claim 1, wherein

R^1 is N-containing heterocyclic ring represented by the following formula:



20



wherein

R⁴ is hydrogen or a group selected from the group consisting of

- 5 (1) lower alkyl optionally substituted with di(lower)alkylamino or hydroxy,
- (2) lower alkoxy,
- (3) aryl optionally substituted with the substituent selected from the group consisting of halogen, lower
- 10 alkanoyl, lower alkylsulfonyl, lower alkoxy and di(lower)alkylamino,
- (4) lower alkanoyl,
- (5) lower alkoxy-carbonyl,
- (6) arylcarbonyl,
- 15 (7) aryl(lower)alkoxy,
- (8) amino optionally mono- or di-substituted with substituent(s) selected from the group consisting of lower alkyl, lower alkanoyl and cycloalkyl,
- (9) halo(lower)alkyl,
- 20 (10) aryloxy,
- (11) aryl(lower)alkyl optionally substituted with hydroxy,
- (12) carboxyl,
- (13) nitro,
- 25 (14) cyano,
- (15) halogen,
- (16) heteroaryl,
- (17) non-aromatic heterocycle optionally substituted with lower alkyl,
- 30 (18) hydroxy,
- (19) (lower)alkylsulfonylcarbonyl and
- (20) non-aromatic heterocycle carbonyl,

R⁵ is hydrogen or a group selected from the group consisting

of lower alkyl and aryl(lower)alkyl, and
R⁶, R⁷ and R⁸ are each hydrogen or lower alkyl,
R⁹ is hydrogen or a group selected from the group consisting
of

- 5 (1) lower alkyl optionally substituted with
di(lower)alkylamino,
(2) aryl optionally substituted with lower alkoxy,
(3) (lower)alkoxy-carbonyl,
(4) cyano,
10 (5) carbamoyl optionally mono- or di-substituted with
(lower)alkyl,
(6) halogen,
(7) (lower)alkyl-carbonyl,
(8) arylcarbonyl and
15 (9) cyclo(lower)alkyl,

R¹⁰ is hydrogen or a group selected from the group consisting
of

- (1) (lower)alkylcarbamoyl,
(2) di(lower)alkylcarbamoyl,
20 (3) aryl optionally substituted with halogen,
(4) (lower)alkoxy-carbonyl,
(5) carboxy,
(6) non-aromatic heterocycle carbonyl,
(7) halogen,
25 (8) (lower)alkyl optionally substituted with hydroxy,
(lower)alkoxy, non-aromatic heterocycle, aryl,
di(lower)alkylamino or halogen and
(9) adamantyl,

R¹¹ is hydrogen or aryl(lower)alkyl in which the aryl portion
30 is substituted with lower alkoxy,

R¹² is hydrogen or a group selected from the group consisting
of lower alkyl and aryl optionally substituted with
halogen,

R¹³ is hydrogen or a group selected from the group consisting
35 of lower alkyl and aryl, and

R¹⁴ is hydrogen or lower alkyl,

R² is hydroxyamino,

R³ is hydrogen or lower alkoxy,


L¹ is -(CH₂)_n- (wherein n is 1 to 5) optionally substituted with one or more substituent(s) selected from lower alkyl(s) and aryl(lower)alkyl, and wherein one methylene may be replaced with an oxygen atom, and

L² is vinylene,

or a salt thereof.

10 3. The compound of claim 2, wherein

R¹ is N-containing condensed heterocyclic ring represented by the following formula:



wherein R⁴ and R⁵ are each as defined in claim 2.

15

4. The compound of claim 3, wherein

R⁴ and R⁵ are each hydrogen,

R² is hydroxyamino,

R³ is hydrogen,


20 L¹ is -CH₂-, and

L² is vinylene,

or a salt thereof.

5. The compound of claim 2, wherein

25 R¹ is N-containing heterocyclic ring represented by the following formula:



wherein R⁹, R¹⁰ and R¹¹ are each as defined in claim 2.

30

6. The compound of claim 5, wherein

R⁹ is hydrogen or aryl optionally substituted with lower alkoxy,

R¹⁰ is hydrogen or aryl optionally substituted with halogen, and

R¹¹ is hydrogen,

5 R² is hydroxyamino,

R³ is hydrogen,




L¹ is -CH₂-, and

L² is vinylene,


or a salt thereof.

10

7. A compound of the following formula




or



15 or a salt thereof.

8. A compound having the following formula (I'):



wherein

R¹ is N-containing condensed heterocyclic ring optionally substituted with one or more suitable substituent(s),

5 R² is hydroxyamino,

R³ is hydrogen or a suitable substituent,

L¹ is -(CH₂)_n- (wherein n is an integer of 0 to 6) optionally substituted with one or more suitable substituent(s), wherein one or more methylene(s) may be replaced with suitable heteroatom(s),


10 and

L² is lower alkenylene,

or a salt thereof.

9. The compound of claim 8, wherein

15 R¹ is N-containing condensed heterocyclic ring represented by the following formula:



20


wherein

R⁴ is hydrogen or a group selected from the group consisting of

(1) lower alkyl,

- (2) lower alkoxy,
(3) aryl optionally substituted with the substituent selected from the group consisting of halogen, lower alkanoyl, lower alkylsulfonyl, lower alkoxy and
5 di(lower)alkylamino,
(4) lower alkanoyl,
(5) lower alkoxy-carbonyl,
(6) arylcarbonyl,
(7) aryl(lower)alkoxy,
10 (8) amino optionally mono- or di-substituted with substituent(s) selected from the group consisting of lower alkyl, lower alkanoyl and cycloalkyl,
(9) halo(lower)alkyl,
(10) aryloxy,
15 (11) aryl(lower)alkyl optionally substituted with hydroxy,
(12) carboxyl,
(13) nitro,
(14) cyano,
20 (15) halogen,
(16) heteroaryl and
(17) non-aromatic heterocycle optionally substituted with lower alkyl,
R⁵ is hydrogen or a group selected from the group consisting
25 of lower alkyl and aryl(lower)alkyl, and
R⁶, R⁷ and R⁸ are each hydrogen or lower alkyl,
R² is hydroxyamino,
R³ is hydrogen or lower alkoxy,
L¹ is -(CH₂)_n- (wherein n is 1 or 2) optionally substituted with
30 one or more substituent(s) selected from lower alkyl(s) and aryl(lower)alkyl, and wherein one methylene may be replaced with an oxygen atom, and
L² is vinylene,
or a salt thereof.
35

10. A compound having the following formula (I''):



wherein

R¹ is N-containing condensed heterocyclic ring optionally substituted with one or more suitable substituent(s),

5 R² is hydroxyamino,

L¹ is -(CH₂)_n- (wherein n is an integer of 0 to 6) optionally substituted with one or more suitable substituent(s), and


L² is lower alkenylene,

or a salt thereof.

10

11. The compound of claim 10, wherein

R¹ is N-containing condensed heterocyclic ring represented by the following formula:



15

wherein

R⁴ is hydrogen or a group selected from the group consisting of lower alkyl and aryl, and

R⁵ is hydrogen or a group selected from the group consisting of lower alkyl and aryl(lower)alkyl,

20

R² is hydroxyamino,

L¹ is -(CH₂)_n- (wherein n is 1 or 2) optionally substituted with aryl(lower)alkyl, and

L² is vinylene,

25 or a salt thereof.

12. A histone deacetylase inhibitor comprising the compound of any of claims 1, 7, 8 and 10.

13. A pharmaceutical composition for treating or preventing inflammatory disorders, diabetes, diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), organ transplant rejections, autoimmune diseases, protozoal infections or tumors, which comprises the compound of any of claims 1, 7, 8 and 10.

14. A pharmaceutical composition containing the compound of any of claims 1, 7, 8 and 10 as an active ingredient, in association with a pharmaceutically acceptable, substantially non-toxic carrier or excipient.

15. The compound of any of claims 1, 7, 8 and 10 for use as a medicament.

16. A method for inhibiting histone deacetylase, comprising using the compound of any of claims 1, 7, 8 and 10.

17. Use of the compound of any of claims 1, 7, 8 and 10 for the manufacture of a medicament for inhibiting histone deacetylase.

18. A method for treating or preventing inflammatory disorders, diabetes, diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), organ transplant rejections, autoimmune diseases, protozoal infections or tumors, which comprises administering an effective amount of the compound of any of claims 1, 7, 8 and 10 to a human being or an animal.

19. Use of the compound of any of claims 1, 7, 8 and 10 for the manufacture of a medicament for treating or preventing inflammatory disorders, diabetes, diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), organ transplant rejections, autoimmune diseases, protozoal infections or tumors.

20. A commercial package comprising the pharmaceutical composition

of claim 13 and a written matter associated therewith, the written matter stating that the pharmaceutical composition may or should be used for treating or preventing inflammatory disorders, diabetes, diabetic complications, homozygous thalassemia, fibrosis, cirrhosis, acute promyelocytic leukaemia (APL), organ transplant rejections, 5 autoimmune diseases, protozoal infections or tumors.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/JP2004/000157

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D235/16 C07D471/04 C07D487/04 C07D235/12 C07D491/04
C07D401/04 C07D403/04 C07D233/64 C07D233/90 C07D233/68
A61K31/4184 A61K31/4164 A61K31/4178 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 03/087066 A (SK CHEMICALS CO LTD ; IN2GEN CO LTD (KR); KIM DAE-KEE (KR); KIM JAE-SU) 23 October 2003 (2003-10-23) examples 1-5,7,8,15 claims 1,4	1-20
P, X	WO 03/082288 A (ROMERO-MARTIN MARIA-ROSARIO ; FINN PAUL W (GB); RITCHIE JAMES (GB); PR) 9 October 2003 (2003-10-09) page 61 - page 72; compounds 6,7,12-15,28-41,49-51,54-58,65-70,75,76,82 -90 claim 67	1-20

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

7 June 2004

Date of mailing of the international search report

18/06/2004

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kollmannsberger, M

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/JP2004/000157

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/30872 A (MARUHA CORP ; MORI KENJI (JP); TAMAI TADAKAZU (JP); NISHIKAWA MASAZUMI) 18 April 2002 (2002-04-18) claims 25,72,136 page 170; example 117 -----	1-20
X	WO 01/38322 A (METHYLGENE INC) 31 May 2001 (2001-05-31) cited in the application claims 26,32,41 examples 31-96,99,104 -----	1-20
X	WO 02/22577 A (NOVARTIS ERFIND VERWALT GMBH ; NOVARTIS AG (CH); BAIR KENNETH WALTER () 21 March 2002 (2002-03-21) cited in the application examples 1-265 -----	1-20
X	WO 95/13264 A (TERUMO CORP ; ISOZAKI MASASHI (JP); KASUKAWA HIROAKI (JP); NAKAZAWA KE) 18 May 1995 (1995-05-18) page 18 - page 23; compounds 20-27 -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/JP2004/000157

Patent document cited in search report	A	Publication date	Patent family member(s)	Publication date
WO 03087066	A	23-10-2003	WO 03087066 A1	23-10-2003
WO 03082288	A	09-10-2003	WO 03082288 A1	09-10-2003
WO 0230872	A	18-04-2002	AU 9424601 A CA 2425923 A1 EP 1325904 A1 WO 0230872 A1 US 2004014816 A1	22-04-2002 11-04-2003 09-07-2003 18-04-2002 22-01-2004
WO 0138322	A	31-05-2001	AU 1876801 A CA 2391952 A1 EP 1233958 A1 WO 0138322 A1 JP 2003514904 T US 6541661 B1	04-06-2001 31-05-2001 28-08-2002 31-05-2001 22-04-2003 01-04-2003
WO 0222577	A	21-03-2002	AU 8212901 A BR 0113669 A CA 2420899 A1 CN 1450991 T CZ 20030573 A3 WO 0222577 A2 EP 1318980 A2 HU 0300880 A2 JP 2004509105 T NO 20030867 A SK 2422003 A3 US 2004024067 A1 US 2003018062 A1	26-03-2002 03-06-2003 21-03-2002 22-10-2003 18-06-2003 21-03-2002 18-06-2003 28-08-2003 25-03-2004 25-02-2003 11-09-2003 05-02-2004 23-01-2003
WO 9513264	A	18-05-1995	WO 9513264 A1 JP 7278086 A	18-05-1995 24-10-1995